

Review

Current Status of Antibiotic Resistance in the Six Major Foodborne Bacteria

Fabio Granados-Chinchilla^{1,2,*}

¹ Centro de Investigación en Enfermedades Tropicales (CIET), Facultad de Microbiología, Universidad de Costa Rica, Sede Rodrigo Facio, San Pedro Montes de Oca, San José 11501-2060, Costa Rica

² Escuela de Química, Facultad de Ciencias Básicas, Sede Rodrigo Facio, Universidad de Costa Rica, San Pedro Montes de Oca, San José 11501-2060, Costa Rica

* Corresponding author. E-mail: fabio.granados@ucr.ac.cr (F.G.-C.)

Received: 16 April 2026; Revised: 17 June 2026; Accepted: 25 August 2026; Available online: 14 September 2026

ABSTRACT: Bacterial pathogens still represent a serious public health issue. Special attention should be paid to bacteria that are usually transmitted through food. More relevantly, several isolates found in food exhibit (usually high) antibiotic resistance, which, in the case of human or animal infection, will reduce treatment efficacy. Hence, this review seeks to summarize and make available to researchers on the subject the most recent findings regarding antibiotic resistance profiles of common pathogens that are transferred into food. The bacterial pathogens responsible for most poisonings and illnesses include *Salmonella* spp., *Listeria* spp., *Campylobacter* spp., *Staphylococcus* spp., *E. coli* pathotypes, and *Cronobacter* spp. Therefore, we will focus this review on these pathogens. In this way, this writing will not only provide valuable information regarding which matrices are more susceptible to bacterial contamination, but in the end, the reader will be able to identify which antibiotics have been tested on these isolates and where the gaps are in the phenotypic determination of resistance to antibiotics, while highlighting several genotypic traits specific to some of these bacteria.

Keywords: Pathogens; Bacteria; Foodstuffs; Antibiotic; Resistance

1. Introduction

1.1. Foodborne Bacterial Pathogens

Foodborne pathogens have gained notoriety because, despite sustained quality practices and regulations regarding bacterial load in food production, foods are still found to be contaminated with bacteria. Pathogens are usually poor competitors and do not survive in the presence of existing food microflora [1] or after food processing or unit operations for food preservation (e.g., vacuum, pasteurization, nano/microfiltration) [2]. However, this ecological defense is not a guarantee of safety; food extended, erroneous manipulation, or bacterial survival mechanisms favor pathogen persistence. FDA food recalls are targeted actions to remove or correct food products that violate U.S. food safety laws. They are primarily driven by microbiological contamination, undeclared major allergens (such as milk or peanuts in mislabeled packages), and foreign objects. For example, of the 233 total food and beverage recalls from 2023, 70 (30.0%) were associated with possible or confirmed foodborne illness/bacterial contamination.

Understanding the relevance of these recalls involves knowing how the FDA classifies and communicates the associated health risks. The most common contaminants were found to be 28 (41.2%), 32 (47.0%), 3 (4.4%), 4 (5.9%), and 1 (1.5%) for *Salmonella* spp., *Listeria monocytogenes*, *E. coli*, *Cronobacter sakazakii*, and *Clostridium botulinum*, respectively (Table 1). *C. sakazakii* has been the subject of contamination and product recalls of powdered and dairy formulas in Chile as well [3]. More recently, the most serious food recalls have involved *Salmonella* contamination in foods such as ice cream [4], cereals [5], peanuts [6], soft cheese [7], shelled walnuts [8], crispy fried onions [9], and hummus (Table 1) [10]. Recent French data indicate that, with some yearly decline in the number of cases, meats, dairy, and fishery products are among the most recalled products [11].

Table 1. List of Governmental food recalls requested within a two-year window ¹.

Product Description ²			Recall Reason and Description		
Year 01–12/2023					
<i>Salmonella</i>	<i>Listeria monocytogenes</i>	<i>Escherichia coli</i>	<i>Cronobacter</i> spp.	<i>Clostridium botulinum</i>	Summary
28 (41.2%)	32 (47.0%)	3 (4.4%)	4 (5.9%)	1 (1.5%)	68
Alfalfa Sprouts			Contamination with Shiga toxin-producing <i>E. coli</i>		
All Purpose Flour, bleached and unbleached			Potential to be contaminated with <i>Salmonella</i>		
Bagged Collard Greens			Possible <i>Listeria monocytogenes</i> contamination		
Bean Sprouts			<i>Listeria monocytogenes</i>		
Biltmore Smoked Wild Sockeye Salmon			Potential Foodborne Illness/ <i>Listeria</i>		
Black Fungus (Nam Meo)			<i>Salmonella</i>		
Breeders Choice Raw Pet Food 2 lb			<i>Salmonella</i> contamination		
Cantaloupe Chunks and Cubes and Fruit Mixes and Medleys Containing Cantaloupes (×5)			Potential <i>Salmonella</i> Contamination		
Cheese products			Potential <i>Listeria monocytogenes</i> contamination		
Cocktail Shrimp			<i>Listeria monocytogenes</i>		
Dairy and Non-Dairy ice cream (parve), sorbet, cakes, and novelty items			Potential to be contaminated with <i>Listeria monocytogenes</i>		
Diced onions			<i>Salmonella</i>		
Diced Organic Butternut Squash			Potential <i>Escherichia coli</i> O45 contamination		
Dog and Cat food			Potential to be contaminated with <i>Salmonella</i>		
Dog food, Select Beef Meal & Brown Rice Formula			Potential <i>Salmonella</i> contamination		
Dry Dog, Dry Cat, and Catfish Food			Potential <i>Salmonella</i> contamination		
Enfamil Prosbee Simply Plant-Based Infant Formula in 12.9 oz containers			Potential <i>Cronobacter sakazakii</i> contamination		
Enoki Mushrooms (×4)			<i>Listeria monocytogenes</i>		
Fresh Cantaloupe (×2)			Potential to be contaminated with <i>Salmonella</i>		
Frozen Fruit			Potential for <i>Listeria monocytogenes</i> contamination		
Fruit cups and trays containing cantaloupe (×4)			Potential to be contaminated with <i>Salmonella</i>		
GEISHA Medium Shrimp, 4 oz can (×2)			Potential contamination with <i>Clostridium botulinum</i>		
Gift Baskets with Quaker Chewy Granola Bars			Potential <i>Salmonella</i> contamination		
Granola Bars and Granola Cereals			Potential for <i>Salmonella</i> contamination		
Ground Cumin			<i>Salmonella</i>		
Ice Cream, Yogurt, Ice Cream Bars, and Gelato in a Variety of Flavors			Potential to be contaminated with <i>Listeria monocytogenes</i>		
Kale, Spinach, Collard Green products (×2)			Potential for <i>Listeria monocytogenes</i>		
Lettuce & Salad Kits (×2)			<i>Listeria monocytogenes</i>		
Mini Chunk Chicken Recipe Dry Dog Food			Potential <i>Salmonella</i> contamination		
Multiple frozen fruits with mango products			Potential <i>Listeria monocytogenes</i> contamination		
Mung Bean Sprouts (×2)			<i>Listeria monocytogenes</i>		

Numerous human foods, animal (pet) food, medical devices, and drug products	Potential <i>Salmonella</i> contamination and presence of rodent activity at the distribution center & temperature abuse				
Nutramigen Powder infant formula in 12.6 and 19.8 oz cans	Potential <i>Cronobacter sakazakii</i> contamination				
Organic frozen Pineapple and Frozen Fruit Blend Containing Organic Frozen Pineapple	Potential to be contaminated with <i>Listeria monocytogenes</i>				
Organic green kiwifruit	Possible <i>Listeria monocytogenes</i> contamination				
Organic Tahini	Potential Foodborne Illness/ <i>Salmonella</i>				
Pepper Jack Raw Milk Cheese	Presence of <i>Listeria monocytogenes</i>				
Powdered Infant Formula (×2)	Potential <i>Cronobacter sakazakii</i> contamination				
Premade Salads	Potential <i>Listeria monocytogenes</i> contamination				
Ready-to-Eat (RTE) Sandwiches, Salads, Yogurt, Wraps, and related products	Potential <i>Listeria monocytogenes</i> contamination				
Refrigerated and Frozen Cooked Lobster	Possible <i>Listeria monocytogenes</i> contamination				
Salad Greens	Due to potential <i>Escherichia coli</i> O157:H7 contamination				
Salads Kits	<i>Listeria monocytogenes</i>				
Sambhar Masala and Garam Masala Spices	Potential Foodborne Illness- <i>Salmonella</i>				
Soft Serve ice cream and sorbet cups	Possible <i>Listeria monocytogenes</i> contamination				
Sophelise, Tobasi, and Berkshire Bloom Cheeses (×2)	Potential contamination with <i>Listeria monocytogenes</i>				
Soybean Sprouts in 1 lb bags	Potential contamination of <i>Listeria monocytogenes</i>				
Spinach	<i>Listeria monocytogenes</i>				
Strawberry Granola & Greek Yogurt Parfait Bar	<i>Listeria monocytogenes</i>				
Various Deli Salads	The product may be contaminated with <i>Listeria monocytogenes</i>				
Whole Cantaloupe	Potential to be contaminated with <i>Salmonella</i>				
Whole Peaches, Plums, and Nectarines	Potential to be contaminated with <i>Listeria monocytogenes</i>				
Year 01–12/2024					
<i>Salmonella</i>	<i>Listeria monocytogenes</i>	<i>Escherichia coli</i>	<i>Cronobacter</i> spp.	<i>Clostridium botulinum</i>	Summary
43 (38.7%)	57 (51.3%)	7 (6.3%)	1 (0.9%)	3 (2.7%)	111
Falafel Bites	Potential Shiga toxin-producing <i>Escherichia coli</i> O121:H19 contamination				
Crab meat	Potential foodborne illness— <i>Listeria monocytogenes</i>				
Cashews	Potential to be contaminated with <i>Salmonella</i>				
Aged Cojita Mexican Cheese	Potential to be contaminated with <i>Listeria monocytogenes</i>				
Alfalfa and Onion Sprouts	<i>Listeria monocytogenes</i>				
Atovaquone oral suspension, USP, 750 mg/5 mL	Potential <i>Bacillus cereus</i> contamination				
Turkey Sandwiches	Potential foodborne illness— <i>Listeria monocytogenes</i>				
Baby Arugula	Potential to be contaminated with <i>Salmonella</i>				
Bacon Ranch Crunch Chopped Salad Kit (×2)	Potential to be contaminated with <i>Listeria monocytogenes</i>				
Beef and Chicken Dog Foods	Due to Potential <i>Salmonella</i> and <i>Listeria</i>				
Black bean six-layer dip	Potential to be contaminated with <i>Listeria monocytogenes</i>				
Southwest Chipotle Crunch Salad Kit	Potential to be contaminated with <i>Listeria monocytogenes</i>				
California Shelled Walnuts, Organic Light Halves and Pieces	Due to potential <i>Escherichia coli</i> O157:H7 contamination				
Chicken Chips Dog Treats (×2)	Potential to be contaminated with <i>Salmonella</i>				
Cereal, Bars, and Snacks	Potential to be contaminated with <i>Salmonella</i>				

Cheese, Yogurt, and Sour Cream	Expanded recall for potential <i>Listeria monocytogenes</i> contamination
Chicken Enchiladas Verde, Cilantro Salad Dressing, Elote Chopped Salad Kit, and Southwest Salad	Potential to be contaminated with <i>Listeria monocytogenes</i>
Chicken Street Taco Kit (×6)	Potential to be contaminated with <i>Listeria monocytogenes</i>
Chillimami Sauce	Potential foodborne illness— <i>Clostridium botulinum</i>
Chocolate Caramel Corn and Candy Tray	Possible <i>Salmonella</i> contamination
Cilantro Line Crema, Everything Sauce Fiesta, Cilantro Cotija Dressing, Poblano Caesar Dressing, Cilantro Dressing, Street Taco Express Meal Kit	Potential to be contaminated with <i>Listeria monocytogenes</i>
Coffee Products	Potential to be contaminated with <i>Clostridium botulinum</i>
Cold Smoked Capelin	Potential to be contaminated with <i>Clostridium botulinum</i>
Cucumbers and Salads with Kit	Potential Foodborne Illness— <i>Salmonella</i>
Curly Mustards Greens	Potential foodborne illness— <i>Listeria monocytogenes</i>
Cucumber Slices	Potential Foodborne Illness— <i>Salmonella</i>
Eggs (×2)	Potential to be contaminated with <i>Salmonella</i>
Enchiladas	Potential to be contaminated with <i>Listeria monocytogenes</i>
Enoki Mushroom (×3)	Potential to be contaminated with <i>Listeria monocytogenes</i>
Fresh Express Salsa Ensalada Kit, Southwest Chopped Salada Kit (×2)	Potential to be contaminated with <i>Listeria monocytogenes</i>
Fresh Guacamole Products	Potential to be contaminated with <i>Listeria monocytogenes</i>
Fresh Organic Basil (×2)	Potential to be contaminated with <i>Salmonella</i>
Frozen Waffle Products (×2)	Potential foodborne illness— <i>Listeria monocytogenes</i>
Chicken Caesar Salad Bowl	Potential foodborne illness— <i>Listeria monocytogenes</i>
Green Onions	<i>Salmonella</i>
Ground black pepper	Potential Contamination with <i>Salmonella</i>
Gyro Family Kit	Potential Foodborne Illness— <i>Salmonella</i>
Ham & Cotija Torta Sandwich on Telera Roll	Potential to be contaminated with <i>Listeria monocytogenes</i>
Hedgehog food (×2)	Potential Contamination with <i>Salmonella</i>
Honey Roasted Peanuts and Deluxe Lightly Salted Mixed Nuts	Potential to be contaminated with <i>Listeria monocytogenes</i>
Ice Cream Products	<i>Listeria monocytogenes</i>
Individually Wrapped Sandwiches	Potential to be contaminated with <i>Listeria monocytogenes</i>
Jalapeños, Green Peppers, Green Beans	Potential foodborne illness— <i>Listeria monocytogenes</i>
Smoked Salmon	Potential foodborne illness— <i>Listeria monocytogenes</i>
Kitten Grind, Kitten Mix, Puppy Mix	Potential Contamination with <i>Salmonella</i> and <i>Listeria monocytogenes</i>
Macadamia in the Raw	Potential Contamination with <i>Salmonella</i>
Mango bars in the Variety Pack	Potential Contamination with <i>Salmonella</i>
Meal Kits Containing Chicken	Potential to be contaminated with <i>Listeria monocytogenes</i>
Mediterranean Inspired Party Tray	
Mexican Style Street Corn Bites	Potential to be contaminated with <i>Listeria monocytogenes</i>
Moong Dal, Crunchy Green Gram	Potential Contamination with <i>Salmonella</i>
Multiple Cucumber Products (×3)	Potential Foodborne Illness— <i>Salmonella</i>
Thyroid Formula	Potential to be contaminated with <i>Salmonella</i>

Organic Carrots and Celery	Potential Foodborne Illness—Shiga toxin-producing <i>Escherichia coli</i> O121:H19
Organic Chia Seeds	Potential Presence of <i>Salmonella</i>
Organic Whole Carrots and Baby Carrots	Potential Foodborne Illness—Shiga toxin-producing <i>Escherichia coli</i> O121:H19
Parrot Food	Due to Potential <i>Salmonella</i> contamination
Pepperjack Cheeseburger, Bacon Cheeseburger	Potential foodborne illness— <i>Listeria monocytogenes</i>
Plain Whipped Cream Cheese, Plain Cream Cheese, and Cookies & Cream Mix	Potential to be contaminated with <i>Salmonella</i>
Powdered Goat Milk Infant Formula	<i>Cronobacter</i> spp. contamination
Puppy Mix	Potential Foodborne Illness— <i>Salmonella</i>
Queso de Mano PAISA	Potential to be contaminated with <i>Listeria monocytogenes</i>
Raw Cheddar Cheese	Possible Contamination <i>E. coli</i> O157:H7
Raw Dog and Cat Food	<i>Listeria monocytogenes</i>
RTE Fruit and Vegetables	Potential foodborne illness— <i>Listeria monocytogenes</i>
Shelled Walnuts	Potential foodborne illness— <i>Listeria monocytogenes</i>
Smoked Norwegian Salmon Slices—Toast	Potential to be contaminated with <i>Listeria monocytogenes</i>
Soft Ripened Cheese (×2)	Potential to be contaminated with <i>Listeria monocytogenes</i>
Soybean Sprouts	Potential to be contaminated with <i>Listeria monocytogenes</i>
Spinach and Salad Kits (×3)	Potential to be contaminated with <i>Listeria monocytogenes</i>
Saint Jerome Cheese	Potential foodborne illness— <i>Listeria monocytogenes</i>
Tahini (×2)	Potential to be contaminated with <i>Salmonella</i>
Beef & Lamb Gyro Sandwich Express Meal Kits	May be Contaminated with <i>Salmonella</i>
Torta Sandwiches	Potential to be contaminated with <i>Listeria monocytogenes</i>
Various Confectionery Products (×2)	Potential Contamination with <i>Salmonella</i>
Vegetable Medleys and Whole Organic Carrots	Potential Foodborne Illness—Shiga toxin-producing <i>Escherichia coli</i> O121:H19
Vegetable products (×4)	Potential foodborne illness— <i>Listeria monocytogenes</i>
Whole Cucumbers (×5)	<i>Salmonella</i>
Yogurt-covered pretzels	Potential to be contaminated with <i>Salmonella</i>

¹ Adapted from the United States Food and Drug Administration (US FDA) [12]. ² Counters expressed as ×*n* represent the number of times a specific food item has been subjected to a safety callback.

1.2. Official Guidelines: Regulatory Documents, Methods, and Institutes

The presence of pathogens, especially in RTE foods, and the evidence of antibiotic resistance among these strains have evolved into a worldwide public health challenge [13,14]. Hence, detection and constant surveillance programs of these pathogens in foods are paramount. A primer on novel approaches to analyzing these bacteria in food can be found in recent work [15,16]. From the regulatory standpoint, the European Commission decision 2020/1729 includes the surveillance and antibiotic resistance report of the specific pathogens *Salmonella* spp., *E. coli* (especially those strains with extended spectrum β-lactamases (ESBL), AmpC β-lactamases, or carbapenemases), *C. coli*, *C. jejuni*, *Enterobacter faecium*, *E. faecalis* in foods such as broilers, laying hens, fattening turkeys and pigs, bovine animals under a year of age, fresh meat from broilers, turkeys, pigs, and bovine animals. Analogously, it has regulations regarding microbiological criteria for foodstuffs, specifically, 2073/2005/EC.

Official methods for detection of pathogens include ISO 6579-1:2017, ISO 11290-1:2017, ISO 10272-1:2017/Amd 1:2023, ISO 6888-1:2021, ISO 22964:2017 [17], ISO 16649-1:2018, DIS 13136-1, for *Salmonella* spp., and for *Listeria* spp., *Campylobacter* spp., *Staphylococcus* spp., *Cronobacter* spp., and *E. coli*, respectively. Relevant methods for food analysis also comprise ISO 20976-1:2019, 23418:2022, and 20976-2:2022. These methods have also evolved with the advent of new technologies for detecting and controlling foodborne pathogens [18–24]. Rapid, reliable, and cost-effective methods for bacteria detection are of paramount importance in food safety monitoring [25].

In the case of regulations for antimicrobial susceptibility testing, the European Committee on Antimicrobial Susceptibility Testing (EUCAST) sets epidemiological cut-off values (ECOFFs) and tentative ECOFFs (TECOFFs). For the USA, the Clinical & Laboratory Standards Institute CLSI sets these guidelines, and a workbook is available [26]. Additional guidelines for monitoring and reporting in zoonotic and commensal bacteria include 2013/652/EU (repealed by 2020/1729/EU) and 2003/99 [27]. Finally, ISO 20776-1/-2:2022 sets standards for micro broth dilution methods and other *in vitro* diagnostic tools for testing minimal inhibitory concentration (MIC).

Official methods for detecting pathogens and performing Antimicrobial Susceptibility Testing (AST) combine traditional phenotypic cultures with rapid molecular diagnostics. Standardized by global authorities, these tools are vital for guiding life-saving therapies, managing antimicrobial resistance (surveillance, policy, and stewardship), and ensuring optimal patient outcomes [28].

Only phenotypic results from recognized, standardized *in vitro* antimicrobial susceptibility testing of isolates, along with interpretive criteria, can successfully help classify multiple, extended, pan, and resistant bacteria [29].

Phenotypic testing is the recognized gold standard; it works by exposing isolated bacteria to specific antimicrobial concentrations and observing their actual growth response. While molecular/genotypic tests (such as PCR or sequencing) are excellent for identifying specific resistance genes, they cannot determine the cumulative effect of multiple resistance mechanisms. Because of this, standardizing definitions of drug resistance relies exclusively on phenotypic measurements [28].

1.3. Foodborne Bacterial Pathogens and Antibiotic Resistance

The long-term, overuse and misuse of antibiotics in food animals and humans create ideal conditions for the rising threat of antibiotic resistance. Resistant bacteria in animals may reach humans directly or indirectly through food, water, mud, and manure used as fertilizer [30]. Some types of bacteria (high-burden resistant pathogens such as *Salmonella*, *Shigella*, *Neisseria gonorrhoeae*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus*) [31,32] that cause serious infections in humans have already developed resistance to most or all the available treatments, and there are very few promising options in the research pipeline [33,34].

Interventions that restrict antibiotic use in food-producing animals, such as biosecurity, improved animal management, vaccination, improved hygiene, and changes in animal housing and husbandry practices, have reduced antibiotic-resistant bacteria in these animals [35]. Hygiene practice alone can attenuate the effect of antibiotic selection pressure [36]. These practices, in conjunction with One Health approaches, should be implemented to counter antibiotic resistance [37,38]. In this regard, the One Health approach is a collaborative strategy recognizing that human, animal, plant, and environmental health are deeply interconnected. Because antibiotic resistance easily spreads across these domains, containing it requires integrated monitoring, responsible prescribing, and environmental stewardship across all sectors [39].

The overall reduction in the use of antibiotics in food-producing animals, including the complete restriction of these substances for growth promotion and disease prevention, without proper diagnosis. Malaise animals should be assessed to determine the most effective and prudent antibiotic to treat their specific infection. Meanwhile, healthy individuals should only receive antibiotics to prevent disease if, within a particular population, an animal has already been diagnosed [40].

Antibiotic use is further exemplified by the application of streptomycin (STR), oxytetracycline (OXY), copper-based products, and certain fungicides (e.g., triazoles) in plant agriculture. Their use is correlated with increased resistance among plant pathogens [41–43]. Horizontal gene transfer is responsible for genes being exchanged among a variety of bacteria in the plant production setting, including phytopathogens, soil bacteria, and zoonotic bacteria that are occasionally present in that environment and the food chain [30,42]. Through gene transfer, co-resistance, cross-resistance, and gene up-regulation, resistance to one compound may confer resistance and multi-drug resistance to other similar, or even very dissimilar, compounds [42].

Genetic information encoding bacterial resistance mechanisms can be chromosomal or carried by mobile elements called plasmids. Some bacteria are also inherently resistant to a particular antibiotic if they lack the structure on which the antibiotic acts. Several mechanisms of resistance have continuously evolved, including target modification or mutation, permeability reduction, expression of efflux pump genes, production of hydrolases or inactivating enzymes, metabolic enhancement or auxotrophy, synthesis of target-protective proteins, self-repair systems, changes in cell morphology, and biofilm protection (Figure 1) [44,45]. For a thorough primer on specific mechanisms of resistance by antibiotic class, the work by van Hoeck et al. [46] is suggested.




Figure 1. Schematic representation of common mechanisms of antibiotic resistance exhibited by bacteria. Yellow/Light blue spheres: Antimicrobial/active ingredient. Green arrows: antimicrobial introduction routes or target cell structures. Red arrows: antimicrobial export or target inhibition (e.g., outward export from the cell). Figure prepared using CorelDRAW® Graphics Suite 2026 (Corel Corporation, Ottawa, ON, Canada).

Plasmids are major vectors of bacterial antibiotic resistance [47]. Recent evidence demonstrates that antibiotic resistance-related plasmids are differentially dispersed among humans and livestock, reflecting known antibiotic prescribing practices [48]. For example, using a large-scale multivariate approach, the authors found that only 0.42% of livestock plasmids carried carbapenem resistance (compared with 12% of human plasmids); conversely, tetracycline resistance was enriched in livestock vs. human plasmids [48]. However, they also found that the prevalence of carbapenem and colistin plasmid-related resistant genes is on the rise.

Despite the long list of antibiotics commercially and clinically available (Table 2), resistance has made some of these drugs therapeutically useless against multidrug-resistant bacteria. Hence, the discussion of alternatives to antibiotics has resulted in natural product-led research [49–51]. Several primers regarding natural products with antimicrobial capabilities are advised [52–57]. Interestingly, recent research suggests

that constant bacterial exposure to even these alternatives can affect their resistance profile (that is, decreased sensitivity). Exposure to biocides, pesticides, common home cleaners, heavy metals, food preservatives, and even natural products can lead to a cross-adaptation to antibiotics induced by exposure to non-antibiotic antimicrobials [58–60]. This co-resistance occurs when bacterial exposure to sub-lethal concentrations of non-antibiotic agents, used within the clinical, agricultural, and domestic settings, inadvertently selects strains that are resistant to therapeutic antibiotics (*i.e.*, the physiological changes often double as defense mechanisms against antibiotics) [61]. More worrisomely, evidence of this behavior in pathogenic foodborne bacteria such as *S. aureus*, *E. coli*, *S. typhimurium*, *P. aeruginosa*, *K. pneumoniae*, *L. monocytogenes*, *A. baumannii*, *C. sakazakii*, and other species is clear [59].

Table 2. The ninety-three antimicrobials mentioned during this survey, their mechanisms of action, and their abbreviations.

Drug Class		
Action Mechanism ¹	Antibiotics	Abbreviations
Aminoglycosides		
Inhibition of protein synthesis.	Amikacin	AMK
	Apramycin	APM
	Gentamycin	GEN
	Kanamycin	KAN
	Neomycin	NEO
	Netilmicin	NTM
	Spectinomycin	SPC
	Streptomycin	STR
	Tobramycin	TMP
Ansamycins/Rifamycins		
Inhibits bacterial DNA-dependent RNA synthesis (RNA polymerase).	Rifampicin	RIF
β -lactam/ β -lactamase inhibitor		
Bind to plasmid-mediated β -lactamases and inactivate them.	Amoxicillin-clavulanic acid	AMC
	Ampicillin-sulbactam	SAM
	Cefotaxime-clavulanic acid	CTC
	Ceftazidime-avibactam	CVI
	Ceftazidime-clavulanic acid	FOX
	Piperacillin-tazobactam	TZP
Carbapenems/ β -lactams		
Fix to PBPs ² , inhibiting bacterial cell wall synthesis.	Ertapenem	ETP
	Imipenem	IMP
	Meropenem	MEM
Carboxypenicillins/ β -lactams		
Inhibit cell-wall synthesis.	Carbenicillin	CAB
	Ticarcillin	TIC
Carboxylic acids		
Pseudomonic acid inhibits isoleucine—tRNA ligase in bacteria, leading to depletion of isoleucyl-tRNA & accumulation of the corresponding uncharged tRNA. Depletion of isoleucyl-tRNA results in inhibition of protein synthesis.	Mupirocin (Bactroban)	MCP
Cephalosporins/ β -lactams		
Inhibit cell-wall synthesis.	Cefalexin (1st class)	CPX
	Cefazolin (1st class)	CEZ
	Cefepime (4th class)	CFP
	Cefixime (3rd class)	CFM
	Cefotaxime (3rd class)	CFX
	Cefotetan (2nd class)	CTT
	Ceftiofur (3rd class)	CFT

	Cefovecin	CFO
	Cefoxitin (2nd class)	CEX
	Ceftachlor (2nd class)	CFC
	Ceftazidime (3rd class)	CZA
	Cefpodoxime (3rd class)	CPD
	Cefoperazone (3rd class)	CPZ
	Cefquinome (4th class)	CFQ
	Ceftaroline (5th Class)	CRN
	Ceftriaxone (3rd class)	CTX
	Cefuroxime (2nd class)	CXM
	Cephalothin (1st class)	CEP
Diaminopyrimidines/Sulfonamides		
	Trimethoprim/sulfamethoxazole (Bactrim® or Septra®)	STX
Dihydrofolate reductase inhibitor & Competitive inhibitors of the enzyme dihydropteroate (folate) synthase.	Sulfadiazine	SDZ
	Sulfamethaxole	SMZ
	Sulfafurazole/Sulfisoxazole	INN
	Trimethoprim	TMP
(Fluoro)quinolones		
	Ciprofloxacin	CIP
	Enrofloxacin	ENR
	Flumenique	FMQ
	Gamifloxacin (4th generation)	GMX
Interferes with DNA replication by preventing bacterial DNA from unwinding and duplicating. Specifically, they inhibit the ligase activity of the type II topoisomerases, DNA gyrase, and topoisomerase IV, which cut DNA to introduce supercoiling, while leaving nuclease activity unaffected.	Gatifloxacin (4th generation)	GTX
	Levofloxacin	LFX
	Nalidixic acid	NAL
	Marbofloxacin	MBF
	Moxifloxacin	MXF
	Norfloxacin	NOR
	Ofloxacin	OFX
	Oxolinic acid	OXO
	Pefloxacin	PFX
Phosphonic acid derivatives		
Bactericidal inhibits bacterial cell wall biogenesis by inactivating the enzyme UDP-N-acetylglucosamine-3-enolpyruvyltransferase (MurA)	Fosfomycin (Monurol)	FOS
Glycopeptides		
Bind to acyl-D-alanyl-D-alanine in lipid II, preventing the addition of new units to the peptidoglycan.	Vancomycin (1st gen)	VAN
	Teicoplanin (Targocid)	TEI
Macrolides/Polyketide/Ketolide		
Bacteriostatic, protein synthesis inhibitors. Prevent peptidyltransferase from adding the growing peptide attached to tRNA to the next amino acid. Inhibit bacterial ribosomal (bind reversibly to the P site on the 50S) translation. Dissociate the peptidyl-tRNA from the ribosome prematurely.	Azithromycin	AZM
	Erythromycin	ERY
	Clarithromycin	CAM
	Midecamycin	MDY
	Telithromycin	TYM
	Tylosin	TYL
Nitrofurans		
Rapid reduction inside the bacterial cell by flavoproteins to multiple reactive intermediates that attack ribosomal proteins, DNA, respiration, pyruvate metabolism, and other macromolecules within the cell.	Furazolidone	FZD
	Nitrofurantoin	NFT
Lincosamides		
Bacteriostatic, interferes with the synthesis of proteins. Bind close to the peptidyl transferase center on the 23S portion of the 50S subunit of bacterial ribosomes.	Lincomycin	LIN
	Clindamycin	CDA

Oxazolidinones		
Blocks the initiation of protein production	Linezolid	LZD
Penicillins/β-lactams		
Inhibit the completion of the synthesis of peptidoglycans, the structural component of the bacterial cell wall. Inhibits the activity of enzymes that are needed for the cross-linking of peptidoglycans during the final step in cell wall biosynthesis.	Amoxicillin	AMO
	Ampicillin	AMP
	Aztreonam (monobactam)	ATM
	Cloxacillin	CLX
	Methicillin	MET
	Oxacillin	OXA
	Penicillin G	PEN
	Temocillin	TMC
Phenicol		
Bacteriostatic, inhibits protein synthesis. Prevents protein chain elongation by inhibiting the peptidyl transferase activity of the bacterial ribosome. It specifically binds to A2451 and A2452 residues in the 23S rRNA of the 50S ribosomal subunit, preventing peptide bond formation.	Chloramphenicol	CHL
	Florfenicol	FLO
Polymyxins		
Bind to lipopolysaccharide in the outer membrane of Gram-negative bacteria, polymyxins disrupt both the outer and inner membranes. The hydrophobic tail is important in causing membrane damage, suggesting a detergent-like mode of action (increase membrane permeability/disorganization).	Colistin (Polymyxin E)	COL
	Polymyxin B	PMB
Steroids		
Binds to EF-G after translocation and GTP (guanosine-5'-triphosphate) hydrolysis.	Fusidic acid (Fucidin)	FAS
Streptogramins		
Bacteriostatic, binds to the bacterial 50S ribosomal subunit and inhibits the elongation process of protein synthesis.	Pristinamycin	PRI
Tetracyclines		
Protein synthesis inhibitors. Prevent the initiation of translation in various ways by binding to the 30S ribosomal subunit. Hinder the binding of aminoacyl-tRNA to the mRNA translation complex.	Doxycycline (Semi-synthetic)	DOX
	Minocycline (Semi-synthetic)	MIN
	Oxytetracycline (Naturally occurring)	OXY
	Tetracycline (Naturally occurring)	TET
	Tigecycline (Glycacyclines)	TIG
Ureidopenicillins/β-lactams		
Inhibit cell-wall synthesis	Mezlocillin	MEZ
	Piperacillin	PIP

¹ Mechanisms of action for antimicrobials adapted from Singh et al. [62]. ² PBPs, Penicillin-binding proteins.

2. *Salmonella* spp.

2.1. Foods for Human Consumption

Salmonella is a genus of rod-shaped (bacillus), motile, non-spore-forming, chemotrophic, facultative anaerobe, and Gram-negative bacteria of the family Enterobacteriaceae. It is known for its two species, *S. enterica* and *S. bongori*. *S. enterica* is further divided into six subspecies that include over 2600 serotypes [63,64].

Nontyphoidal *Salmonella* serotypes are zoonotic and can be transferred from animals to humans. Infection by this pathogen is usually due to ingestion of contaminated, undercooked food. *Salmonella* is one of the most common causative agents of foodborne illness in humans, causing approximately 1.35 million infections, 26,500 hospitalizations, and 420 deaths in the United States alone every year [65].

Treatment of salmonellosis is further complicated by the fact that isolates may also carry resistance genes conferring resistance to antimicrobials. For example, *Salmonella* species carrying antimicrobial resistance genes have been isolated from animal- and plant-derived foods, including poultry, eggs, fish,

pork, beef, vegetables, fruits, nuts, and flour [56]. Outbreak studies have recognized food commodities, including fresh produce, vegetables, cantaloupes, cereals, alfalfa sprouts, fruit/fruit pulp, pistachios, poultry meat, turkeys, tuna, ground beef, pork, dried/shredded coconut, tomatoes, and mangos, as sources of foodborne *Salmonella*-related outbreaks in the past decade [66–69].

Soubeiga et al. [70] found *Salmonella*-related virulence genes associated with vegetable/fruit sources. Multidrug resistance profiles were consistent determinants in the 148 isolates, as genes *bla*_{TEM}, *temAB* (β -lactams), *StrA*, *aadA* (aminoglycosides), *sul1*, *sul2* (sulfonamides), and *tetAB* (tetracyclines) were consistently present (Figure 2).




Figure 2. Reported resistance profiles for (A) Tetracycline and (B) Ampicillin for foodstuffs and related matrices (assorted meats, RTE, feces, aquatic products, and poultry) from China from 2016–2024. China was selected as it is the country with the most entries (nine) as *per* Table 3. Figure elaborated using Sigma Plot 16 (Grafiti LLC, Palo Alto, CA, USA).

Similar gene profiles [e.g., *aadA*, *aac*, *aph*, *bla*_{CARB}, *bla*_{PSE}, *bla*_{TEM}, *bla*_{CTX-M}, *bla*_{CMY}, *floR* (phenicols), folate pathway inhibitors *sul* and *dfrA* (folate pathway inhibitors), *qnrAB*, *oqxAB* (fluoroquinolones), *tetABM*, and *mph(A)* and *lnu(F)* (macrolides)] were found recently, but in retail American poultry [71], Mexican ground beef samples [72], and Chilean poultry production line samples [45]. Analogous gene profiles were also found in food chain links in Poland [73]. On the other hand, recent Italian isolates from swine mostly exhibited resistance determinants *parC*, *gyrA*, *catA*, *nfsAB*, *bla*_{TEM}, and *tetAB*. For a detailed

primer regarding mechanisms of resistance for *Salmonella*, the work by Monte et al. [74] and Wang et al. [75] is recommended.

Furthermore, as global trends indicate that more people are basing their diets on fresh produce and vegetables [76], *Salmonella* contamination is relevant, as an increase in produce-related outbreaks has been documented [77]. Additionally, multidrug-resistant bacteria can disseminate more quickly due to an ever-expanding regional and international food trade [77]. More problematic still, it is easy to forget that foods, in general, and especially fruit and vegetables [78], can carry more than one pathogen, and these pathogens can carry resistance genes and even share them amongst themselves and other bacteria. In a very thorough Cuban study, 14 different food groups were analyzed for the presence of *Salmonella* spp., *E. coli*, *Staphylococcus* spp., and *V. cholerae* [79]. Meat and dairy products and RTE foods showed not only multiple pathogen contamination but also multiple drug resistance profiles [79]. A recent study saw lettuce, cabbage, spinach, carrot, tomato, and green peppers contaminated with other antibiotic-resistant species alongside *Salmonella* (e.g., *E. coli*, *S. aureus*, *S. epidermis*) [78].

On the other hand, Cao et al. [80] found a high prevalence of *Salmonella* (isolating the bacteria at a rate of 29.4% per sample tested) contamination in cow's milk. A similar account was reported by Ejo et al. [81], who found both a higher prevalence of contamination and a drug resistance profile in raw meat, eggs, and milk.

Quesada et al. [82] reviewed *Salmonella* reports from 2003–2014 from Latin America, as well as the antibiotic-resistant profiles of these isolates (especially recovered from the Brazilian and Mexican poultry industry). Among their findings, *S. Enteritidis*, *S. Typhimurium*, *S. Paratyphi*, and *S. Heidelberg* were the most commonly encountered serovars. Similar to other reports [83,84] most strains were multidrug-resistant and showed a high tolerance to STR, NAL, AMP, TET, CHL, and STX (Table 3). Relatedly, a 22-year historical analysis of antibiotic-resistant *Salmonella* from Bangladesh found 81% tetracycline resistance [85], rendering this antibiotic effectively without any therapeutic or clinical use. However, the continued use of tetracyclines in other agricultural settings (e.g., crop production, apiculture, and aquaculture) still provides a selective pressure on bacteria [40].

As macrolides (the first-choice drugs) for *Salmonella* infections in humans have failed, cephalosporins are now the paramount therapeutics. However, ESBL enzymes encoded by *bla*_{CMY2}, *bla*_{SHV}, and *bla*_{TEM} emerged, thus reducing the efficacy of cephalosporins. Also, generalized *mcr* genes have been described in *Salmonella* and encode colistin resistance, an antibiotic regarded as critical. In both cases, resistance genes were carried by conjugative plasmids (e.g., IncI2, IncHI2, IncX3, IncX4) [86].

Interestingly, Siddique et al. [87] not only found *Salmonella* strains carrying β -lactam, cephalosporins, and carbapenems resistance genes (e.g., *bla*_{CMY-2}, *bla*_{OXA-1}, *bla*_{PSE-1}, *bla*_{TEM-1}, *bla*_{NDM-1}, and *ampC*), and the zoonotic potential virulence gene *sopE*, but also noted that the strains recovered had moderate-to-high abilities to aggregate into biofilms at 30 and 37 °C. Other virulence factors (e.g., *fimA*, *invA*, *spvC*) co-occurring with similar resistance gene patterns have been recently reported in Poland [73], Italy [88] and China [89–92].

For Brazilian poultry farms, Rau et al. [93] found, in two different sampling moments (*i.e.*, 3 years apart), that traditional *Salmonella* serovars Typhimurium and Enteritidis have been supplanted by other serovars like Heidelberg or Kentucky. In the same period, Brazil saw an increase in the number of isolates exhibiting antimicrobial resistance by almost double. Similar phenomena seem to be occurring of late in countries like the USA and Costa Rica [94,95].

As the debate about rendering animals without antibiotics rages on, evidence suggests that alternative meat production practices reduce the frequency of bacteria exhibiting antibiotic resistance, which favors “organic” practices [71]

2.2. The Relation Between the Poultry Industry and the Emerging *Salmonella* Infantis (pESI)-Plasmid

Salmonella enterica serovar Infantis (or *S. Infantis*) is an emergent serotype worldwide (Table 3) [96]. This serotype has been reported as one of the most prevalent non-typhoidal *Salmonella* (NTS) in humans in Europe and the USA [97–99]. Several studies describe *S. Infantis* as the most prevalent serotype in poultry [98,99]. Recent data from over 10,000 isolates demonstrated that *S. Infantis* and *S. Seftenberg* with pESI plasmid carrying a *bla*_{CTX-M-65} gene constitute one of the biggest clusters of *Salmonella* along the United States, both originating from turkeys [100].

Moreover, multidrug-resistant phenotypes of this serotype are of public health concern [101]. For example, Mattock et al. [102] recently related the presence of pESI megaplasmid with 73% of the poultry isolates from 74 different countries showing multidrug resistance. Hence, the presence of *Salmonella* in poultry production has attracted serious attention, and several studies have found in poultry a vector for several species of this bacterium. Whole genome sequencing from broiler integrations determined that the most prevalent non-typhoid *Salmonella* serotype was *S. Infantis* [103]. Not only do most isolates show potential for resistance against medically important antibiotics [especially aminoglycosides (mutations in genes *aac*, *ant*, *aph*) and quinolones (*parC*, *gyrA*)] and disinfectants, but also almost all isolates harbored the pESI-like megaplasmid, which plays an important role in the global dissemination of antimicrobial resistance. Among analogous Latin American studies, the *bla*_{CTX-M-65} ESBL antibiotic resistance gene is also prevalent in isolates. This was more recently confirmed by Costa Rican-led research, in which ampicillin resistance in the more prevalent *S. Infantis* reached 81.6% [95]. In terms of plasmid prevalence, a similar trait has been observed in *S. Muenchen*, *S. Schwarzengrund*, *S. Agona*, and *S. Senftenberg* in a large-scale broiler and chicken isolate study [104]. Previous studies, again in the Andean Region (mostly Ecuador, Chile, and Peru), seem to support similar findings [105–109]. A more recent Chilean study [47] has reported mobile elements carrying multiple drug resistance determinants in the poultry production line, predominantly related to *S. Infantis* and *S. Heidelberg* (IncFIB and Inc11-*Ia* plasmids, respectively).

Furthermore, the ESI clone carrying an extended spectrum β -lactamase *bla*_{CTX-M-65} on a pESI-like plasmid and a mutation in the *gyrA* gene has also been found in poultry meat in regions such as Russia, USA, Italy, Germany, Hungary, Poland, and Slovenia [110–114]. All these data suggest that pESI is not only widespread but is acquired by *Salmonella* spp. with relative ease. Finally, recent evidence has described pESI-like megaplasmids as carriers of additional antimicrobial resistance genes, such as *tet*, *dfr*, and *sul*, and has demonstrated that these structures are genetically plastic and can also harbor virulence and fitness genes, as well as toxin/antitoxin systems that enhance their persistence in the *S. Infantis* host [97].

2.3. Animal Feeds and Related Matrices

Salmonella monitoring and surveillance in animal feed are not only related to production costs and animal welfare but are also instrumental in understanding the behavior of foodborne pathogens at the start of the food chain and how this relates (if any) to prevalence, drug-resistant profiles, and specific species found downstream in human or clinical isolates.

Relatively few studies have tackled feed contamination despite being considered a main source of contamination on farm grounds (Table 3) [65]. In Thailand, the prevalence of *Salmonella* contamination in compound feeds (a specially formulated, balanced mixture of feed components designed for feeding livestock and poultry) was registered at 12.5% [115]. Similar data have been recovered from isolates from American and Australian feeds [71,116–118]. The authors found more prevalence in the compound feed than in any of its respective ingredients. However, Parker et al. [118] recently related contamination hotspots with the milling equipment during feed manufacturing. Hsieh et al. [117] found an 84% contamination in fish meals and 48% in animal products. From these samples, the acquisition of the *bla*_{CMY-2} gene, which confers resistance to cephalosporins, was most commonly observed. Interestingly, multidrug-resistant isolates were

recovered from monensin (an ionophore/Rumensin[®])-medicated dry beef cattle feeds. A possible association between the use of ionophores in livestock and antibiotic resistance has recently been raised [119].

Of all the bacterial pathogens discussed, *Salmonella* seems to have received the most attention in the feed sector (both in surveillance and research). Antibiotic resistance in *Salmonella* often stems from agricultural practices, where antimicrobial usage in animals selects resistant strains that spread into the human food supply. This poses a major public health issue, as these pathogens can trigger severe infections that are difficult to treat with standard medications. The cycle of how antibiotic resistance moves from animal feed to human consumers operates as follows: i. Contaminated Feed, ii. Livestock Exposure, iii. Indirect Transmission to Humans [120].

As stated, however, there is a very close link between livestock and human pathogen contamination. For instance, the main source of *Salmonella* infections in humans is contaminated animal-derived foodstuffs, with pork products being among the most important contributors [65].

Table 3. Summary of bacterial isolates from foodstuffs and other food-related matrices for *Salmonella*.

Country	Isolates	Source	Antibiotics Tested	Resistance to ¹	Testing Method	References
<i>Feeds</i>						
USA	<i>S. Senftenberg</i> , <i>S. Montevideo</i> , <i>S. Mbandaka</i>	Animal feeds (finished feeds), feed ingredients, pet foods, treats, & supplements (<i>n</i> = 2058)	AMC, AMP, CEX, CHL, CIP, GEN, KAN, NAL, NOR, STX, INN, STR, TET	TET, INN, AMP, NAL	Sensititre system (micro version of the broth dilution method)	[116]
Costa Rica	<i>S. Give</i> , <i>S. Rissen</i> , <i>S. Havana</i> , <i>S. Soerenga</i> , <i>S. Schwarzengrund</i>	Feed & ingredients (<i>n</i> = 1725; meat and bone meal, poultry feed, pet foods)	TET	TET (7.4%)	E-test	[121]
USA	<i>S. Newport</i> , <i>S. Typhimurium</i> var. O 5-(Copenhagen), & <i>S. Lexington</i> var. 15+ (Manila)	Dry beef (<i>n</i> = 3) & dry cattle feeds (<i>n</i> = 1), cabbage culls (<i>n</i> = 2), animal protein products (<i>n</i> = 2), & fish meal (<i>n</i> = 1)	AMC, AMP, CEX, CFT, CHL, KAN, STR, INN, TET, SXT	AMC, AMP, CEX, CFT, CHL, STR, INN, TET	Sensititre automated antimicrobial susceptibility system	[117]
Tanzania	<i>Salmonella</i> spp.	Commercial chicken feed samples (<i>n</i> = 384; Broiler starter mash (<i>n</i> = 121), Broiler grower mash (<i>n</i> = 47), Broiler finisher mash (<i>n</i> = 102) & Layers mash (<i>n</i> = 114)	STR, AMO, AMK, CIP, TET, KAN, GEN, STX	TET, AMK, KAN, GEN, STX	Kirby-Bauer disk-diffusion	[122]
Kenya	<i>Salmonella</i> spp.	Poultry feed (<i>n</i> = 150)	AMP, CTX, STX, TET, CHL, CIP	AMP (41%), STX, CTX	Kirby-Bauer disk-diffusion	[123]
Colombia	<i>Salmonella</i> spp.	Feed for swine, poultry, canine, feline, leporine, & equine species (<i>n</i> = 1758)	AMP, SAM, CEZ, CFP, CEX, CTX, CIP, IMP, MEM, TZP	AMP (44.5%), SAM, CEZ	Becton Dickinson Phoenix TM Automated Dilution System	[124]
Costa Rica	<i>S. Typhimurium</i> , <i>S. Braenderup</i> , <i>S. Newport</i> , <i>S. Anatum</i> , <i>S. Westhampton</i>	Non-human primate feces (<i>n</i> = 180) & feed (<i>n</i> = 43)	AMP, AMC, TPZ, CFX, CZA, CFP, IMP, MEM, AMK, GEN, NAL, CIP, NFT, COL, STX	CIP (9.8%)	VITEK [®] 2 AST-N279 cards	[125]
Thailand	<i>S. Rissen</i> , <i>S. Mbandaka</i> , & <i>S. Livingstone</i>	Compound feed (<i>n</i> = 60), Soybean (<i>n</i> = 10), pork (<i>n</i> = 5), fish (<i>n</i> = 3) & poultry meal (<i>n</i> = 2)	AMP, AMC, PIP, CPX, CFP, CFT, CFO, AMK, GEN, TMP, IPM, STX, CHL, ENR, MBF, TET, DOX, MIN, NFT	TMP (100%), CPX, AMK, GEN	VITEK [®] 2 Card type AST-GN65	[115]
<i>Foods and related matrices</i>						
Thailand	<i>S. Rissen</i> , <i>S. Weltevreden</i> & <i>S. Bredeney</i> for pork; <i>S. Albany</i> & <i>Typhimurium</i> for chicken; <i>S. Typhimurium</i> & <i>S. Hvitittingfoss</i> for vegetables	Pork (<i>n</i> = 41) & chicken meat (<i>n</i> = 40) & fresh vegetables (<i>n</i> = 80)	TET, CIP, CHL, AMP, NOR, NAL, STR, STX, CEP, GEN	STR (92%), TET, NAL, CHL, AMP, CEP	Kirby-Bauer disk-diffusion	[126]

Türkiye	<i>Salmonella</i> spp.	RTE salads & vegetables ($n = 81$)	KAN, NEO, NAL, TET, SPC, STX, AMP, AMC, CHL, FLO, GEN, CIP, STR, CFT	NEO, NAL	Micro dilution method	[127]
Nigeria	<i>Salmonella</i> spp.	Fecal samples of cattle, pig, sheep, goat, & chicken ($n = 200$ each)	AMP, AMO, AMC, CHL, STR, TET, CEP, GEN, CIP, SMZ, NAL	Cattle [TET (85%), AMP, CIP, NAL], Sheep [AMP (25%), STR, CHL, TET] Goat [TET (44%), AMP, CHL, STR, CEP, SMZ, NAL], Kirby-Bauer disk-diffusion [TET (90%), NAL, SMZ, AMP, CHL, STR, CEP], Pig [TET (75%), AMP, CHL, CEP, SMZ, NAL]		[128]
Nigeria	<i>S. Typhi</i> , <i>S. Paratyphi</i> & <i>S. Typhimurium</i>	Cabbage, lettuce, cucumber, tomatoes, green pepper & spinach	AMC, STX, CIP, CTX, GEN, TET, KAN, CEP	CEP, STX, TET, KAN, multidrug-resistant profile	Kirby-Bauer disk-diffusion	[129]
Poland	<i>S. Enteritidis</i>	Foods other than meat: confectionery products, eggs, fruits, vegetables, spices	ATM, AMC, AMP, CFP, CFX, CEX, CZA, CTX, CHL, CIP, ETP, GEN, IMP, NAL, STR, TMP, TET, STX	NAL (35.2%)	Kirby-Bauer disk-diffusion	[130]
Malaysia	<i>Salmonella</i> spp., <i>S. Enteritidis</i> & <i>S. Typhimurium</i>	Ulam (salad, fresh leaves, vegetables, or fruits; $n = 96$)	CHL, CIP, AMP, ERY, CEP, STR, AMC, NAL, GEN, STX, TET	AMP (100%), ERY (100%), AMC, CEP, STR, CIP	Kirby-Bauer disk-diffusion	[131]
Ghana	<i>S. Kentucky</i> , <i>S. Nima</i> , <i>S. Muenster</i> , & <i>S. Enteritidis</i>	Laying hens & broilers (feces $n = 75$, dust $n = 75$, feed $n = 10$ & drinking water $n = 10$)	AMP, CFX, CEX, GEN, CZA, TET, CHL, TMP, SMZ, NAL	NAL, TET, CIP	Kirby-Bauer disk-diffusion	[132]
Ethiopia	<i>Salmonella</i> spp.	Animal-origin food samples ($n = 384$: egg sandwich, minced & raw meat, burger patties, cottage cheese, cream cake, beef pizza, raw egg & pasteurized & raw milk from restaurants, hotels, cafeterias, & retail shops)	AMO, AMP, CEP, CTX, GEN, NAL, NFT, STX, TET	TET (42.6%), STX, AMP	Kirby-Bauer disk-diffusion	[81]
Nigeria	<i>Salmonella</i> spp.	Raw chicken (retail table) eggs ($n = 340$)	NEO, DOX, STX, AMO, VAN, NFT, PEN, CIP, ERY, GEN, STR, TET, OXA, AMC	STR (85%), AMC, GEN, VAN, DOX	Kirby-Bauer disk-diffusion	[133]
China	<i>S. Typhi</i> , <i>S. Enteritidis</i> , & <i>S. Typhimurium</i>	Pork ($n = 85$) & chicken ($n = 200$) at retail stores	CFX, GEN, LFX, TET, DOX	TET	Kirby-Bauer disk-diffusion	[89]
South Africa and Brazil	<i>Salmonella</i> spp.	Broiler chicken caecum samples ($n = 200$)	GEN, AMO, ERY, CHL, TET, TMP, AMP, STR, STX	TET (93%), STX, TMP, KAN, GEN, AMP	Kirby-Bauer disk-diffusion	[134]
China	<i>Salmonella</i> spp.	Pork ($n = 137$), chicken ($n = 91$), & beef ($n = 99$) from supermarkets	AMO, AMC, AMP, CFT, CIP, OXY, LFX, TMP, GEN	OXY (80.9%), TMP, AMP, LFX	Agar dilution method	[58]

Malaysia	<i>S. Corvallis</i> , <i>S. Typhimurium</i> , <i>S. Braenderup</i> , <i>S. Albany</i>	Spent hens ($n = 470$)	AMP, CTX, CHL, CIP, ERY, GEN, KAN, NAL, STR, STX, TET	ERY (97%), TET, STR, STX, AMP, NAL	Kirby-Bauer disk-diffusion	[135]
Latvia	<i>S. Typhimurium</i> , <i>S. Derby</i> , <i>S. Enteritidis</i>	Meat and meat products ($n = 3152$)	AMP, ATM, CFX, CZA, CHL, MEM, CIP, TET, TIG, NAL, COL, GEN, TMP, SMX.	SMX (38%), NAL, CIP, AMP, TET	Broth microdilution (Sensititre® EUVSEC3® & EUVSEC2® plates)	[136]
Ghana	<i>Salmonella</i> spp.	Cabbage and lettuce samples	AMP, PEN, CHL, CIP, CTX, GEN, ERY, OFX, STX, TET	AMP (72.2%), ERY, OFX	Kirby-Bauer disk-diffusion	[137]
Vietnam	<i>Salmonella</i> spp.	Fecal samples ($n = 240$) & Environmental samples (feed, water, barn floor & wastewater; $n = 640$) from household and pig farms	AMC, CZA, CXM, COL, GEN, AMK, STR, TET, DOX, CHL, STX, AMP, OFX, LFX	STX (64.5%), CHL, AMP	Kirby-Bauer disk-diffusion	[138]
Bangladesh	<i>Salmonella</i> spp.	Fuska ($n = 55$), sugarcane juice ($n = 58$), Borhani ($n = 30$)	AMP, AMO, CIP, ENR, COL, OXY, TET, AZM, ERY, PFX, CTX	High level of resistance to all antibiotics tested, especially TET, AZM, ERY, AMP, AMO	Kirby-Bauer disk-diffusion	[139]
Ethiopia	<i>S. Dublin</i> , <i>S. Virchow</i>	Fecal samples of slaughtered cattle ($n = 720$)	AMP, AMC, CHL, CTX, CEP, CIP, CEX, GEN, KAN, STX, TMP, TET, SMZ, STR, NFT, NAL	STR (89.3%), CEP, AMP, AMC	Kirby-Bauer disk-diffusion (BD BBL™ Sensi-Disc™)	[140]
USA	<i>S. Newport</i>	Humans ($n = 2763$), animals ($n = 901$), & retail-meats ($n = 64$)	AMK, APM, GEN, KAN, STR, AMC, TZP, CEP, CEX, CTX, CFT, CZA, CFX, CFQ, CFP, INN, STX, AZM, ATM, IMP, AMP, CHL, CIP, NAL, TET	Multidrug resistance-AMP & single drug resistance-TET	Agar dilution Method	[141]
South Africa	<i>S. bongori</i> , <i>S. Pullorum</i> , <i>S. Typhimurium</i> , <i>S. Weltevreden</i>	Fecal samples (poultry pens of broilers, layers & indigenous chickens, $n = 20$ each)	AMP, OXY, CIP, STR, GEN, STX, CHL, ERY, NOR, CEP, NAL	ERY (100%), STR, STX, OXY, AMP	Kirby-Bauer disk-diffusion	[142]
Nigeria	<i>S. Enteritidis</i> , <i>S. Typhimurium</i>	Ready to eat shrimp ($n = 1440$)	KAN, GEN, STR, ERY, TOB, AMP, AMO, IMP, SAM, MEM, CEP, TMP, AMC, COL, CHL, CHL, PEN, PMB, OXY, DOX, TET, OFX, CIP	ERY, PEN (100%), AMO (92.9%), AMC, AMP (85.7%)	Kirby-Bauer disk-diffusion	[143]
China	<i>S. Enteritidis</i> , <i>S. Typhimurium</i> , <i>S. Kentucky</i>	Feces, cecum, & milk ($n = 1578$) from chicken, pig, duck, & cows	AMC, AMP, CFT, CZA, GEN, ENR, OFX, MEME, STX, SPC, APM, FLO, INN, TET, COL	AMP (59.6%)	Not indicated	[80]

China	<i>S. Derby</i> , <i>S. London</i> , <i>S. Rissen</i>	Pork, chicken & the Environment (<i>n</i> = 1112)	AMP, TET, PIP, CZA, DOX, CTX, MIN, NOR, CHL, SMX, OFX, CIP	TET (73.04%), AMP, DOX	Kirby-Bauer disk-diffusion	[144]
Thailand	<i>S. Brunel</i> & <i>S. Hvittingfoss</i> for vegetable, <i>S. Stanley</i> for fermented pork, <i>S. Tennessee</i> for fermented fish	Vegetables (<i>n</i> = 100) & 72 Thai fermented foods (<i>n</i> = 72, pla-som & nham)	GEN, TET, CIP, STX, NAL, SMZ, CEP, NEO, NFT	SMZ (20%), TET, STX, NAL, GEN	Kirby-Bauer disk-diffusion	[145]
USA	<i>S. Typhimurium</i>	Humans (<i>n</i> = 6381), animals (<i>n</i> = 2940), & retail meats (<i>n</i> = 2126)	AMK, APM, GEN, KAN, STR, AMC, TZP, CEP, CEX, CTX, CFT, CZA, CFX, CFQ, CFP, INN, STX, AZM, ATM, IMP, AMP, CHL, CIP, NAL, TET	AMP, CHL, STR, STX, TET, AMC, CTX, CFT	Agar dilution Method	[146]
China	<i>S. Derby</i> , <i>S. Typhimurium</i> , <i>S. London</i> , <i>S. Rissen</i>	Pork (<i>n</i> = 287), beef (<i>n</i> = 161), mutton (<i>n</i> = 92), dumplings (<i>n</i> = 212), & smoked pork (<i>n</i> = 55).	AMP, AMC, CEZ, CEX, CTX, CZA, CFX, CFT, CFP, ATM, IMP, GEN, KAN, AMK, STR, TET, CIP, ENR, NAL, STX, CHL, FLO	TET (65.6%), AMP, STX, STR, NAL	Kirby-Bauer disk-diffusion	[147]
Brazil	<i>S. Mbandaka</i> , <i>S. Braenderup</i>	Layer hens' fecal samples (<i>n</i> = 2008)	NAL, ENR, NOR, CIP, CHL, KAN, STR, GEN, AMK, STX, CFX, CFP, CFT, AMP, AMO, AMC, IMP, ATM, TET, OXY, PMB	STX, STR, CIP	Kirby-Bauer disk-diffusion & dilution method for PMB	[148]
Zambia	<i>Salmonella</i> spp.	Raw retail table eggs (<i>n</i> = 1080)	AMC, AMP, CTX, IMP, CHL, NAL, CIP, COL, TET	TET, COL (83.3%)	Kirby-Bauer disk-diffusion	[149]
Ecuador	<i>S. Infantis</i> , <i>S. Gallinarum</i>	Retail fresh meats (<i>n</i> = 1095, chicken, pork, veal, lamb, beef, and turkey)	STR, SXT, CTX, AMC, CIP, NAL, TET, CHL, GEN, AMP, AZM, KAN	TET (97.1%), NAL, STR, CIP, AMP, CTX (all > 90% resistance)	Kirby-Bauer disk-diffusion	[150]
Cuba	<i>Salmonella</i> spp.	Meat, dairy, eggs & seafood, RTE foods	NAL, AMK, AMP, AZM, CAB, CFX, CZA, CTX, CIP, CHL, DOX, ERY, STR, GEN, KAN, OXA, PEN, STX, TET	TET (59.3%), NAL, AMP, CAB	Kirby-Bauer disk-diffusion	[79]

China	<i>Salmonella</i> spp.	Fecal samples patients with diarrhea ($n = 14,579$), chicken meat ($n = 1237$), pork ($n = 1354$), aquatic products ($n = 814$)	AMP, SAM, CZA, CFX, IMP, MEM, STX, GEN, TET, CIP, NAL, CHL	Human: AMP (68%), TET (68%), NAL (56%); Aquatic food products: TET (56%), CHL (48%), AMP (42%); Chicken meat: NAL (82%), AMP (53%); Pork: TET (85%), AMP (68%), NAL (68%), STX (53%)	Broth dilution method and Biofosun Gram-negative bacterial panels	[90]
Rwanda	<i>Salmonella</i> spp.	Pig fecal samples ($n = 180$)	AMC, AMP, AZM, CEX, CTX, CHL, CIP, COL, MEM, NAL, STR, TET	All pan-susceptible	Kirby-Bauer disk-diffusion	[151]
Vietnam	<i>Salmonella</i> spp.	Chicken feces ($n = 417$), Environmental samples [bedding ($n = 170$), feed ($n = 219$), & drinking water ($n = 142$)], & Pest animals [rats ($n = 32$), geckos ($n = 808$), flies ($n = 450$), ants ($n = 530$), & cockroaches ($n = 287$)	AMP, AMK, AMC, CZA, CHL, COL, CXM, DOX, GEN, OFX, LFX, STR, STX	CHL (63.0%), TET, AMP, STX	Kirby-Bauer disk-diffusion	[152]
Brazil	<i>S. Heidelberg</i> , <i>S. Minnesota</i> & <i>S. Saintpaul</i>	Poultry meat	AMP, AZM, CZA, CHL, CIP, COL, CFX, GEN, MEM, NAL, STX, TET, TIG	CIP (91.3%), NAL, AMP, CZA, CFX, TET	Broth microdilution	[93]
Pakistan	<i>S. Typhimurium</i> , <i>S. Enteritidis</i>	Poultry feces ($n = 180$), organs ($n = 70$), meat ($n = 60$), & eggs ($n = 60$)	AMK, CHL, TET, AMC, CIP, GEN, NAL, STX, AMP, IMP, MEM, STR, ERY, MIN, KAN, CFM, CFP, VAN, LZD, RIF, ENR, OXA, CDA	All isolates are resistant to OXA, CDA, ERY, NAL, LZD, RIF, TET, MIN & VAN. ENR (95%), GEN (93%), KAN (91%), STX (91%), AMP & AMC (81%), CFP (76%)	Kirby-Bauer disk-diffusion	[87]
USA	<i>Salmonella</i> spp.	Poultry (Chicken, breasts with bone-in & skin on, legs, wings, & thighs, $n = 27,591$ & ground turkey products, $n = 19,346$). Conventional & “organic”	GEN, STR, AMP, AMC, CEX, CFT, CTX, AZM, CHL, NAL, CIP, INN, STX, TET	TET (57.6–75.0%), GEN/STR (27.0–52.9%), AMP (17.7–44.3%)	Broth microdilution	[71]
Italy	<i>S. Infantis</i> , Monophasic Variant of <i>S. Typhimurium</i> 4 [5], 12:I (MVST), <i>S. Typhimurium</i> , <i>S. Kentucky</i>	Meat (broiler, swine & cattle), mixed meat, mussels, sheep milk & cheese & feed. Carcasses, feces, swabs from poultry (broiler and laying hens), cattle, sheep & pigs.	AMP, AZM, AMK, GEN, TIG, CZA, CFX, COL, NAL, TET, TMP, SMZ, CHL, MEM, CIP, IMP, CFP,	NAL (63.6%), CIP, TET, TMP, AMP, CFX, CZA	Broth microdilution (Sensititre® EUVSEC3® & EUVSEC2® plates)	[153]
Italy	<i>S. Typhimurium</i> , <i>S. Enteritidis</i> , <i>S. Rissen</i>	Pig slaughterhouses ($n = 150$) carcasses, cecal samples ($n = 30$), environmental samples ($n = 180$) & traditional pork dry sausages ($n = 75$).	AMP, AMC, CPD, CFT, TMP, MBF, PIP, GEN, AMK, ENR, CHL, TET, NTF, STX	AMK, TMP, GEN (86.1%), TET, STX	VITEK® 2 AST GN-65	[154]

Iran	<i>Salmonella</i> spp.	Chicken meat samples (<i>i.e.</i> , breast muscle, $n = 150$)	PEN, TYL, TET, ERY, TIA, FOS, SXT, DFX, LIN/SPC, FLM, FLO	TET, PEN, TYL, ERY, TIA (100%)	Kirby-Bauer disk-diffusion	[155]
Russia	<i>S. Infantis</i> , <i>S. Enteritidis</i> , <i>S. Typhimurium</i>	Poultry ($n = 112$), pork ($n = 91$), beef ($n = 103$), & minced meat ($n = 168$)	AMP, IMP, AMK, STR, TOB, CFX, CEZ, STX, CHL, ATM, AMC, NFT, TET	STR (75%), STX, TET, ATM, CEZ	Kirby-Bauer disk-diffusion	[156]
Chile	<i>S. Typhimurium</i> , <i>S. Infantis</i> , <i>S. Enteritidis</i> , <i>S. Derby</i>	Stool samples from pigs ($n = 500$), from chickens ($n = 300$)	AMP, AMC, CZA, CFT, CTX, CIP, GEN, NAL, STX, TET, STR, ENR, ATM, TMP, SMZ, CHL	Chickens: NAL (94.7%), Pigs: TET (53.3%)	Kirby-Bauer disk-diffusion	[157]
Italy	<i>S. Derby</i> , <i>S. Typhimurium</i> , MVST, & <i>S. Infantis</i>	Pigs ($n = 5$), broiler ($n = 8$), pigeon, turkey, and laying hen ($n = 1$ each). Pork ($n = 20$), bovine ($n = 4$), & chicken meat ($n = 7$), mollusks ($n = 4$). Food processing ($n = 2$) & slaughter rooms ($n = 4$) & poultry farms ($n = 3$)	AMP, CFX, CZA, AMC, CEX, MEM, TET, CHL, CIP, NAL, PFX, GEN, STR, STX, SMZ, TMP	INN (86%), TET, AMP, STR, NAL	Kirby-Bauer disk-diffusion	[158]
Burkina Faso	<i>Salmonella</i> spp.	Sesames ($n = 480$), mango juices ($n = 156$), lettuces ($n = 92$), salads ($n = 324$)	AMC, AMP, CHL, STX, NAL, TET, IMP, MEM, GEN, AMK, CIP, CFP, CZA, CFX, ATM	GEN (26.8%), AMP, CEX, NAL	Kirby-Bauer disk-diffusion	[70]
Bangladesh	<i>S. Enteritidis</i> , <i>S. Typhimurium</i>	Carcass dressing water, chopping board, & knife swabs ($n = 290$)	AMK, GEN, STR, MEM, CTX, CFX, CZA, ATM, AMC, AMP, AZM, CIP, NAL, STX, TET, CHL	CIP (68.9%), NAL, TET, AMP, STR	Kirby-Bauer disk-diffusion	[159]
Poland	<i>Salmonella</i> spp.	Food chain links	AMP, SAM, AMC, PIP, TZP, CTX, CFP, CZA, IMP, MEM, ATM, CIP, PFX, LFX, OFX, NOR, AMK, GEN, TOB, CHL, STX	AMK (58.5%), GEN, TOB	Kirby-Bauer disk-diffusion	[73]
China	<i>S. Derby</i> , <i>S. Meleagridis</i> , <i>S. Enteritidis</i> , <i>S. Senftenberg</i>	RTE food samples ($n = 74$ cooked meat, $n = 175$ cooked poultry, $n = 30$ cold vegetable dishes, $n = 21$ fried rice/noodles)	AMC, AMP, CEZ, CEX, CZA, CTX, CFX, CFT, CFP, IMP, ATM, SRT, GEN, AMK, KAN, TET, NAL, CIP, ENR, STX, FLO, CHL	TET (70%), AMP, STR, STX	Kirby-Bauer disk-diffusion	[91]

Ethiopia	<i>Salmonella</i> spp.	Lettuce, cabbage, spinach, carrots, tomatoes, & green peppers (<i>n</i> = 30 each).	AMC, AMP, PEN, STX, CIP, CHL, GEN, ERY, TET, DOX, CEX, CTX, IMP, MEM, CFX, CZA	AMC & AMP (87.5%), STX, GEN, ERY, TET, DOX, CEX	Kirby-Bauer disk-diffusion	[78]
Mexico	<i>S. Montevideo</i> , <i>S. Meleagridis</i>	Calves, periparturient cow feces, & maternity beds (<i>n</i> = 377)	AMP, CIP, FLX, AMK, GEN	AMP, GEN (10%)	Kirby-Bauer disk-diffusion	[160]
Mexico	<i>S. Aantum</i> , <i>S. Adelaide</i> , <i>S. Newport</i> , <i>S. Infantis</i>	Retail ground beef (<i>n</i> = 115)	AMP, AMC, CTX, CFP, MEM, AMK, STR, CIP, AZM, TET, CHL, STX	TET (39.7%), CHL, STR, STX, AMP	Kirby-Bauer disk-diffusion	[72]
India	<i>Salmonella</i> spp.	Fresh eggs (backyard & commercial chicken & duck eggs, <i>n</i> = 60 each)	AMP, CFX, CIP, GEN, OXY	CFX (50%), AMP (39.3%)	Kirby-Bauer disk-diffusion	[161]
Italy	<i>S. Infantis</i> , <i>S. Typhimurium</i>	Poultry (<i>n</i> = 145) & pig meat (<i>n</i> = 106), beef (<i>n</i> = 54), bivalve molluscs (<i>n</i> = 109), eggs (<i>n</i> = 43) & sprouted seeds (<i>n</i> = 36).	KAN, GEN, STR, TOB, AMP, AMC, CFX, IMP, NAL, CIP, ENR, CTX, CZA, LFX, STX, TET, CHL	TET (50.7%), NAL, STX, KAN	Kirby-Bauer disk-diffusion	[88]
Vietnam	<i>Salmonella</i> spp.	Poultry fecal samples (<i>n</i> = 200), house floors (<i>n</i> = 150), drinking water (<i>n</i> = 45), feed (<i>n</i> = 45), sewages (<i>n</i> = 45) & tools (<i>n</i> = 30)	AMP, CIP, CZA, GEN, NAL, NFT, NOR, STR, TMP, TET	STR (89.2%), TET, AMP, NAL	Kirby-Bauer disk-diffusion	[162]
Chile	<i>S. Infantis</i> & <i>S. Heidelberg</i>	Poultry meat from the production line	AMP, CFP, CZA, CTX, CIP, GEN, AMK, IMP, MEM, ETP, TET, CVI, TZP, STX, SAM, NFT, CHL, ATM	GEN, KAN, STR, AMP, CEX, CTX, CHL, TET, STX	Kirby-Bauer disk-diffusion	[47]
Costa Rica	<i>S. Infantis</i> , <i>S. Anatum</i> , & <i>S. Kentucky</i>	Poultry meat (<i>n</i> = 38), Broiler caecum (<i>n</i> = 63)	AMK, GEN, AMP, CEX, CFX, CFP, ATM, AMP, AMC, TZP, CHL, AZM, NAL, CIP	NAL (92.1%), CHL, AMP, CFX	Kirby-Bauer disk-diffusion	[95]
Kenya	<i>Salmonella</i> spp.	Fresh Nile tilapia (<i>Oreochromis niloticus</i>) fish	PEN, VAN, RIF, AMP, CFP, CPD, CHL, NFT, CTX, MEM, STR	PEN (22.2%)	Kirby-Bauer disk-diffusion	[163]
Bhutan	<i>S. Typhimurium</i> , <i>S. Paratyphi</i>	Healthy broiler carcasses (<i>n</i> = 180) from commercial farms	AMP, AMO, STR, STX, GEN	TET (95.7%), STX, AMO, AMP	Kirby-Bauer disk-diffusion	[164]
China	<i>S. Enteritidis</i> & <i>S. Typhimurium</i>	Pork (<i>n</i> = 140), chicken (<i>n</i> = 128), vegetables (<i>n</i> = 232)	CIP, GEN, STR, ERY, CFX, AMP, AMO, ENR, KAN, SMZ, CEP, TET	TET (85.7%), CIP (71.4%)	Kirby-Bauer disk-diffusion	[92]

Morocco	<i>S. Kentucky</i> , <i>S. Muenster</i> , <i>S. Typhimurium</i> , <i>S. Menston</i>	Miscellaneous foods	AMP, CFX, CEX, CZA, CTX, CHL, CIP, GEN, ERT, NAL, TET, TMP, STX	TET (46.7%)	Kirby-Bauer disk-diffusion	[165]
Spain	<i>S. Derby</i>	Human clinical samples ($n = 20$), pig carcasses, pork- or pork & beef-derived products, or wild boar ($n = 16$)	AMP, AMC, CFP, CFX, CEX, ERT, AMK, GEN, STR, NAL, CIP, KAN, ATM, NFT, TET, TOB, CHL, TMP	TET (66.7%), STR, NAL, AMP	Kirby-Bauer disk-diffusion	[166]
China	<i>S. Enteritidis</i> , <i>S. Corvallis</i> , <i>S. Indiana</i> , <i>S. Typhimurium</i> , & <i>S. Kentucky</i>	Poultry meat	TET, CHL, SMZ, AMP, CTX, CZA, IMP, GEN, NAL, CIP, ATM, COL, CEX, CEZ	TET (59.2%), NAL, CHL	Broth microdilution	[167]

¹ Numbers in round brackets next to the first antibiotic represent the percentage of resistance reported for the antibiotic for which the strains analyzed showed the most tolerance. If no number is indicated, please assume 100% resistance.

3. *Listeria* spp.

Listeria is a genus of Gram-positive, rod-shaped, facultatively anaerobic, and non-endospore-producing bacteria that acts as an intracellular parasite in mammals. Although in food analysis *L. monocytogenes* is the most targeted species, early reports have focused on other strains, such as *L. innocua*, *L. seeligeri*, and *L. welshimeri*. Though no resistance was found in the latter, 19.3% of the *L. innocua* isolates, mostly from frozen food items, were resistant to one or more antibiotics [168]. A more recent study showed that 29% of samples from butcher shops were positive for *Listeria* spp. (48, 28, and 7% for *L. innocua*, *L. seeligeri*, and *L. monocytogenes*, respectively) [169].

L. monocytogenes resistance has been attributed to conjugation (e.g., plasmid horizontal transfer from other Gram-positive bacteria such as *Enterococcus*, *Streptococcus*, and *Staphylococcus*), efflux pumps [170,171] genetic alteration, and biofilm formation (especially contamination during food packaging) [172]. An excellent primer on *Listeria*-specific mechanisms of resistance can be found in the work of Luque-Sastre and collaborators [173].

Listeria isolates are differentiated by probing for immunoreactivity to two cell-surface structures, i.e., the somatic O and flagellar H antigens. For *L. monocytogenes*, four O-serogroups (1/2, 3, 4, 7) have been recognized, which are further differentiated into serotypes 1/2a, 1/2b, 1/2c, 3a, 3b, 3c, 4a, 4b, 4c, 4d, 4e, and 7. Three of these (1/2a, 1/2b, and 4b) cause most of the human illness cases. Specifically, serotype 4b is most associated with outbreaks [174,175]. For example, in Mexico, from 300 food samples, 5.6% of recovered strains were identified as *L. monocytogenes*, of which 68.4% belonged to serogroups 4b, 4d, and 4e, and 92.1% exhibited at least four virulence genes (i.e., *actA*, *hly*, and *plcB*) [176]. A similar case was found in Romania, where *L. monocytogenes* isolates from RTE, meat, and dairy foods included 47.1 and 29.4% of the 1/2a-3a and 4b-4d-4e serotypes, respectively [177]. An additional study in this country found that 75% of *L. monocytogenes* strains recovered from RTE foods were serotype 1/2a (with regulatory factor *prfA* present in all isolates). Analysis of antimicrobial resistant genes identified tetracycline-resistant genes, particularly *tet(C)*, *tet(M)*, and *tet(K)*. The presence of *ampC* and *dfrD* was also noticeable [178].

From a total of 50 *Listeria* strains isolated from meat and dairy products, consisting of seven *L. monocytogenes* and 43 *L. innocua* strains, which showed the presence of antimicrobial resistance genes: *tet M*, *tet L*, *mef A*, *msr A*, *erm A*, *erm B*, *lnu A*, and *lnu B*. Multidrug resistance was identified in 27 *Listeria* strains, four belonging to *L. monocytogenes*. Resistance to CDA (a lincosamide) was the most common resistance phenotype, identified in 45 *Listeria* strains. Hence, the preferred first-line therapy remains AMP or PEN, often combined with GEN or RIF for synergistic effects [170]. Worryingly, recent data suggest *L. monocytogenes* strains recovered from foods exhibited 87.3% of multidrug resistance and 92.4% to β -lactams [179]. Similarly, South African isolates exhibit resistance determinants for sulfonamides (*sulI* and *sul2*, 64.4%), aminoglycosides [*strA*, *aadA*, *aph (3)-IIA*, 47.9%], and β -lactams (*bla_{TEM}*, and *bla_z*, 44.3%). For patients allergic to β -lactams, STX is the standard alternative, as *Listeria* remains broadly susceptible to it (Table 4) [170]. Additionally, in some cases, high resistance to FOS has been documented as a natural feature of the pathogen (together with NAL, FMQ, and cephalosporins) (Table 4) [179–181].

Interestingly, a longitudinal study found that significant resistance genes in *L. monocytogenes* included *fosX*, *lin*, *abc-f* (macrolide), and *tet(M)*. No increase in antimicrobial resistance determinants was observed over 11 years [182]. There is additional evidence suggesting co-selection of resistance traits in food production systems and bacterial responses to stress and disinfectants (e.g., *bcrABC*) [181]. Over eight years, acquired antimicrobial phenotypes were shown to confer resistance to tetracyclines (*tetM*), trimethoprim (*dfrD*), lincosamides (*lnuG*), macrolides (*ermB* and *mphB*), and phenicols (*fexA*). A similar trend was recently described in isolates recovered from Chile, in which 100% of the *L. monocytogenes* strains carried the *fosX*, *lin*, *norB* (fluoroquinolones), *mprF* (lipopeptides), *tetA*, and *tetC* resistance genes. In addition, the *arsBC*, *bcrBC*, and *clpL* genes were detected, which conferred resistance to stress and

disinfectants [183]. Recent research also supports these findings, with next-generation sequencing of *L. monocytogenes* showing that *fosX* determinants are ubiquitous across 58 studies and, in up to 60.38% of reports, the presence of the *lin* and *norB* genes [184].

Since *L. monocytogenes* is ubiquitous in agricultural environments, the cultivation of fresh produce predisposes it to contamination. Cold-stored dairy, meat products, fresh produce, and novel foods made thereof (e.g., hummus or guacamole) are considered high-risk foods [185].

In this scenario, the contamination of RTE foods originates either from raw ingredients (of both plant and animal origin) or, more commonly, from cross-contamination within food-processing facilities [163]. Interestingly, recent research has also demonstrated that stressors in food-processing environments (e.g., heat, acidic and alkaline pH, UV irradiation, and high osmolarity) can induce tolerance that may contribute to the predominance of *Listeria* and the development of antimicrobial resistance, even without antibiotic selection pressure [186]. Said stressors induce bacterial tolerance by triggering global stress responses that alter cellular physiology. When bacteria encounter stressors such as sublethal antimicrobials, temperature shifts, or nutrient starvation, they temporarily slow their metabolism, deploy protective chaperones, and alter membrane permeability, thereby effectively blocking antibiotics from reaching their targets [187].

For example, in carcasses obtained from poultry slaughterhouses in Northern Greece, an 80 and 99% of samples were positive for *Listeria* spp. and *L. monocytogenes*, respectively [166]. All isolates were resistant to NAL (a naphthyridone) and OXO (both used to treat Gram-negative bacteria), most were resistant to CDA, and only a few were resistant to OXY and TET [188].

Finally, although relatively recent attention has been drawn to the resistance of *Listeria* species in food, some countries, possibly compelled by listeriosis outbreaks, have already amassed considerable research on the matter [189,190]. Notably, some other African countries have also identified human breast milk as a reservoir for *Listeria* [191].

Table 4. Summary of bacterial isolates from foodstuffs and other food-related matrices for *Listeria* species.

Country	Isolates	Source	Antibiotics Tested	Resistance to ¹	Testing Method	References
United Kingdom	<i>L. monocytogenes</i>	Foods	AMP, CHL, COL, ERY, GEN, KAN, STR, SPC, SMZ, TET, TMP, VAN, FZD, NAL, CIP	TET (2–5%), CIP (2.2%), <i>Listeria</i> spp. are constitutively resistant to NAL, COL & cephalosporins	Agar dilution Breakpoint	[192]
Ireland	<i>L. monocytogenes</i> , <i>L. innocua</i> , <i>L. seeligeri</i> , <i>L. welshimeri</i> .	<i>n</i> = 67 retail food samples	AMP, CHL, ERY, GEN, PEN, STR, TET, VAN	TET (6.3%), PEN (3.7%), AMP (2.0%)	Kirby-Bauer disk-diffusion	[168]
Greece	<i>L. monocytogenes</i>	Poultry slaughterhouses (<i>n</i> = 4 establishments; 100 samples)	AMO, AMP, CFX, CEP, CHL, CIP, CDA, ENR, ERY, GEN, KAN, NAL, NEO, OXO, OXY, PEN	OXO & NAL (100%), CDA (83.6%), OXY (12.7%)	Kirby-Bauer disk diffusion	[188]
Türkiye	<i>L. monocytogenes</i>	Raw or cooked foods, including sausage, raw minced beef, chicken breast, chicken gizzard, raw meat, raw milk, cooked turkey döner, anchovy, and mussel	PEN, VAN, TET, CHL, RIF, ERY, GEN, TMP, FOS, STR	FOS (92.9%), STR (7.1%)	Kirby-Bauer disk diffusion	[180]

Spain	<i>L. monocytogenes</i> , <i>L. innocua</i>	RTE products of animal origin (meat and dairy products)	AMP, CIP, CDA, ERY, GEN, PEN, RIF, TET, VAN	CDA (100%), CIP (52%), AMP (32%)	Kirby-Bauer disk-diffusion [193]
South Africa	<i>L. monocytogenes</i>	Environmental, raw seafood, raw meats, & RTE foods (e.g. fresh produce, dairy, deli meats), <i>n</i> = 177 isolates	AMP, CHL, ERY, GEN, TET	RTE foods 25% at least to <i>n</i> = 2 (CHL, ERY, TET) antibiotics, raw meats 50% at least to <i>n</i> = 4 antibiotics	Kirby-Bauer disk-diffusion [189]
South Africa	<i>L. monocytogenes</i>	RTE foods (<i>n</i> = 194 isolates)	FOS, CHL, OXY, CIP, VAN, STR, AMK, GEN, STX, TRP, CAM, CTT, CTX, SMX, ERT, MEM, ERY, IMP, AMO, SAM, AMP, PEN	CTX (53.1%), TMP (56.2%), SMZ & CTT (61.9%), VAN & OXY (62.9%)	Kirby-Bauer disk diffusion [194]
Chile	<i>L. monocytogenes</i>	RTE foods (<i>n</i> = 436). Cheeses (<i>n</i> = 161), cooked meats (artisanal ham, pâté, sausages, & blood sausage) (<i>n</i> = 186), pre-processed fruit & vegetables (chopped fruit, fruit salads with strawberries, melon, & peaches, & leafy vegetable salads) (<i>n</i> = 22), & meals and mixed dishes with raw & cooked ingredients (<i>n</i> = 67)	AMP, PEN, STX, ERY, VAN, TET, CIP, CHL	AMP (78% sensitivity)	Kirby-Bauer disk diffusion [183]
Romania	<i>L. monocytogenes</i>	<i>n</i> = 130 RTE products (ham sausages, smoked specialties), <i>n</i> = 47 raw meat samples (minced pork & pork organs, snails), <i>n</i> = 44 dairy samples (assorted cheese)	PEN, IMP, GEN, CIP, MFX, FOS, TEL, TIG, VAN, LIN, CDA, ERY, LZD, FAS, RIF, STX, TET	PEN & FAS (100%), MFX (88.2%), FOS (82.4%), CDA (76.5%), IMP (52.9%)	Vitek 2 Compact automated system [177]
France	<i>L. monocytogenes</i>	Food isolates (<i>n</i> = 2424)	AMO, AMP, CTX, CIP, CDA, CHL, ERY, FOS, FAS, GEN, IMP, TET, LVX, MXF, NAL, KAN, PEN, RIF, STR, TMP, VAN	NAL (100%), CTX (99%), FOS (96%)	E-test [181]
South Africa	<i>Listeria</i> spp. (<i>L. monocytogenes</i> , <i>L. innocua</i> , & <i>L. welshimeri</i>)	AMC, PEN, AMP, CFX, STR, GEN, KAN, TET, DOX, NAL, CIP, ENR, AZM, CDA, STX	NAL 100%, CFX & CDA 66.7%, TET 33.3%, AMC 22.2%	NAL (100%), CFX (80.4%)	Kirby-Bauer disk-diffusion [195]
Mexico	<i>L. monocytogenes</i>	Raw poultry (<i>n</i> = 60), ground beef (<i>n</i> = 60), seafood (<i>n</i> = 20 fish, <i>n</i> = 20 shrimp, & <i>n</i> = 20 crabs), cheese (<i>n</i> = 30 fresh cheese & <i>n</i> = 30 mature cheese), & milk (<i>n</i> = 30 pasteurized milk and <i>n</i> = 30 raw milk)	AMP, CHL, CIP, ERY, GEN, LFX, MEM, STR, STX/TMP, TET, VAN	STX (88.4%), STR & MEM (30.7%)	Kirby-Bauer disk-diffusion [176]

Ethiopia	<i>L. monocytogenes</i>	Bovine milk from dairy farms ($n = 166$), shops ($n = 120$), & cafeterias ($n = 98$). Bovine meat from abattoirs ($n = 240$), butcher shops ($n = 84$), & restaurants ($n = 60$)	PEN, AMO, AMP, VAN, CIP, STR, NAL, GEN, ERY, CDA, CLX, TET	NAL (96.1%), TET (53.8%), STR (15.4%), PEN (11.5%)	Kirby-Bauer disk-diffusion	[196]
Romania	<i>L. monocytogenes</i>	RTE foods ($n = 8151$)	AMP, CIP, CDA, CHL, GEN, OXA, CEP, LVX, MXF, PEN, RIF, STM, TET, VAN	STM (26.9%), OXA (23.1%)	Kirby-Bauer disk-diffusion	[178]
Spain	<i>L. monocytogenes</i>	$n = 104$, $n = 52$ carcasses & $n = 52$ chicken cuts	AMP, OXA, CEX, CFX, CFP, GEN, ERY, VAN, SXT, RIF, TET, CHL, CIP, ENR, NFT	OXA + CEX, CTX + CEP (48.7%)	Kirby-Bauer disk-diffusion	[197]
Italy	<i>L. monocytogenes</i>	Meat products ($n = 173$), seafood ($n = 54$), dairy products ($n = 19$), sauces ($n = 2$), confectionery products ($n = 1$), RTE rice dishes ($n = 1$), & food-processing environments ($n = 19$).	AMP, LIN, OXA, FMQ, FOS, GEN, MER, ERY, STX, TET	OXA (88.48%), FOS (85.87%) & FMQ (78.44%).	Kirby-Bauer disk-diffusion	[179]
Ethiopia	<i>L. monocytogene</i> , <i>L. innocua</i> , & <i>L. seeligeri</i>	Raw meat from abattoirs & butcher shops	AMO, CHL, CLX, PEN, TET, VAN, SMZ, GEN, NAL, AMP	CLX & NAL (100%), PEN (76%), TET (59%), CHL (24%)	Kirby-Bauer disk-diffusion	[169]
South Africa	<i>L. monocytogenes</i> ($n = 24$) & <i>Listeria</i> spp. ($n = 122$)	Beef and beef products	TET, STX, STR, PEN, NAL, KAN, ERN, DOX, CDA, CEX, GEN, CIP, AZM, AMP, AMC	NAL (100%), STR (45.8%), ENR (41.7%), DOX (37.5%), AMP & ATM (33.3%)	Kirby-Bauer disk-diffusion	[198]
India	<i>L. monocytogenes</i>	Local ethnic foods ($n = 2829$). Raw meats, cooked foods, fermented products, dairy, street foods, & beverages	AMP, PEN, SXT, MEM, CHL, CIP, GEN, TET, ERY, CDA, VAN	CDA (48%), ERY (45%), AMP (42%), PEN (38%)	Microbroth dilution technique	[171]
Romania	<i>L. monocytogenes</i>	$n = 10,306$ samples (<i>i.e.</i> , milk, milk-based beverages, cheese, cream, butter, yogurt, ice cream, dairy desserts, milk, and whey powders)	AMP, PEN, OXA, MET, CIP, LFX, MFX, CDA, TET, GEN, CHL, RIF, SMZ	OXA & SMZ (13.95%), AMP, TET (11.62%)	Kirby-Bauer disk-diffusion	[199]
Poland	<i>Listeria</i> spp.	$n = 750$ raw fish samples (Crucian carp, Sturgeon, Trout, Herring)	OXA, PEN, MER, ERY, RIF, VAN, TET, SMZ, CHL, CIP, ENR, NIT, CDA	OXA, CEF, MER, RIF, SMZ (100%), CIP (91.7%), TET (75.0%)	Kirby-Bauer disk-diffusion	[200]

¹ Numbers in round brackets next to the first antibiotic represent the percentage of resistance reported for the antibiotic for which the strains analyzed showed the most tolerance. If no number is indicated, please assume 100% resistance.

4. *Campylobacter* spp.

Campylobacter spp. are Gram-negative spiral, non-spore-forming, chemoorganotrophic, rod-shaped, or curved bacteria. These bacteria are a leading cause of foodborne gastroenteritis [201,202]. The advent of molecular detection of genes responsible for antibiotic resistance (e.g., *gyr* for fluoroquinolones, *erm* for macrolides, *tet* for tetracyclines, or *aph* for aminoglycosides) has facilitated clinical therapy and surveillance of the emergence and dissemination of antibiotic-resistant bacteria [203]. For an excellent

primer regarding *Campylobacter* resistance mechanisms (including fluoroquinolones, macrolides, FLO, β -lactams, aminoglycosides, and tetracyclines, the work by Iovine, Shen et al., Tang et al., and, more recently, Aleksic et al., are proposed [204–207].

In a very thorough work regarding contamination in a slaughterhouse, Schreyer et al. [208] found not only *Campylobacter* spp. in chicken cecum but also evisceration knives and workers' hands, and samples before and after the chiller. Interestingly, *C. coli* contamination was significantly higher than that of *C. jejuni*. This trait has been observed in analogous (poultry-related) studies in other latitudes [209,210]. Not only was the contamination within the meat chain relatively high, but the strains also exhibited several genes responsible for resistance, such as *cmeA/B*, *tet(O)*, and *bla_{OXA-6}*.

Efflux pump CmeABC expels substances such as antimicrobials, toxic heavy metals, dyes, and detergents from the interior of the bacterial cell [211]. Interestingly, it has been correlated with surface polysaccharides, which serve as a resistance mechanism against phenolic compounds [212]. It is also largely responsible for macrolide [213] and multidrug resistance in Gram-negative bacteria [211,214], and its role in the clinical context is well defined, as it contributes significantly to high-level resistance to CIP and TET in *C. jejuni* and *C. coli* [215].

Additionally, the *tet(O)* determinant is a common trait and has been described in similar studies [216,217]. Such evidence adds to the previous body of research, which heavily linked fluoroquinolone-resistant *Campylobacter* spread from poultry (since *Campylobacter* is a normal enteric flora in many avian species) to humans [209,218,219]. However, as a zoonotic infection, it can be present in domestic and wild birds and other animals [220]. Hence, campylobacteriosis warrants a One Health approach to tackling this problem [221].

Alike results have also been obtained from pig slaughterhouses. *Campylobacter* spp. have been isolated from pig guts (200 samples, 47.6% contaminated), feces, and surfaces. 90.8% of strains exhibited multidrug resistance [222]. A general increase in *Campylobacter* resistance toward fluoroquinolones (e.g., NAL/CIP) and TET has been reported worldwide [220,223–225]. EFSA reported that rates of resistance to CIP were very high in *C. jejuni* and *C. coli* isolates from humans and very high to extremely high in *C. jejuni* and *C. coli* isolates from poultry, pigs, and calves [224]. A recent review that reported South American *Campylobacter* isolates indicated that although the bacteria showed an overall low resistance to ERY (a drug of choice to treat campylobacteriosis), it was widespread [225], a trait that persists in Europe [224]. On the other hand, in general, cattle and chickens seem to be important hosts in the Sub-Saharan African region, and may play an important role in transmitting to humans. Most *Campylobacter* isolates were resistant to ERY (44%), AMP (39%), TET (33%), NAL (31%), and CIP (30%) [226].

A recent study [227] found multidrug resistance in almost all isolates from cattle, chicken, and water samples, and also determined that the most resistance determinant within the strains was the *tet(O)* gene. Despite several restrictions in some countries, tetracyclines are still among the most systematically used as growth promoters in animal feed, which maintains selective pressure over gut microbiota [228–231]. A recent Canadian study [232] found that, even in 2021, the most used antibiotic in turkey farming includes bacitracin (at 472.7 mg·kg⁻¹ per biomass), followed by β -lactams, orthomycin (a combination of isoniazid and a streptomycin derivative) [233], and tetracyclines (at 58.8 mg·kg⁻¹ per biomass). Other genes coding for resistance were those for β -lactams (*bla_{OXA-61}*), aminoglycosides (*aph-3-I*), (fluoro)quinolones (*gyrA*), and multidrug efflux pump (*cmeB*). Tunisian laying hens and eggs have been recently found to harbor similar resistance determinants [234]. The authors also found the *erm(B)* gene in almost all the isolates from these samples. Byambajav et al. also found the *erm(B)* gene or a nucleotide substitution at nt 2075 of 23S rDNA in all *C. coli* isolates from Mongolia [235].

An interesting finding that supports the (over)use of antibiotics in livestock as a major driver of the selection of resistant bacteria is reported by Paintsil et al. [210]. The authors found that fecal samples from

animals on smallholder farms exhibited not only significantly lower *Campylobacter* contamination but also reduced antibiotic resistance.

Notably, a Belgian study [236] showed that human isolates (from diarrheal cases) had resistance profiles similar to those found in animals, with a tendency toward increased resistance to NAL, CIP, and TET. Whole genome sequencing of human isolates from the USA [237] found antimicrobial resistance genes encoding mostly resistance to the β -lactam (*bla*_{OXA-193}, *bla*_{OXA-460}), tetracyclines [*tet*(O), *tet*(O/32/O)], fluoroquinolones (*gyrA*), and aminoglycosides (*aph*(3')-III). Another African study reports similar results in pediatric cases with clinical diarrhea, supporting the transfer of genetic determinants of antibiotic resistance from animals to humans [238].

In conclusion, evidence indicates that the relatively high rates of fluoroquinolone resistance (Table 5) have limited their usefulness in treating *Campylobacter* infections. Macrolides like azithromycin are the current drugs of choice when antibiotic treatment is indicated. Resistance to macrolides in *Campylobacter* has remained stable. The increased prevalence of antibiotic-resistant *Campylobacter* cannot be solely attributed to antibiotic overuse. Interspecies transmission and subsequent clonal expansion also contribute to the dissemination [221].

There seems to be a certain plasticity in *Campylobacter* species, and genetic material from *C. jejuni* to *C. coli* is a possibility, which then increases virulence [234]. The result is a higher number of antibiotic-resistant infections, which could become a health burden. Furthermore, genetic exchange between *Campylobacter* and Gram-positive bacteria occurs in the natural environment and is more frequent than previously realized. These antibiotic-resistance genes present in *Campylobacter* species can further disseminate by clonal expansion or horizontal gene transfer [221].

Table 5. Summary of *Campylobacter* bacterial isolates from foodstuffs and other food-related matrices.

Country	Isolates	Source	Antibiotics Tested	Resistance to ¹	Testing Method	References
Poland	<i>C. jejuni</i> & <i>C. coli</i>	Chicken meat and giblets (<i>n</i> = 218)	CIP, ERY, TET, GEN	CIP (97.9%)	Kirby-Bauer disk-diffusion	[209]
Poland	<i>C. jejuni</i> (<i>n</i> = 132) & <i>C. coli</i> (<i>n</i> = 189)	Retail raw meat beef (<i>n</i> = 105), pork (<i>n</i> = 85), poultry (<i>n</i> = 368)	CIP, NAL, STR, TET, ERY	CIP (91% <i>C. jejuni</i> ; 86.1% <i>C. coli</i>) & NAL (89.3% <i>C. jejuni</i> ; % <i>C. coli</i>)	Kirby-Bauer disk-diffusion	[239]
France	<i>C. jejuni</i> (<i>n</i> = 175)	Chicken meat products: carcasses (<i>n</i> = 120), legs (<i>n</i> = 121) & fillets (<i>n</i> = 120)	GEN, STR, CIP, NAL, TET, ERY, CHL		Broth microdilution	[240]
Nigeria	<i>C. jejuni</i> (<i>n</i> = 242)	Retail chicken (<i>n</i> = 252)	NAL, GEN, ERY, ENR, CHL, STR, TET	NAL (100%), GEN, ERY, ENR, CHL, STR, TET	Kirby-Bauer disk-diffusion	[241]
Canada	<i>C. jejuni</i> (<i>n</i> = 53) & <i>C. coli</i> (<i>n</i> = 3); (23.5 & 14.2% from chicken and turkey samples)	Retail chicken (<i>n</i> = 204), turkey (<i>n</i> = 110), ground beef (<i>n</i> = 145) & pork (<i>n</i> = 147)	TMY, AZM, CDA, ERY, GEN, NAL, FLO, TET	TET (48.1%)	Broth microdilution	[216]
United Kingdom	<i>C. jejuni</i> & <i>C. coli</i> (<i>n</i> = 393)	Chilled chickens at retail (skin samples)	CIP, ERY, TET, GEN, STR	CIP (52% <i>C. jejuni</i> ; 48% <i>C. coli</i>) & TET (52% <i>C. jejuni</i> ; 60% <i>C. coli</i>)	Broth microdilution	[242]

Egypt	<i>Campylobacter</i> spp. (n = 98)/ <i>C. jejuni</i> (n = 65)	Chicken (n = 315), Animal Products (n = 122) & Human Stool Swabs (n = 128)	AMP, CEP, ERY, GEN, NAL, OXY	CEP (94.7%), TET, ERY, NAL	Kirby-Bauer disk-diffusion	[243]
Canada	<i>C. jejuni</i> & <i>C. coli</i> (n = 178)	Chicken, Turkey, Duck, Game Birds	GEN, TMY, CDA, AZM, ERY, FLO, CIP, NAL, TET	TET (56.25%)	Broth microdilution	[244]
Mongolia	<i>C. jejuni</i> (n = 25) & <i>C. coli</i> (n = 6)	Cloacal Swabs of Chickens (Broiler/Laying, n = 71)	CIP, ERY, NAL, NOR, TET	NAL (51.6%)	E-Tests	[235]
Russia	<i>C. jejuni</i> (n = 40)	Raw milk, poultry products, and washings from processing equipment	ERY, AZM, CDA, CAM	ERY (27.5%)	Kirby-Bauer disk-diffusion	[213]
Brazil	<i>C. jejuni</i> (n = 54)	Broiler carcass/Poultry slaughterhouses	CHL, CIP, ERY, GEN, NAL, TET	TET (51.8%), ERY	Broth microdilution	[217]
Japan	<i>C. jejuni</i> (n = 13) & <i>C. coli</i> (n = 3)	Pork (n = 14) & chicken (n = 53)	AMP, CEZ, STX, IPM, MIN, GEN, KAN, ERY, AZM, NAL, CIP, LFX	CEZ, STX, NAL, CIP, LFX	Kirby-Bauer disk-diffusion	[245]
Benin	<i>C. jejuni</i> & <i>C. coli</i>	Cutting Tables at Swine Slaughterhouse and Taverns	AMP, GEN, ERY, CIP, TET, AMC	CIP (44%), AMP, ERY, GEN, AMC	Kirby-Bauer disk-diffusion	[222]
Italy	<i>C. jejuni</i> (n = 2734)	Domestic/wild animals & humans	NAL, CIP, CHL, ERY, GEN, STR, TET	NAL (67.39%), CIP, TET	Broth microdilution	[246]
Tunisia	<i>C. jejuni</i> & <i>C. coli</i> (n = 177)	Layer hens and eggs	AMP, AMC, GEN, TET, ERY, NAL, CIP, CHL	ERY (98.87%) & TET	Kirby-Bauer disk-diffusion	[234]
Malaysia	<i>C. jejuni</i> & <i>E. coli</i>	Wild birds (n = 38), chickens (n = 71), humans (n = 47), & environment (n = 153)	CIP, ENR, SAM, TET, GEN, STR, ERY, NAL, SXT, CPD	CPD & TET	Kirby-Bauer disk-diffusion	[247]
Ghana	<i>C. jejuni</i> & <i>C. coli</i>	Livestock fecal samples from commercial & smallholder farms	CIP, TET, STR, AMP, KAN, ERY, CHL	CIP (41.3% <i>C. jejuni</i> ; 58.7% <i>C. coli</i>) & TET (52.9% <i>C. jejuni</i> ; 43.6% <i>C. coli</i>)	Kirby-Bauer disk-diffusion	[236]
Argentina	<i>C. jejuni</i> & <i>C. coli</i>	Poultry slaughterhouse (n = 333)	CIP, ERY, TET, ENR, GEN, AMP, CHL, STR	<i>C. jejuni</i> : CIP, TET, ENR. <i>C. coli</i> : CIP, ERY, TET, ENR, AMP	Kirby-Bauer disk-diffusion	[208]
Estonia/Latvia/Lithuania	<i>C. jejuni</i> (n = 46) & <i>C. coli</i> (n = 15)	Fresh broiler chicken meat and human patients	ERY, CIP, STR, NAL, GEN	NAL (100–86.7%), CIP, TET, STR (68.8–26.7%)	Broth microdilution	[248]
Pakistan	<i>Campylobacter</i> spp. (n = 38/6)	Chicken, beef & mutton (n = 300)	NAL, CFX, CDA, TYL, ENR, STR, DOX, TET	TET (100%), CFX, TYL, ENR, CDA, NAL	Kirby-Bauer disk-diffusion	[249]

Canada	<i>C. jejuni</i> (n = 871) & <i>C. coli</i> (n = 383)	Turkeys (fecal samples, n = 293 flocks)	CIP, AZM, ERY, GEN, NAL, TET, CIP, NAL, TET TYM, CDA, FLO	Broth microdilution [232]
Kenya	<i>C. jejuni</i> (n = 74) & <i>C. coli</i> (n = 29)	Chicken, cattle, and cattle-trough water samples	AMP, TET, GEN, ERY, CIP, AMP (100%), NAL TET, ERY, CIP	Kirby-Bauer disk-diffusion [227]

¹ Numbers in round brackets next to the first antibiotic represent the percentage of resistance reported for the antibiotic for which the strains analyzed showed the most tolerance. If no number is indicated, please assume 100% resistance.

5. *Staphylococcus* spp.

Staphylococci are clump-forming Gram-positive cocci of 1 µm in diameter. They are salt-tolerant and often hemolytic. *S. aureus* and *S. intermedius* are coagulase-positive. All other staphylococci are coagulase negative. Staphylococcal food poisoning is a rapid-onset illness caused by eating foods contaminated with *S. aureus*, often transmitted by food handlers or through dairy/meat products.

An early report on whey and cheese showed that staphylococci coagulase-negative (CoNS, 54% of the isolates) included species such as *S. lentus*. Coagulase-positive staphylococci identified were divided into 3 groups: *S. aureus* (40%), *S. intermedius* (2%), and *S. hyicus* (4%), indicating a high incidence of PEN resistance. *S. lentus* isolates exhibited resistance to OXA, ERY, and LIN [250].

More recently, Osman et al. [251] found *S. hyicus*, *S. aureus*, *S. schleiferi* subsp. *coagulans*, *S. intermedius*, and *S. lentus* in beef meat. These strains were characterized by possessing genes *gyrA*, *gyrB*, and *griA* (encoding essential type II/IV topoisomerases in bacteria that regulate DNA supercoiling during replication and transcription). Though phenotypically already resistant to OXA, only 40.7 and 7.4% carried the *mecA* and *cfr* genes, respectively. The chloramphenicol-florfenicol resistance gene is a mobile genetic element that encodes a 23S rRNA methyltransferase, conferring broad-spectrum, multidrug resistance to Phenicol, Lincosamides, Oxazolidinones, Pleuromutilins, and Streptogramin A [251].

Although *S. aureus* is mostly considered a clinically relevant pathogen [252], several reports have contributed to establishing its incidence in food samples, food handlers, food-producing animals, and food processing environments worldwide (e.g., Brazil [253,254], Portugal [255], China [256], Italy [257]). Of special interest are reports that include the presence of Methicillin-Resistant *Staphylococcus aureus* (or MRSA) in foods. MRSA (resistance primarily driven by the *mecA* gene, which encodes a low-affinity penicillin-binding protein [PBP2a]) is one of the leading causes of infections and a major threat to human health worldwide due to its numerous toxins [252,258].

One particularity among food *Staphylococcal* contamination is the specific attention given to direct handling (Table 6) [259]. *S. aureus* is known to easily colonize the hands and nose. In fact, among health professionals, carriage of *S. aureus* on hands and nose was found to be 8.9 and 39.6% (4.7 and 17.2% for MRSA), respectively. In addition to β-lactams, the majority of nasal MRSA were also resistant to ERY, and CIP [260].

Mekhloufi et al. [261] found that the *S. aureus* strains recovered from foods belonged to seven different staphylococcal enterotoxin gene (SEg)-types harboring the following combination of *se* and *sel* genes: (1) *seI*W, *seI*X; (2) *egc* (*seG*, *seI*, *seM*, *seN*, *seO*), *seI*W, *seI*X; (3) *seA*, *seH*, *seK*, *seQ*, *seI*W, *seI*X; (4) *seB*, *seI*W, *seI*X; (5) *seD*, *seI*J, *seR*, *seI*W, *seI*X; (6) *seH*, *seI*W, *seI*X, *seI*Y; and (7) *seA*, *egc* (*seG*, *seI*, *seM*, *seN*, *seO*), *seI*W, *seI*X. Meanwhile, 2.1% and 4.2% tested positive for the *tst*- (toxic shock syndrome toxin-1) and *mecA*- (staphylococcal chromosomal cassette *mec*-type IV) genes, respectively.

Interestingly, Elsalkh et al. [262] found that 48.6 and 8.6% of isolates recovered from pizza were identified as methicillin- and vancomycin-resistant *S. aureus* through detection of *mecA* and *vanA* genes, respectively.

In yet another report [263], the *hla* (encoding an α-hemolysin) gene was detected in 87.1% of the isolated strains. Antimicrobial treatments heavily influence the *hla* gene; while some antibiotics (like β-

lactams) can inadvertently trigger its expression, others (like CDA) actively suppress toxin production to help limit tissue damage [263].

Analogously, do Nascimento Pereira et al. [264] evaluated *S. aureus* strains isolated from food handlers of pilot kitchens. In this case, 83.8 and 18.9% of isolates exhibited *seA* and *tst* genes, respectively. Also, *agr* (the accessory gene regulator locus is a critical quorum-sensing system in *S. aureus* that controls virulence factor expression based on cell density). Typing classified 58.3% of the strains as type I, and 18.9% were classified as MRSA. The SCCmec typing characterized strains as type II, III, IV, and V. For a thorough description of the staphylococcal cassette chromosome *mec* (SCCmec) typing, please refer to Uehara [265].

Achek et al. [266] found in both food and clinical isolates, the *tetM/K*, *blaZ*, *aacA-aphD*, *ermC*, and *mecA* genes. More recently, Anas et al. [267] found that the prevalence of *Staphylococcus* was very high (75 and 50%) in street kebab and raw buffalo meat, respectively. These isolates (82%) showed not only multidrug resistance (MDR) patterns but also produced β -lactamases or extended-spectrum β -lactamases, at rates of 48% and 24%, respectively. The most detected resistance genes were *mecA*, *sul1*, and *qnrS/qnrB* (29.2, 25, 20.8%, respectively). Similarly, Urimi et al. [268] found that 75% of *S. aureus* isolates from fast food restaurants were MDR, but their indices of multiple antibiotic resistance (MAR) ranged from 0.2 to 0.6. For a comprehensive list of resistance genes found in *Staphylococcus*, food, and food-related environments, the work by Chieffi et al. [257] is suggested.

Some researchers have used specific antibiotics against *Staphylococci* species, such as MCP (effective against topical infections caused by MRSA and *S. pyogenes*) [268], PRI (to treat severe Gram-positive infections, including MRSA) [250,261], or CRN (used to treat acute MRSA skin and skin structure infections) to test sensitivity [267]. Notably, from these three antimicrobials, resistance has already been proven towards MCP [269–271].

Finally, several approaches have been suggested as alternatives to antimicrobials for controlling *S. aureus* disease (e.g., bacteriophages, plant-based antimicrobials, nanoparticle- or light-based therapeutics, and probiotics) [272].

Table 6. Summary of *Staphylococcus* bacterial isolates from foodstuffs and other food-related matrices.

Country	Isolates	Source	Antibiotics Tested	Resistance to ¹	Testing Method	References
Spain	<i>Staphylococcus aureus</i>	Foods (<i>n</i> = 50 isolates) & Food handlers (<i>n</i> = 14)	AMP, PEN, OXA, MET, GEN, AMK, KAN, MCP, TMP, TET, ERY, CDA, CHL, CIP, MFX, RIF, LZD, VAN, TIG, TMP, STX	AMP & PEN (61.3%), ERY (25.8%) MCP (19.4%), OXA & MET disk-diffusion (6.5%)	Kirby-Bauer disk-diffusion	[269]
Egypt	<i>Staphylococcus</i> spp. & coagulase-negative staphylococci	<i>n</i> = 30 each, fresh normal cow milk, pasteurized milk, pasteurized yogurt, beef burger & minced beef meat	CIP, ERY, TET, MET, RIF, PEN, VAN	PEN, MET, & TET (33.3%)	Kirby-Bauer disk-diffusion	[273]
Egypt	<i>S. hyicus</i> , <i>S. aureus</i> , <i>S. schleiferi</i> subsp. <i>coagulans</i> , <i>S. intermedius</i> , & <i>S. lentus</i>	<i>n</i> = 100 retail whole beef	MET, PEN, SAM, OXA, CHL, CIP, CDA, GEN, ERY, STX, TET, VAN	ERY (48.1%), CIP (59.3%), MET (63%), TET (66.7%), VAN (74.1%), OXA (77.8%), PEN (92.6%), CDA (96.3%), GEN (96.3%), STX (100%)	Kirby-Bauer disk-diffusion	[251]

Algeria	<i>Staphylococcus aureus</i> & MRSA	Raw milk ($n = 30$), minced beef meat ($n = 25$), chicken meat ($n = 18$), creamy cake ($n = 14$), pizza ($n = 10$), beef meat ($n = 10$) & sausages ($n = 5$)	PEN, OXA, CEX, AMC, GEN, ERY, KAN, TET, VAN, CDA, RIF, STX	PEN (100%), TET (45%)	Kirby-Bauer disk-diffusion/E-test	[266]
Türkiye	<i>Staphylococcus aureus</i>	$n = 387$ Cheese samples	PEN, OXA, AMP, MET, LZD, KAN, STR, GEN, NTM, VAN, TEI, ERY, ATM, RIF, CDA, LIN, TET, DOX, CEX, CEZ, CFP, CHL, TMP, FZD, SMZ, CIP, ENR, NOR, OFX	PEN (84.7%), AMP (72.9%), OXA (63.5%), LIN (57.7%), CEX (51.8%), CDA (50.6%), GEN (31.8%), MET (30.6%)	Kirby-Bauer disk-diffusion	[274]
Morocco	<i>Staphylococcus</i> spp.	Hospital food, work surfaces & food handlers. $n = 608$ samples: 300 food, 238 food contact surfaces & 70 nasal & hand	PEN, CEX, OXA, GEN, TMP, CIP, NOR, OFX, STX, FAS	OXA & PEN (100%)	Kirby-Bauer disk-diffusion	[275]
Algeria	<i>Staphylococcus aureus</i> & MRSA	$n = 207$ RTE foods (e.g., pastries $n = 43$ & cereals $n = 17$)	PEN, OXA, CEX, GEN, KAN, TMP, OFX, ERY, LIN, CDA, PRI, LZD, TEI, VAN, TET, FOS, NTF, FAS, RIF, STX	PEN (83.3%), TET & KAN (16.7%), OFX, ERY & LIN (8.3%)	Vitek 2	[261]
Malaysia	<i>Staphylococcus aureus</i>	Food handlers & cooked food	PEN, CEX, ERY, CDA, MCP	PEN (58%)	Kirby-Bauer disk-diffusion	[271]
Bangladesh	<i>Staphylococcus</i> spp.	$n = 60$ samples of fast foods sold in different restaurants	ATM, ERY, PEN, AMO, AMP, CIP, DOX, GEN, CHL, CFX, CFM	ERY & ATM (95%), CFM (75%), AMP (50%), AMO (37.5%)	Kirby-Bauer disk-diffusion	[268]
Brazil	<i>Staphylococcus aureus</i>	$n = 41$ food handlers	PEN, OXA, CAM, ERY, CEX, TET, LFX, AMO, CDA, LZD, VAN	CAM & ERY (67.6%), OXA (64.9%), PEN (59.5%)	Kirby-Bauer disk-diffusion	[264]
Ethiopia	<i>Staphylococcus aureus</i> & <i>E. coli</i>	$n = 525$ Street Foods & $n = 175$ water samples	AMC, CHL, CIP, ERY, STR, SXT	ERY (78.6%), AMC (36.4%), STR (36.4%)—ERY (67.9% for <i>E. coli</i> strains)	Kirby-Bauer disk-diffusion	[276]
Ethiopia	<i>Staphylococcus aureus</i>	Milk, beef, eggs, and cheese, $n = 96$ each	CIP, ERY, TET, PEN, KAN, CEX, TMP, GEN, IMP, CRN	KAN (39%), TET (36%)	Kirby-Bauer disk-diffusion	[270]
India	<i>Staphylococcus</i> spp.	Raw buffalo minced meat ($n = 70$) & street meat kebabs ($n = 20$)	MET, CFC, SDZ, GTX, GMX, CFP, CZA, CTC, ATM, VAN, NAL, ERT, MEM, IMP, CFP, CTX, TET, OXA, CHL, CIP, KAN, ERY	NAL (80%), MET (68%), OXA (54%), CFP (36%)	Kirby-Bauer disk-diffusion	[267]
Poland	<i>Staphylococcus epidermis</i>	RTE foods including cheese, sushi, salad, & juice ($n = 26$ isolates)	AMK, GEN, OXA, MXF, VAN, CDA, ATM, CAM, ERY, NFT, LZD, FAS, TMP, STZ, TET	ERY (80.8%), CDA (73.1%), AZM (65.4%), CAM (61.5%)	Broth microdilution	[277]

Egypt	<i>Staphylococcus aureus</i>	<i>n</i> = 100 each, meat, chicken, & canned tuna pizzas	RIF, LZD, CIP, GEN, ATM, TET, AMP, PEN, KAN, CTX, CDA, CXM, STX, NFT	AMP, OXA & PEN (100%), CXM (87%), CTX (77%), KAN (76%)	Kirby-Bauer disk-diffusion [262]
South Africa	<i>Staphylococcus aureus</i>	<i>n</i> = 168 samples, <i>n</i> = 42 each, maize meal/pap, chicken, pork & salad	TET, PEN, CEX, ERY, GEN, CIP	PEN (52%), CEX (46%), CIP (44%), GEN (24%), TET (14%)	Kirby-Bauer disk-diffusion [278]

¹ Numbers in round brackets next to the first antibiotic represent the percentage of resistance reported for the antibiotic for which the strains analyzed showed the most tolerance. If no number is indicated, please assume 100% resistance.

6. *Escherichia coli*

E. coli is a gram-negative, facultative anaerobic, rod-shaped, coliform bacterium. As a fecal indicator, it may suggest fecal contamination or poor, unhygienic food handling. As an indicator organism of contamination, *E. coli*'s resistance profile is very relevant and has been documented (see discussion below and Table 7).

Babines-Orozco et al. [279] concluded that in Latin America, most antibiotic resistance genes present in *E. coli* strains are carried through IncF-type plasmids or class 1 integrons, including *qnr* (fluoroquinolones), *dfrA1* (diaminopyrimidines), *bla_{SHV}*, *bla_{TEM-1}*, *bla_{CTX-M}* (β -lactams), *tetA*, *tetB* (tetracyclines), *aac* (6)-*Ib*, *aadA1* (aminoglycosides), *sul* (sulfonamides), and *cat-1*, *cmlA* (phenicols), being the most commonly found in diarrheagenic strains, food, water, and some livestock animals [279–282]. Similar findings have been documented elsewhere, e.g., China, Italy, Korea, or Nigeria in foods (including rabbits, poultry, pigs, dairy/beef cattle) [280].

Pathogenic *E. coli* utilizes a combination of specific virulence factors (molecules that enable it to cause disease) and antimicrobial resistance mechanisms to evade host immune responses and survive antibiotic treatments. These traits allow the bacteria to colonize tissues and cause severe intestinal or extraintestinal infections [281]. In this regard, at least 11 distinct pathotypes have been described in regional research [279].

For example, a recent study found that *E. coli* was present in 100% of food samples, with higher contamination rates in vegetables and fruits. Enterotoxigenic (ETEC: *elt*) and Uropathogenic *E. coli* (UPEC: *vat*, *cnf1*, *hylA*) genes were also identified in all samples; meanwhile, Diffusely Adhering *E. coli* (DAEC: *afa*) was found in 27% of the assay objects. The most effective antibiotics against the recovered strains were TET, STX, polymyxins, and quinolones [282]. Other virulence factors of pathogenic *E. coli*, tested elsewhere, include Enteropathogenic (EPEC: *eaeA*, *bfpA*), Enterohemorrhagic (EHEC: *hylA*), and Enteroinvasive *E. coli* (EIEC: *ial*) [283].

Li et al. [284] found a detection rate of *E. coli* isolated from foods as follows: pork (59.6%), mutton (52.6%), retail fresh milk (52.4%), duck (36.4%), beef (35.3%), chicken (33.3%), RTE food (12.9%), and fish and vegetables were <11%. The detection rate of resistance genes was as follows: *tetA* (38%), *tetB* (27%), *bla_{OXA}* (40%), *bla_{TEM}* (20%), *floR* (20%), *sul1* (16%), *sul2* (27%), *aad_{Ala}* (19%), *aadB* (11%), *strA* (28%), and *strB* (24%); *tetC* and *bla_{PSE}* were not detected. Virulence genes *fimC*, *agg*, *stx2*, *fimA*, *fyuA*, *papA*, *stx1*, and *eaeA* were found. Finally, they also found an interesting association between *fimC* and resistance toward CIP, GEN, AMK, LFX, and STR.

Likewise, Ronald et al. [285] isolated 30 *E. coli* with multiple antibiotic resistance (MAR) index of greater than 0.2 from meat samples; 88.9% of the samples tested positive. Isolates showed that the *tetA* (for TET resistance) and *sul1* (for STX resistance) genes were present in 100.0% of isolates, while the *bla_{TEM}* (for AMP resistance) and *strA* (for STR) genes were present in 2 isolates. O157 lipopolysaccharide O-antigen synthesis gene *rfbE-O157* was absent in all strains. Similar findings (*i.e.*, high resistance to TET and STX and high homology in genes *bla_{TEM}* and *gryA*) have been further corroborated in food/clinical *E. coli* isolates from China [286].

A recent review of 61 studies from 18 countries showed that most food samples (*i.e.*, chicken meat and eggs) had prevalence rates of 8.9% and 80.5% for Shiga toxin-producing *E. coli* (STEC) and serotype O157, respectively. Additionally, the isolates were resistant to the following antibiotics: AMC, CHL, TET, CIP, GEN, AMP, NEO, and AMO [287]. An independent study found similar trends in eggs and poultry sausage, with prevalence of 59.3% and 14.4% for *E. coli* and 2.2% and 5.6% for *Salmonella*, respectively [288].

Finally, some reports demonstrate a close link between agriculture [289] and aquaculture [290] based human activities and the dissemination of bacteria with antimicrobial resistance traits. For example, van Hamelsveld et al. [291] showed that polluted water (with normal contamination levels of *E. coli*) in continuous contact with shellfish magnifies their microbial contamination levels, rendering the food above the threshold apt for human consumption. The authors also demonstrated that the bacteria found within the shellfish show a regular and increased rate of plasmid horizontal transfer. Osman et al. [292] demonstrated that β -lactam resistance genes (*bla*_{OXA}, *bla*_{MOX}, *bla*_{CIT-like}, *bla*_{SHV}, *bla*_{FOX}) in *E. coli* were abundant both in poultry hatchlings and in the hatchery environment. Isolates were also able to form biofilms, thereby increasing their tolerance to antibiotics [293].

Furthermore, at least eight studies, published from 1940–2016, representing 18% of all revised literature, could directly link the transmission of antimicrobial resistance in *E. coli* from food animals to humans, 25 studies (56%) suggested transmission between animals and humans with no direction specified, and 12 studies (26%) did not support transmission [294].

Table 7. Summary of *Escherichia coli* bacterial isolates from foodstuffs and other food-related matrices.

Country	Isolates	Source	Antibiotics Tested	Resistance to ¹	Testing Method	References
China	<i>Escherichia coli</i>	RTE food samples (supermarkets & street markets, <i>n</i> = 150, from each)	AMP, AMC, CFX, CPZ, CTX, CHL, CIP, GTX, GEN, KAN, NAL, STR, AMK, TET, STX	AMC (80.9%), STX (79.0%), NAL (77.8%), AMP (75.6%), TET (73.1)	Agar diffusion [295]	
USA	<i>Escherichia coli</i>	all available brands of retail chicken (<i>n</i> = 1367, 88% prevalence) & turkey (<i>n</i> = 546, 91% prevalence)	AMK, AMP, SAM, CEZ, CEX, CTX, CIP, GEN, IMP, NAL, TET, STX	AMP (62%), SAM (51%), CEZ (52%), TET (76%). Prevalence is highest in isolates from turkey meat	Kirby-Bauer disk-diffusion [296]	
Qatar	<i>ESBL-Escherichia coli</i>	<i>n</i> = 456 fecal samples from food handlers	AMP, AMC, GEN, CIP, CHL, TET, TMP, SMZ, CFX	SMZ (33%), AMP (32%), TMP (31%), TET (26%)	E-test & Double Disk Sinergy (for ESBL)	[297]
Egypt	<i>Escherichia coli</i> & hemolytic <i>E. coli</i> Serotypes: O1, O2, O8, O78, O119, & O126	<i>n</i> = 450 hatchlings & <i>n</i> = 24 environmental samples per hatchery. Water, broken eggshells, infertile eggs, swabs from the workers' hands, air tunnels, floors, incubators, hatchery machines & egg refrigerators	COL, FOS, GEN, NEO, STR, CHL, ENR, CIP, NOR, FMQ, CDA, AMO, AMP, STX, ERY, SPC, RIF, OXY, DOX	CIP (62.9% environment & 30% hatchlings)	Kirby-Bauer disk-diffusion [294]	
China	<i>Escherichia coli</i>	<i>n</i> = 431 retail foods. Pork, mutton, fresh milk, duck, beef, chicken, RTE foods, fish & vegetables	AMP, CFX, CZA, GEN, IMP, CIP, LFX, TET, CHL, AMK, PIP, STX, ERY, STR, NAL, PMB	TET (52%), AMP (42%), STX (37%), AMO (33%), NAL (32%)	Kirby-Bauer disk-diffusion [284]	

Iran	<i>Escherichia coli</i>	<i>n</i> = 350 Pastry cream samples	GEN, STR, NAL, CFM, CHL, CIP, TET, OXA, VAN, PEN	TET, VAN, OXA (100%)	Kirby-Bauer disk-diffusion [298]
Guadeloupe	ESBL- <i>Escherichia coli</i>	<i>n</i> = 216 fecal samples from food-producing animals. <i>n</i> = 124 pigs, <i>n</i> = 75 beef cattle, & <i>n</i> = 17 poultry litter.	AMP, AMC, TMC, CFX, CZA, CEX, ERT, GEN, AMK, STR, ENR, NAL, CIP, TET, STX, FOS	Usage in farm: Tetracyclines 100–33%, β -lactams $\leq$ 54.5, aminoglycosides: $\leq$ 45.5%. Resistance: TET (50–82%), AMP (61–100%), STR (46–73%)	Kirby-Bauer disk-diffusion [289]
Poland	<i>Escherichia coli</i> & <i>Klebsiella</i> spp.	<i>n</i> = 4496 <i>E. coli</i> isolates from liver, spleen, heart, and lungs of chickens from commercial farms	AMO, AMC, COL, DOX, ENR, FLO, NEO, NOR, SPC, STX	AMC (73%), DOX (52%), ENR (49%), NOR (45%), FLO (38%)	Broth microdilution [299]
Emirate of Abu Dhabi	<i>Escherichia coli</i>	Isolated from different animal species (camel <i>n</i> = 42, sheep <i>n</i> = 42, goat <i>n</i> = 54, & poultry <i>n</i> = 27)	TET, OXY, PEN, AMP, AMO, OXA, CEX, GEN, STX, ENR, CFT	AMP, TET, STX, GEN, ENR (95–83%)	Kirby-Bauer disk-diffusion [300]
Ethiopia	<i>Escherichia coli</i> (54.7% prevalence) & <i>Escherichia coli</i> O157:H7 (6.5% prevalence)	<i>n</i> = 384 different samples of foods of bovine origin. Carcass swabs (<i>n</i> = 162), udder milk (<i>n</i> = 146) & milk tank (<i>n</i> = 6), yoghurt (<i>n</i> = 36) & cottage cheese (<i>n</i> = 9), & beef swabs (<i>n</i> = 25) from abattoirs, butcher shops & restaurants	AMO, AMP, PEN, OXA, CIP, NOR, ERY, AMK, GEN, KAN, NAL, TET, DOX, STX	PEN, OXA, & VAN (100%), ERY (92%), AMO (40%), TET & AMK (4%)	Kirby-Bauer disk-diffusion [301]
Sierra Leone	<i>E. coli</i> & <i>Salmonella</i> spp.	<i>n</i> = 100 Samples Fresh Poultry Excreta Used for Vegetable Farming	AMP, CHL, ERY, CEX, PEN, STR, INN, TET	TET, ERY, CEX (100%), AMP (12%)	Kirby-Bauer disk-diffusion [302]
Bangladesh	Pathogenic <i>Escherichia coli</i>	Potato smash & chicken curry, 25 samples each	AMP, ATM, CIP, COL, CHL, GEN, LFX, TET	AMP (91%), COL (52%)	Kirby-Bauer disk-diffusion [283]
Kenya	<i>Escherichia coli</i>	<i>n</i> = 105 RTE meat products	AMK, AMP, TET, NFT, NAL, STX, CIP	STX (80%), TET (73%), STR & AMP (67%)	Kirby-Bauer disk-diffusion [285]
New Zealand	<i>Escherichia coli</i>	Aquatic wild foods/“mahinga kai”: water, watercress, <i>Austrovenus stutchburyi</i> , <i>Nasturtium officinale</i> & <i>Perna canaliculus</i>	CHL, TET, CIP, TMP, GEN, KAN, AMP, CFX, CZA	AMP, CIP, CHL	Combination disk method [291]
Bangladesh	ESBL-producing <i>E. coli</i>	<i>n</i> = 400 meat samples (beef & sheep, 200 each)	STX, CIP, COL, ERY, STR, GEN	STR, ERY, AMP (100%) ESBL (21%) resistant to CFX & CTX	Kirby-Bauer disk-diffusion Double Disk Sinergy (for ESBL) [303]
Burundi	<i>Escherichia coli</i>	<i>n</i> = 30 heat-treated fresh milk samples	TET, CIP, AMP, CHL, STX, GEN	Amp (100%), TET (75%), STX (33%)	Kirby-Bauer disk-diffusion [304]

Mexico	ETEC/UEPC/DAEC	<i>n</i> = 189 street foods	TET, CPX, STX, AMP, AMK, PMB, LFX, AMC, GEN	AMC (100%), CPX (45%)	Kirby-Bauer disk-diffusion [282]
Benin	<i>E. coli</i> & <i>Salmonella</i> spp.	<i>n</i> = 135 table egg pools, <i>n</i> = 90 poultry sausages <i>n</i> = 81 clinical isolates	CTX, CXM, GEN, STR, TET, NAL, CIP, AMC, CHL, NFT	TET (88%), STR (65%), AMC (55%)	Kirby-Bauer disk-diffusion [288]
Ethiopia	<i>E. coli</i> O157:H7	Fish (<i>n</i> = 120) and fish handling equipment	KAN, TET, SMZ, AMP, ERY, STR	AMP, PEN	Kirby-Bauer disk-diffusion [305]

¹ Numbers in round brackets next to the first antibiotic represent the percentage of resistance reported for the antibiotic for which the strains analyzed showed the most tolerance. If no number is indicated, please assume 100% resistance.

7. *Cronobacter* spp.

The genus *Cronobacter* comprises seven species that are opportunistic pathogenic, Gram-negative, facultative anaerobic, non-spore-forming, bacillus-shaped bacteria. They can cause serious and even fatal infections, affecting infants, immunocompromised patients, and the elderly. In infants, the typical infection can cause meningitis, necrotizing enterocolitis, and bacteremia, and is usually foodborne, whereas in adults, bacteremia and urosepsis are typical and are normally nosocomial [306].

Foodwise *C. sakazakii* has been isolated from various plant- and animal-derived matrices, including confectionery, dried herbs and spices, teas, nuts, pasta, rice, oats, linseed, and cereal products, fresh/frozen produce [306,307], fruits, and powdered infant, follow-up formulas and milk, cheese, and meats (usually powdered foods with low moisture and water activity values). Table 8 provides a detailed account of the prevalence of the bacteria. The work of Sani and Odeyemi, and, more recently, that of Kumar et al., is recommended [308,309].

A pivotal study on *C. sakazakii* in households was conducted in 2012 and demonstrated that the bacterium is not only a relevant foodborne pathogen but is also present in food-related domestic environments. From this study, 76.1% of isolates were resistant to penicillin. Most of the research regarding *C. sakazakii* in foods and its antibiotic resistance profile has been focused on infant formulae. Overall, strains isolated from foods have shown special resistance to β -lactams. For example, Fei et al. [310] found resistance to AMC/AMP with MIC $\leq 8 \mu\text{g}\cdot\text{mL}^{-1}$. Similar values were found for isolates from a milk processing facility in Serbia [311]. The EUCAST threshold for assessing AMC is $8 \mu\text{g}\cdot\text{mL}^{-1}$. They also found that some strains were tolerant to dryness and osmotic pressure, which has a profound impact on food processing and storage [312,313]. More worryingly, at least one study has positively identified *C. sakazakii* as an ESBL-producer (e.g., class C β -lactamases) [312,314]. On the other hand, a Brazilian study demonstrated that 94.4% of isolates from infant foods were tested resistant to cefazolin. Additionally, six antibiotic-resistance genes, particularly those encoding efflux pumps, were identified. Pakbin et al. [315] found that isolates from powdered infant formula milk were highly resistant to AMC, AMP, CEX, CFP, ERY, CTX, CIP, and CHL and susceptible to GEN, TET, NOR, and ATM antibiotics. Meanwhile, the *bla*_{CTX-M-1} gene was detected in 96% of the isolates. Previously, the same research group [316] also found a higher prevalence of *C. sakazakii* in bulky imported domestically packaged powdered milk infant formula samples.

Although isolated strains do seem to possess relevant resistant phenotypes, the overall prevalence is relatively low. For example, in an Iranian study of 200 samples (100 infant formula and 100 baby foods), only eight (4%) samples tested positive [317]. However, the sole presence of a few bacteria and their multidrug resistance profile suggests that this pathogen can disseminate resistance genes with relative ease within these food samples [318]. Worryingly, most reports of *C. sakazakii* have demonstrated multiple drug resistance profiles and biofilm formation in isolates from infant contaminated foods; both bacterial mechanisms result in therapy failure [319]. Biofilms and drug resistance profiles are the primary drivers of chronic, recurrent, and untreatable infections. Together, they cause therapy failure by creating physical, chemical, and genetic barriers that prevent antibiotics from either reaching or eliminating target pathogens [320].

Holý et al. [321] found in their *C. sakazakii* isolates a type IV secretion system-related gene (found on hybrid or cointegrate plasmids, incFIB/incH1 are central components of bacterial conjugation machineries), which plays a pivotal role in bacterial evolution by facilitating the horizontal transfer of genetic material—such as antimicrobial resistance and virulence factors—between bacteria [307,318]. These genes have also been described in *C. malonaticus*, *C. turicensis*, *C. muytjensii*, and *C. universalis* [322]. Several antibiotic target site modification and efflux-related genes have also been identified in *C. sakazakii* isolates (e.g., *mdf(A)*, which encodes a multidrug efflux transporter, *emrR*, *emrB*, *rsmA*, *cRP*, *adeF*, and *marR* for fluoroquinolones or *fos* genes encoding resistance for fosfomycin) [323–325]. Recently, in a large Chinese study performed by Gan et al. [318], Strains isolated from powdered infant formula and supplementary food collected from 29 provinces exhibited at least 7 antimicrobial resistance genes, which corroborates other findings supporting multidrug resistance. Similarly, Fei et al. [325] also isolated *C. sakazakii* from goat milk-based infant formula from Shaanxi Province, China. From the serotypes found, O2 was the most abundant ($n = 15/32$, 46.88%) and showed, as well, mostly resistance to antimicrobials such as CEP (87.5%) and AMO (25%). Resistance phenotyping for other *Enterobacteriaceae* can be found in the work by Mardaneh et al. [326]. Recently, Song et al. [327] found, using a proteomics approach, that there is a certain relation between antibiotic tolerance and the production of enterotoxin and hemolysin.

Antibiotic resistance and the production of enterotoxins and hemolysins are critical virulence characteristics in pathogenic bacteria like *S. aureus*, *E. coli*, and *B. cereus*. Bacteria often acquire genes for drug resistance and virulence simultaneously and therefore are often located together on the same genetic elements (e.g., plasmids, transposons, or pathogenicity islands). When bacteria share genetic material, they can spread both traits to neighboring cells [328].

When antibiotics are used, bacteria with resistance genes survive. Because these genes are linked to toxin-producing genes, antibiotic use promotes the survival and proliferation of the most virulent strains. Hence, it appears to be a correlation between multidrug resistance and an increased presence of hemolysin and enterotoxin genes. Strains exhibiting this combination are associated with severe complications and a higher mortality rate in clinical settings [328].

On the other hand, Oloninefa et al. [329] demonstrated that strains with antibiotic sensitivity (to CIP and GEN) were eliminated in culture conditions after 240 min of exposure to the antimicrobials. Hence, those sensitive strains still seem to respond quickly to antibiosis. Interestingly, Capita et al. [330] described that *C. sakazakii* strains increased their resistance profile after treatment with several biocides (e.g., sodium hypochlorite, peracetic acid, benzalkonium chloride) commonly used in food processing equipment cleaning procedures [331,332]. There is also compelling evidence that suggests that antibiotic resistance increases virulence in *Cronobacter* spp. isolates from food [321,333].

Finally, resistance in isolates from clinical cases of neonatal hemorrhagic diarrhea and meningitis has been reported in countries like Mexico and China [322,324,334–336].

Table 8. Summary of *Cronobacter* bacterial isolates from foodstuffs and other food-related matrices.

Country	Isolates	Source	Antibiotics Tested	Resistance to ¹	Testing Method	References
China	<i>C. sakazakii</i> ($n = 9$)	Infant Formula Milk	AMP, TET, VAN, CHL, NOR, CEP, GEN, ERY	VAN (100%), CEP	Kirby-Bauer disk-diffusion	[337]
USA	<i>C. sakazakii</i>	Kitchens Surfaces ($n = 234/78$)	TET, GEN, STR, AMP, KAN, PEN, CHL, CIP, CEF, NAL	PEN (76.1%), TET, CIP, NAL	Kirby-Bauer disk-diffusion	[338]

Egypt	<i>C. sakazakii</i> (n = 11)	Powdered Infant Formula Milk (n = 173), Powdered Infant Foods (n = 61), Blood (n = 7) & Environment (n = 3)	LFX, NOR, OFX, CIP, NAL, GEN, IMP, CAZ, SXT, ATM, STR, CTX, AMC, CPX, RIF, AMP	RIF (100%), AMP, CPX	Kirby-Bauer disk-diffusion	[339]
Bangladesh	<i>C. sakazakii</i> (n = 6)	Assorted foods (n = 54)	VAN, CIP, NFT, CHL, PEN, TET, AMP, IMP, DOX, NEO, AMK, GEN	PEN (100%), AMP, IMP, NEO	Kirby-Bauer disk-diffusion	[312]
Iran	<i>C. sakazakii</i>	Infant Formula & Baby Foods (n = 200)	TIG, TET, TIC, CHL, CZA, CIP, CFP, IMP, LFX, MIN, PIP, TZP, CAB, TMP, STX, MFX, GEN, COL, AMP, AMO, ATM, CFX, AMK, STR, MEM, MEZ, NAL	AMO, TIG, TIC, ATM, CZA	Kirby-Bauer disk-diffusion	[317]
China	<i>C. sakazakii</i> (n = 66) & <i>C. malonaticus</i> (n = 4)	Powered Infant Formula & Processing Environment	AMK, AMC, AMP, ATM, CFX, CZA, CFP, CHL, CIP, COL, GEN, IMP, LFX, MEM, MFX, PIP, TZP, TET, STX	AMC (100%), AMP, CEZ	BD Phoenix™ 100 Automated Microbiology System	[310]
Iran	<i>C. sakazakii</i> (n = 9)	Powdered Infant Formula Milk (n = 125)	AMO, AMP, ATM, CFX, CZA, IMP, MEM, MEZ, TIC, MFX, NAL, STR, GEN, AMK, TET, CHL, LFX, CIP, CFP, MIN, PIP, TZP, CAB, TMP, STX, TIG, COL	CFX (77.8%), MEZ, STR, MIN, TMP, TIG	Kirby-Bauer disk-diffusion	[326]
Brazil	<i>C. sakazakii</i> (n = 44)	Powdered Infant Cereals (n = 11)	AMO, COL, CIP, CHL, ENR, GEN, STR, MEM, STX, TET, CEZ, CPD	CEZ (6.81%), AMO, STX, CFP	Kirby-Bauer disk-diffusion	[340]
Brazil	<i>Cronobacter</i> spp. (n = 72)	Infant Formulas (n = 77) & Cereals (n = 75)	AMO, ENR, CIP, CHL, GEN, STR, MEM, STX, TET, CEZ, CPD	CEZ (94.4%), AMO, CED, STR, STX	Kirby-Bauer disk-diffusion	[323]
Nigeria	<i>C. sakazakii</i>	Powdered Infant Formula	AMO, AMC, ERY, TET, CHL, AMO, AMC, GEN, CLX, STX, CHL, CIP, STR	ERY, TET, CLX, STX	Kirby-Bauer disk-diffusion	[329]
Iran	<i>C. sakazakii</i> (n = 6)	Imported and Domestic Powdered Milk Infant Formula (n = 118)	TET, LFX, AMC, AMO, CHL, IMP, CIP, CFP	AMP & AMO (100%), CIP & TET (intermediate)	Kirby-Bauer disk-diffusion	[316]
Iraq	<i>C. sakazakii</i> , <i>C. malonaticus</i> , <i>C. Dublinensis</i>	Powdered Infant Formula (n = 130), Spices & Herbs (n = 55) & Bouillon Flavoured Powder (n = 50)	AMP, CHL, CEP, ERY, GEN, KAN, NAL, STR, TET, NOR	CEP (66%), KAN, STR, ERY, NOR	Kirby-Bauer disk-diffusion	[341]
Serbia	<i>C. sakazakii</i> (n = 15)	Milk Powder Processing Facility (n = 100)	TZP, AMC, SAM, MEM, CFX	AMC (7%)	Epsilonometer test (Etest)	[311]
China	<i>C. sakazakii</i> (n = 1048 infant supplementary food & n = 7 powdered infant formula)	Powdered infant formula and supplementary food (n = 12,105)	PEN, AMP, SAM, CTX, CAZ, IMP, GEN, KAN, TET, CIP, NAL, SXT, CHL		Biofosun Gram-negative panels by the broth dilution method	[318]

Iran	<i>C. sakazakii</i> (n = 25)	Powdered Infant Formula Milk (n = 364)	NAL, CEX, AZM, STR, AMC, IMP, KAN, AMO, AMC, AMO, AMP, NOR, AMP, TET, CFP, CEX, CFP, ERY, GEN, COL, CHL, ERY, CTX, CIP, CHL CTX, CIP	Kirby–Bauer disk diffusion [315]
Mexico	<i>C. sakazakii</i> (n = 7), <i>F. helveticus</i>	Powdered and dairy formulas	CAZ, CTX, AMO, AMC, CIP, CEP, NAL, GEN, TET, CHL, AMP	CEP (71.4%), NAL, AMP Kirby–Bauer disk diffusion [324]
Czech Republic	<i>C. sakazakii</i> (n = 15)	Food (n = 997), milk-producing environments (n = 855), & feces from live poultry (n = 148)	AMP, AMC, CZA, CIP, CHL, CFP, GEN, CEP, TET, NAL	CEP (93%), AMP, TET, CFP, AMC Kirby–Bauer disk diffusion [321]
China	<i>C. sakazakii</i> (n = 35)	Powdered Infant Formula and Processing Environment	AMP, PIP, CEZ, CMX, CZA, CTX, CPZ, AMK, GEN, KAN, NEO, TET, DOX, ERY, NOR, OFX, CIP, PMB, STX, CHL	ERY (100%), STX, NEO, CEZ, KAN, CTX, AMK, AMP, CZA, CXM, PMB, GEN, NOR, PIP Kirby–Bauer disk diffusion [342]
China	<i>C. sakazakii</i> (n = 96)	Infant Formula Powder & Processing Environment	AMP, PIP, CAB, CEZ, CXM, CZA, CTX, CPZ, AMK, GEN, KAN, NEO, TET, DOX, ERY, MDY, VAN, LIN, NOR, OFX, CIP, CHL, PMB, VAN, STX, LIN	ERY (100%), MDY, Kirby–Bauer disk diffusion [327]
Iran	<i>C. sakazakii</i> (n = 2/14 suspected)	Raw milk and Instant Formula (n = 120 raw, 60 powdered milk samples)	ERY, GEN, CHL, AMP, NEO, AMO, TET, CIP, SMZ	ERY (100%), AMP, AMO, TET Kirby–Bauer disk diffusion [343]

¹ Numbers in round brackets next to the first antibiotic represent the percentage of resistance reported for the antibiotic for which the strains analyzed showed the most tolerance.

8. Conclusions

Continuous use of an antibiotic that has reached the end of therapeutic usefulness, due to propagated or widespread resistance among bacteria, will only conserve the selective pressure and favor further phenotypic resistance. Consequences of the overuse of antibiotics, even after years of regulatory restrictions and efforts to reduce their application, are still visible today (e.g., overall resistance toward tetracyclines in *Campylobacter*). Recent cumulative findings reinforce that, despite some efforts [344], antibiotic resistance within pathogen species is increasing. The data herein are in line with current predictions, which set *E. coli* and nontyphoidal *Salmonella* resistance for β -lactams (often producing Extended-Spectrum Beta-Lactamases, which are resistant to critical, last-resort antibiotics), sulfonamides, and tetracyclines above 50% in most African, Asian (e.g., China), and American (e.g., Brazil, Chile) territories [345]. Foodborne pathogen contamination *per se* is a worrisome health issue that requires constant surveillance programs and allocation of resources, but disseminated antibiotic resistance among these bacteria exacerbates the problem as therapeutic options to fight infections caused by these microorganisms become more limited. This is especially true since a significant percentage of isolates are resistant to three or more antibiotic classes. Some crucial food industries deserve a closer look in terms of pathogen monitoring and inspection (e.g., Infant powdered milk products for *Cronobacter*, minced or ground bovine beef for *Staphylococcus*, raw milk/dairy for *Listeria*, or the poultry industry for *Campylobacter* and *Salmonella*) as they seem to be hotspots for multidrug-resistant bacterial proliferation and transmission. Resistant strains often originate in animal gut microbiota (often due to farm antibiotic usage) and transfer to humans through contamination

of meat, vegetables, or water. These foodborne pathogens can cause severe, antibiotic-resistant infections, limiting therapeutic options in clinical settings. Nevertheless, despite its relevance, we see few surveillance efforts regarding bacterial contamination and drug resistance within animal feed, an issue that should be addressed, especially in pet foods, considering the proximity of pets to their owners. In this respect, the One Health approach should be the go-to tool for preventing bacterial food contamination throughout the food chain. Food safety-wise, transmission pathways of antimicrobial-resistant bacteria can be diminished by proper handling and cooking of raw meat (especially poultry) to prevent cross-contamination. Also, for some particular foods, ensuring a continuous cold chain and proper refrigeration is a must.

Ethics Statement

Not applicable.

Informed Consent Statement

Not applicable.

Data Availability Statement

Not applicable.

Funding

This research received no external funding.

Declaration of Competing Interest

The author declares that he has no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

1. Lisboa HM, Pasquali MB, dos Anjos AI, Sarinho AM, de Melo ED, Andrade R, et al. Innovative and Sustainable Food Preservation Techniques: Enhancing Food Quality, Safety, and Environmental Sustainability. *Sustainability* **2024**, *16*, 8223. DOI:10.3390/su16188223
2. Jordan K, Dalmasso M, Zentek J, Mader A, Bruggeman G, Wallace J, et al. Microbes versus microbes: Control of pathogens in the food chain. *J. Sci. Food Agric.* **2014**, *94*, 3079–3089. DOI:10.1002/jsfa.6735
3. Parra-Flores J, Holý O, Riffo F, Lepuschitz S, Maury-Sintjago E, Rodríguez-Fernández A, et al. Profiling the Virulence and Antibiotic Resistance Genes of *Cronobacter sakazakii* Strains Isolated From Powdered and Dairy Formulas by Whole-Genome Sequencing. *Front. Microbiol.* **2021**, *12*, 694922. DOI:10.3389/fmicb.2021.694922
4. FSN Food Safety News. Ice Cream Bars Recalled After Testing Finds *Salmonella*. Available online: <https://www.foodsafetynews.com/2024/04/ice-cream-bars-recalled-after-testing-finds-salmonella/> (accessed on 18 June 2026).
5. FSN Food Safety News. Officials Confirm Planned Closure of Quaker Oats Plant Behind Recall of Dozens of Products. Available online: <https://www.foodsafetynews.com/2024/04/officials-confirm-planned-closure-of-quaker-oats-plant-behind-recall-of-dozens-of-products/> (accessed on 19 June 2026).
6. U.S. Food and Drug Administration [US FDA]. Hormel Foods Sales, LLC Recalls a Limited Number of Planters® Honey Roasted Peanuts 4 Oz. and Planters® Deluxe Lightly Salted Mixed Nuts 8.75 Oz. Because of Possible Health Risk. Available online: <https://www.fda.gov/safety/recalls-market-withdrawals-safety-alerts/hormel-foods-sales-llc-recalls-limited-number-plantersr-honey-roasted-peanuts-4-oz-and-plantersr> (accessed on 19 June 2026).
7. U.S. Food and Drug Administration [US FDA]. Outbreak Investigation of *Listeria monocytogenes*: Queso Fresco and Cotija Cheese (February 2024). Available online: <https://www.fda.gov/food/outbreaks-foodborne-illness/outbreak-investigation-listeria-monocytogenes-queso-fresco-and-cotija-cheese-february-2024> (accessed on 19 June 2026).
8. FSN Food Safety News. California Company Recalls Shelled Walnuts Linked to Outbreak of *E. coli* Infections. Available online: <https://www.foodsafetynews.com/2024/04/california-company-recalls-shelled-walnuts-linked-to-outbreak-of-e>

- coli-infections/ (accessed on 19 June 2026).
9. FSN Food Safety News. French's Original Crispy Fried Onions Recalled Over *Staphylococcus aureus* Concerns. Available online: https://www.foodsafetynews.com/2024/05/frenchs-original-crispy-fried-onions-recalled-over-staphylococcus-aureus-concerns/#google_vignette (accessed on 19 June 2026).
 10. FSN Food Safety News. Hummus Recalled in Canada Over Yeast Contamination. Available online: <https://www.foodsafetynews.com/2024/04/hummus-recalled-in-canada-over-yeast-contamination/> (accessed on 19 June 2026).
 11. FSN Food Safety News. Recalls Down But Still More than 2000 in France for 2023. Available online: <https://www.foodsafetynews.com/2024/04/recalls-down-but-still-more-than-2000-in-france-for-2023/> (accessed on 19 June 2026).
 12. U.S. Food and Drug Administration [US FDA]. Recalls, Market Withdrawals, & Safety Alerts. Available online: <https://www.fda.gov/safety/recalls-market-withdrawals-safety-alerts> (accessed on 19 June 2026).
 13. Tropea A. Microbial Contamination and Public Health: An Overview. *Int. J. Environ. Res. Public Health* **2022**, *19*, 7441. DOI:10.3390/ijerph19127441
 14. Abalkhail A. Frequency and Antimicrobial Resistance Patterns of Foodborne Pathogens in Ready-to-Eat Foods: An Evolving Public Health Challenge. *Appl. Sci.* **2023**, *13*, 12846. DOI:10.3390/app132312846
 15. Aladhadh M. A Review of Modern Methods for the Detection of Foodborne Pathogens. *Microorganisms* **2023**, *11*, 1111. DOI:10.3390/microorganisms11051111
 16. Paramithiotis S. Molecular Targets for Foodborne Pathogenic Bacteria Detection. *Pathogens* **2023**, *12*, 104. DOI:10.3390/pathogens12010104
 17. de Benito A, Besse NG, Desforges I, Gerten B, Ruiz B, Tomás D. Validation of standard method EN ISO 22964:2017—Microbiology of the food chain—Horizontal method for the detection of *Cronobacter* spp. *Int. J. Food Microbiol.* **2019**, *288*, 47–52. DOI:10.1016/j.ijfoodmicro.2018.03.025
 18. Bramwell P, Clarke D, de Silva T. The food microbiological analyst: Pairing tradition with the future. *Microbiol. Australia* **2022**, *43*, 57–60. DOI:10.1071/MA22024
 19. Hadi J, Rapp D, Dhawan S, Gupta SK, Gupta TB, Brightwell G. Molecular detection and characterization of foodborne bacteria: Recent progresses and remaining challenges. *Compr. Rev. Food Sci. Food Saf.* **2023**, *22*, 2433–2464. DOI:10.1111/1541-4337.13153
 20. Liu S, Zhao K, Huang M, Zeng M, Deng Y, Li S, et al. Research progress on detection techniques for point-of-care testing of foodborne pathogens. *Front. Bioeng. Biotechnol.* **2022**, *10*, 958134. DOI:10.3389/fbioe.2022.958134
 21. Truchado P, Randazzo W. New Challenges for Detection and Control of Foodborne Pathogens: From Tools to People. *Foods* **2022**, *11*, 1788. DOI:10.3390/foods11121788
 22. Quintela IA, Vasse T, Lin C-S, Wu VCH. Advances, applications, and limitations of portable and rapid detection technologies for routinely encountered foodborne pathogens. *Front. Microbiol.* **2022**, *13*, 1054782. DOI:10.3389/fmicb.2022.1054782
 23. Paparella A, Maggio F. Detection and Control of Foodborne Pathogens. *Foods* **2023**, *12*, 3521. DOI:10.3390/foods12193521
 24. Wang B, Wang H, Lu X, Zheng X, Yang Z. Recent Advances in Electrochemical Biosensors for the Detection of Foodborne Pathogens: Current Perspective and Challenges. *Foods* **2023**, *12*, 2795. DOI:10.3390/foods12142795
 25. Zhao X, Bhat A, O'Connor C, Curtin J, Singh B, Tian F. Limits of Detection Analysis of Advanced Technologies for Bacterial Detection in Food Samples: Review & Future Perspective. **2024**, *preprints*. DOI:10.20944/preprints202401.1268.v1
 26. CLSI VET01S Ed7. Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated From Animals. Available online: <https://clsi.org/shop/standards/vet01s/> (accessed on 19 June 2026).
 27. European Food Safety Authority (EFSA), Amore G, Beloeil P-A, Ferrer-Bustins N, Garcia Fierro R, Rizzi V, et al. Manual for reporting 2025 antimicrobial resistance data under Directive 2003/99/EC and Commission Implementing Decision (EU) 2020/172. EFSA supporting publication 2026:EN-9909. *EFSA* **2026**, *23*, 9909E. DOI:10.2903/sp.efsa.2026.EN-9909
 28. Gajic I, Kabic J, Kekic D, Jovicevic M, Milenkovic M, Mitic Culafic D, et al. Antimicrobial Susceptibility Testing: A Comprehensive Review of Currently Used Methods. *Antibiotics* **2022**, *11*, 427. DOI:10.3390/antibiotics11040427
 29. Sweeney MT, Lubbers BV, Schwarz S, Watts JL. Applying definitions for multidrug resistance, extensive drug resistance and pan drug resistance to clinically significant livestock and companion animal bacterial pathogens. *J. Antimicrob. Chemother.* **2018**, *73*, 1460–1463. DOI:10.1093/jac/dky043
 30. Ma F, Xu S, Tang Z, Li Z, Zhang L. Use of antimicrobials in food animals and impact of transmission of antimicrobial resistance on humans. *Biosaf. Health* **2021**, *3*, 32–38. DOI:10.1016/j.bshealth.2020.09.004

31. Sati H, Carrara E, Savoldi A, Hansen P, Garlasco J, Campagnaro E, et al. The WHO Bacterial Priority Pathogens List 2024: A prioritisation study to guide research, development, and public health strategies against antimicrobial resistance. *Lancet Infect. Dis.* **2025**, *25*, 1033–1043. DOI:10.1016/S1473-3099(25)00118-5
32. Muntean M, Muntean AA, Preda M, Manolescu L, Dragomirescu C, Popa MI, et al. Phenotypic and genotypic detection methods for antimicrobial resistance in ESKAPE pathogens (Review). *Exp. Ther. Med.* **2022**, *24*, 508. DOI:10.3892/etm.2022.11435
33. Butler MS, Henderson IR, Capon RJ, Blaskovich MA. Antibiotics in the clinical pipeline as of December 2022. *J. Antibiotics* **2023**, *76*, 431–473. DOI:10.1038/s41429-023-00629-8
34. Huseby DL, Cao S, Zamaratski E, Sooriyaarachchi S, Ahmad S, Bergfors T, et al. Antibiotic class with potent *in vivo* activity targeting lipopolysaccharide synthesis in Gram-negative bacteria. *Proc. Natl. Acad. Sci. USA* **2024**, *121*, e2317274121. DOI:10.1073/pnas.2317274121
35. Ghimpeanu OM, Pogurschi EN, Popa DC, Dragomir N, Drăgoteiu T, Mihai OD, et al. Antibiotic Use in Livestock and Residues in Food—A Public Health Threat: A Review. *Foods* **2022**, *11*, 1430. DOI:10.3390/foods11101430
36. Aspenberg M, Sasane SM, Nilsson F, Brown SP, Waldetoft KW. Hygiene may attenuate selection for antibiotic resistance by changing microbial community structure. *Evol. Med. Public Health* **2023**, *11*, 1–7. DOI:10.1093/emph/eoac038
37. Kasimanickam V, Kasimanickam M, Kasimanickam R. Antibiotics Use in Food Animal Production: Escalation of Antimicrobial Resistance: Where Are We Now in Combating AMR? *Med. Sci.* **2021**, *9*, 14. DOI:10.3390/medsci9010014
38. Ye J, Chen X. Current Promising Strategies against Antibiotic-Resistant Bacterial Infections. *Antibiotics* **2023**, *12*, 67. DOI:10.3390/antibiotics12010067
39. Velazquez-Meza ME, Galarde-Lóez M, Carrillo-Quiróz B, Alpuche-Aranda CM. Antimicrobial resistance: One Health approach. *Vet. World* **2022**, *15*, 743–749. DOI:10.14202/vetworld.2022.743-749
40. Okaiyeto SA, Sutar PP, Chen C, Ni JB, Wang J, Mujumdar AS, et al. Antibiotic resistant bacteria in food systems: Current status, resistance mechanisms, and mitigation strategies. *Agric. Commun.* **2024**, *2*, 100027. DOI:10.1016/j.agrcom.2024.100027
41. Taylor P, Reeder R. Antibiotic use on crops in low and middle-income countries based on recommendations made by agricultural advisors. *CABI Agric. Biosci.* **2020**, *1*, 1. DOI:10.1186/s43170-020-00001-y
42. Miller SA, Ferreira JP, LeJeune JT. Antimicrobial Use and Resistance in Plant Agriculture: A One Health Perspective. *Agriculture* **2022**, *12*, 289. DOI:10.3390/agriculture12020289
43. Verhaegen M, Bergot T, Liebana E, Stancanelli G, Streissl F, Mingeot-Leclercq MP, et al. On the use of antibiotics to control plant pathogenic bacteria: A genetic and genomic perspective. *Front. Microbiol.* **2023**, *14*, 1221478. DOI:10.3389/fmicb.2023.1221478
44. Zhang F, Cheng W. The Mechanism of Bacterial Resistance and Potential Bacteriostatic Strategies. *Antibiotics* **2022**, *11*, 1215. DOI:10.3390/antibiotics11091215
45. Okpashi VE. Pathogenic Speciation and Phylogenicity: Mechanisms of Antimicrobial Resistance in Food Quality. **2023**, preprints. DOI:10.20944/preprints202307.0568.v1
46. van Hoeck AHAM, Mevius D, Guerra B, Mullany P, Roberts AP, Aarts HJM. Acquired antibiotic resistance genes: An overview. *Front. Microbiol.* **2011**, *2*, 203. DOI:10.3389/fmicb.2011.00203
47. Krüger GI, Pardo-Esté C, Zepeda P, Olivares-Pacheco J, Galleguillos N, Suarez M, et al. Mobile genetic elements drive the multidrug resistance and spread of *Salmonella* serotypes along a poultry meat production line. *Front. Microbiol.* **2023**, *14*, 1072793. DOI:10.3389/fmicb.2023.1072793
48. Orlek A, Anjum MF, Mather AE, Stoesser N, Walker AS. Factors associated with plasmid antibiotic resistance gene carriage revealed using large-scale multivariable analysis. *Sci. Rep.* **2023**, *13*, 2500. DOI:10.1038/s41598-023-29530-y
49. Łojewska E, Sakowicz T. An Alternative to Antibiotics: Selected Methods to Combat Zoonotic Foodborne Bacterial Infections. *Curr. Microbiol.* **2021**, *78*, 4037–4049. DOI:10.1007/s00284-021-02665-9
50. Gupta R, Sharma S. Role of alternatives to antibiotics in mitigating the antimicrobial resistance crisis. *Indian J. Med. Res.* **2022**, *156*, 464–477. DOI:10.4103/ijmr.IJMR_3514_20
51. Helmy YA, Taha-Abdelaziz K, Hawwas HAEH, Ghosh S, AlKafaas SS, Moawad MMM, et al. Antimicrobial Resistance and Recent Alternatives to Antibiotics for the Control of Bacterial Pathogens with an Emphasis on Foodborne Pathogens. *Antibiotics* **2023**, *12*, 274. DOI:10.3390/antibiotics12020274
52. Calo JR, Crandall PG, O'Bryan CA, Ricke S. Essential oils as antimicrobials in food systems—A review. *Food Control* **2015**, *54*, 111–119. DOI:10.1016/j.foodcont.2014.12.040
53. Friedman M. Antibiotic-Resistant Bacteria: Prevalence in Food and Inactivation by Food-Compatible Compounds and Plant Extracts. *J. Agric. Food Chem.* **2015**, *63*, 3805–3822. DOI:10.1021/acs.jafc.5b00778
54. Hintz T, Matthews KK, Di R. The Use of Plant Antimicrobial Compounds for Food Preservation. *Biomed. Res. Int.* **2015**,

- 2015, 246264. DOI:10.1155/2015/246264
55. Fan X, Ngo H, Wu C. *Natural and Bio-Based Antimicrobials for Food Applications*; ACS Symposium Series; American Chemical Society: Washington, DC, USA, 2018; Volume 1287. DOI:10.1021/bk-2018-1287
 56. Nair DVT, Venkitanarayanan K, Johny AK. Antibiotic-Resistant *Salmonella* in the Food Supply and the Potential Role of Antibiotic Alternatives for Control. *Foods* **2018**, *7*, 167. DOI:10.3390/foods7100167
 57. Álvarez-Martínez FJ, Barrajón-Catalán E, Herranz-López M, Micol V. Antibacterial plant compounds, extracts and essential oils: An updated review on their effects and putative mechanisms of action. *Phytomedicine* **2021**, *91*, 153626. DOI:10.1016/j.phymed.2021.153626
 58. Deng W, Quan Y, Yang S, Guo L, Zhang X, Liu S, et al. Antibiotic Resistance in *Salmonella* from Retail Foods of Animal Origin and Its Association with Disinfectant and Heavy Metal Resistance. *Microb. Drug Resist.* **2018**, *24*, 782–791. DOI:10.1089/mdr.2017.0127
 59. Giacometti F, Shirzad-Aski H, Ferrerira S. Antimicrobials and Food-Related Stresses as Selective Factors for Antibiotic Resistance along the Farm to Fork Continuum. *Antibiotics* **2021**, *10*, 671. DOI:10.3390/antibiotics10060671
 60. Mataragka A, Anthi R, Christodouli ZE, Malisova O, Andritsos ND. Presence of Major Bacterial Foodborne Pathogens in the Domestic Environment and Hygienic Status of Food Cleaning Utensils: A Narrative Review. *Hygiene* **2025**, *5*, 60. DOI:10.3390/hygiene5040060
 61. Murray LM, Hayes A, Snape J, Kasprzyk-Hordern B, Gaze WH, Murray AK. Co-selection for antibiotic resistance by environmental contaminants. *NPJ Antimicrob. Resist.* **2024**, *2*, 9. DOI:10.1038/s44259-024-00026-7
 62. Singh SP, Qureshi A, Hassan W. Mechanisms of Action by Antimicrobial Agents: A Review. *McGill J. Med.* **2021**, *19*, 4. DOI:10.26443/mjm.v19i1.217
 63. Diep B, Barretto C, Portmann AC, Fournier C, Karczmarek A, Voets G, et al. *Salmonella* Serotyping; Comparison of the Traditional Method to a Microarray-Based Method and an in silico Platform Using Whole Genome Sequencing Data. *Front. Microbiol.* **2019**, *10*, 2554. DOI:10.3389/fmicb.2019.02554
 64. Banerji S, Simon S, Tille A, Fruth A, Flieger A. Genome-based *Salmonella* serotyping as the new gold standard. *Sci. Rep.* **2020**, *10*, 4333. DOI:10.1038/s41598-020-61254-1
 65. Soliani L, Rugna G, Prosperi A, Chiapponi C, Luppi A. *Salmonella* Infection in Pigs: Disease, Prevalence, and a Link between Swine and Human Health. *Pathogens* **2023**, *12*, 1267. DOI:10.3390/pathogens12101267
 66. Jackson BR, Griffin PM, Cole D, Walsh KA, Chai SJ. Outbreak-associated *Salmonella enterica* serotypes and food Commodities, United States, 1998–2008. *Emerg. Infect. Dis.* **2013**, *19*, 1239–1244. DOI:10.3201/eid1908.121511
 67. Nadi ZR, Salehi TZ, Tamai IA, Foroushani AR, Silanpaa M, Soltan Dallal MM. Evaluation of antibiotic resistance and prevalence of common *Salmonella enterica* serovars isolated from foodborne outbreaks. *Microchem. J.* **2020**, *155*, 104660. DOI:10.1016/j.microc.2020.104660
 68. Dallal MMS, Ahmadi E, Sefidan AM, Zarebavani M, Yazdi SS, Nasser A, et al. Prevalence of Antibiotic Resistance Genes in Different *Salmonella enterica* Serovars Isolated from National Food Outbreaks. **2023**, preprint. DOI:10.21203/rs.3.rs-7052748/v1
 69. Teklemariam AD, Al-Hindi RR, Albiheyri RS, Alharbi MG, Alghamdi MA, Filimban AAR, et al. Human Salmonellosis: A Continuous Global Threat in the Farm-to-Fork Food Safety Continuum. *Foods* **2023**, *12*, 1756. DOI:10.3390/foods12091756
 70. Soubeiga AP, Kpoda DS, Compaoré MKA, Somda-Belemlougri A, Kaseko N, Rouamba SS, et al. Molecular Characterization and the Antimicrobial Resistance Profile of *Salmonella* spp. Isolated from Ready-to-Eat Foods in Ouagadougou, Burkina Faso. *Int. J. Microbiol.* **2022**, 2022, 9640828. DOI:10.1155/2022/9640828
 71. Yin X, M'ikanatha NM, Nyirabahizi E, McDermott PF, Tate H. Antimicrobial resistance in non-Typhoidal *Salmonella* from retail poultry meat by antibiotic usage-related production claims—United States, 2008–2017. *Int. J. Food Microbiol.* **2021**, *342*, 109044. DOI:10.1016/j.ijfoodmicro.2021.109044
 72. Campos-Granados CM, Gómez-Pedroso LdCS, Hernández-Pérez CF, Ballesteros-Nova NE, Rubio-Lozano MS, Sánchez-Zamorano LM, et al. Strong antibiotic resistance profiles in *Salmonella* spp. isolated from ground beef in Central Mexico. *Vet. México OA* **2023**, *10*, 23. DOI:10.22201/fmvz.24486760e.2023.1215
 73. Wójcicki M, Chmielarczyk A, Świder O, Średnicka P, Strus M, Kasperski T, et al. Bacterial Pathogens in the Food Industry: Antibiotic Resistance and Virulence Factors of *Salmonella enterica* Strains Isolated from Food Chain Links. *Pathogens* **2022**, *11*, 1323. DOI:10.3390/pathogens11111323
 74. Monte DF, Lincopan N, Fedorka-Cray PJ, Landgraf M. Current insights on high priority antibiotic-resistant *Salmonella enterica* in food and foodstuffs: A review. *Curr. Opinion Food Sci.* **2019**, *26*, 35–46. DOI:10.1016/j.cofs.2019.03.004
 75. Wang Y, Ge H, Wei X, Zhao X. Research progress on antibiotic resistance of *Salmonella*. *Food Qual. Saf.* **2022**, *6*, fyac035. DOI:10.1093/fqsafe/fyac035

76. Micha R, Khatibzadeh S, Shi P, Andrews KG, Engell RE, Mozaffarian D. Global, regional and national consumption of major food groups in 1990 and 2010: A systematic analysis including 266 country-specific nutrition surveys worldwide. *BMJ Open* **2015**, 5, e008705. DOI:10.1136/bmjopen-2015-008705
77. Kilonzo-Nthenge A, Mukuna W. *Salmonella* and Antimicrobial Resistance in Fresh Produce. In *Salmonella—A Re-Emerging Pathogen*, 1st ed.; Mascellino MT, Ed.; IntechOpen: London, UK, 2018; pp. 75–82. DOI:10.5772/intechopen.72894
78. Asfaw T, Genetu D, Shenkute D, Shenkute TT, Yitayew B. Commonly Consumed Vegetables as a Potential Source of Multidrug-Resistant and β -Lactamase-Producing Bacteria in Debre Berhan Town, Ethiopia. *Infect. Drug Resist.* **2023**, 16, 3693–3705. DOI:10.2147/IDR.S412126
79. Puig-Peña Y, Leyva-Castillo V, Tejedor-Arias R, Illnait-Zaragozi MT, Aportela-López N, Camejo-Jardines A, et al. Antimicrobial Resistance in Bacteria Isolated from Foods in Cuba. *MEDICC Rev.* **2020**, 22, 40–45. DOI:10.37757/MR2020.V22.N3.9
80. Cao T, Liu P, Li Y, Cui M, Zhang C, Wang Y, et al. Prevalence of *Salmonella* and Antimicrobial Resistance in Isolates from Food Animals—Six PLADs, China, 2019. *China CDC Wkly.* **2021**, 3, 514–517. DOI:10.46234/ccdcw2021.132
81. Ejo M, Garedew L, Alebachew Z, Worku W. Prevalence and Antimicrobial Resistance of *Salmonella* Isolated from Animal-Origin Food Items in Gondar, Ethiopia. *BioMed Res. Int.* **2016**, 2016, 4290506. DOI:10.1155/2016/4290506
82. Quesada A, Reginatto GA, Ruiz Español A, Colantonio LD, Burrone MS. Resistencia antimicrobiana de *Salmonella* spp. aislada de alimentos de origen animal para consumo humano. *Rev. Peru. Med. Exp. Salud Publica* **2016**, 33, 32–44. DOI:10.17843/rpmesp.2016.331.1899
83. de Jesús Cortés-Sánchez A, García-Barrientos R, Minor-Pérez H, Dublán-García O, Martín-Azocar ALS. Food Safety and Antimicrobial Resistance an Approach to the Genus *Salmonella* spp. *J. Biosci. Med.* **2017**, 5, 55–71. DOI:10.4236/jbm.2017.52006
84. Rincón-Gamboa SM, Poutou-Piñales RA, Carrascal-Camacho AK. Antimicrobial Resistance of Non-Typhoid *Salmonella* in Meat and Meat Products. *Foods* **2021**, 10, 1731. DOI:10.3390/foods10081731
85. Ripon RK, Motahara U, Ahmed A, Devnath N, Mahua FA, Hashem RB, et al. Exploring the prevalence of antibiotic resistance patterns and drivers of antibiotics resistance of *Salmonella* in livestock and poultry-derived foods: A systematic review and meta-analysis in Bangladesh from 2000 to 2022. *JAC Antimicrob. Resist.* **2023**, 5, dlad059. DOI:10.1093/jacamr/dlad059
86. Mthembu TP, Zishiri OT, El Zowalaty ME. Genomic Characterization of Antimicrobial Resistance in Food Chain and Livestock-Associated *Salmonella* Species. *Animals* **2021**, 11, 872. DOI:10.3390/ani11030872
87. Siddique A, Azim S, Ali A, Andleeb S, Ahsan A, Imran M, et al. Antimicrobial Resistance Profiling of Biofilm Forming Non Typhoidal *Salmonella enterica* Isolates from Poultry and Its Associated Food Products from Pakistan. *Antibiotics* **2021**, 10, 785. DOI:10.3390/antibiotics10070785
88. Gambino D, Gargano V, Butera G, Sciortino S, Pizzo M, Oliveri G, et al. Food Is Reservoir of MDR *Salmonella*: Prevalence of ESBLs Profiles and Resistance Genes in Strains Isolated from Food. *Microorganisms* **2022**, 10, 780. DOI:10.3390/microorganisms10040780
89. Ren D, Chen P, Wang Y, Wang J, Liu H, Liu H. Phenotypes and antimicrobial resistance genes in *Salmonella* isolated from retail chicken and pork in Changchun, China. *J. Food Saf.* **2017**, 37, e12314. DOI:10.1111/jfs.12314
90. Wang W, Zhao L, Hu Y, Dottorini T, Fanning S, Xu J, et al. Epidemiological study on prevalence, serovar diversity, multidrug resistance, and CTX-M-type extended-spectrum β -lactamases of *Salmonella* spp. from patients with diarrhea, food of animal origin, and pets in several provinces of China. *Antimicrob. Agents Chemother.* **2020**, 64, e00092-20. DOI:10.1128/AAC.00092-20
91. Yang X, Huang J, Wu Q, Zhang J, Yang S, Wang J, et al. Occurrence, serovars and antibiotic resistance of *Salmonella* spp. in retail ready-to-eat food products in some Chinese provinces. *LWT-Food Sci. Technol.* **2022**, 154, 112699. DOI:10.1016/j.lwt.2021.112699
92. Qian M, Xu D, Wang J, Zaeim D, Han J, Qu D. Isolation, antimicrobial resistance, and virulence characterization of *Salmonella* spp. from fresh foods in retail markets in Hangzhou, China. *PLoS ONE* **2023**, 18, e0292621. DOI:10.1371/journal.pone.0292621
93. Rau RB, Ribeiro AR, dos Santos A, Barth AL. Antimicrobial resistance of *Salmonella* from poultry meat in Brazil: Results of a nationwide survey. *Epidemiol. Infect.* **2021**, 49, e228. DOI:10.1017/S0950268821002156
94. Foley SL, Nayak R, Hanning IB, Johnson TJ, Han J, Ricke SC. Population dynamics of *Salmonella enterica* serotypes in commercial egg and poultry production. *Appl. Environ. Microbiol.* **2011**, 77, 4273–4279. DOI:10.1128/AEM.00598-11
95. Molina A, Thye T, Muñoz-Vargas L, Zamora-Sanabria R, Chercos DH, Hernández-Rojas R, et al. Molecular characterization of antibiotic resistant *Salmonella enterica* across the poultry production chain in Costa Rica: A cross-

- sectional study. *Int. J. Food Microbiol.* **2024**, *416*, 110663. DOI:10.1016/j.ijfoodmicro.2024.110663
96. Alvarez DM, Barrón-Montenegro R, Conejeros J, Rivera D, Undurraga EA, Moreno-Switt AI. A review of the global emergence of multidrug-resistant *Salmonella enterica* subsp. *enterica* Serovar Infantis. *Int. J. Food Microbiol.* **2023**, *403*, 110297. DOI:10.1016/j.ijfoodmicro.2023.110297
 97. Alba P, Carfora V, Feltrin F, Diaconu EL, Sorbara L, Dell'Aira E, et al. Evidence of structural rearrangements in ESBL-positive pESI(like) megaplasms of *S. Infantis*. *FEMS Microbiol. Lett.* **2023**, *370*, fnad014. DOI:10.1093/femsle/fnad014
 98. U.S. Department of Health and Human Services. Centers for Disease Control and Prevention (CDC). Antibiotic Resistance Threats in the United States. 2019. Available online: <https://ndc.services.cdc.gov/wp-content/uploads/Antibiotic-Resistance-Threats-in-the-United-States-2019.pdf> (accessed on 28 May 2024).
 99. EFSA (European Food Safety Authority), Beloeil P-A, Guerra B, Stoicescu AV. Manual for reporting on antimicrobial resistance within the framework of Directive 2003/99/EC and Decision 2013/652/EU for information derived from the year 2017. *EFSA J.* **2018**, *15*, 1369E. DOI:10.2903/sp.efsa.2018.EN-1369
 100. Li C, Tate H, Huang X, Hsu CH, Harrison LB, Zhao S, et al. The spread of pESI mediated extended-spectrum cephalosporin resistance in *Salmonella* serovars—Infantis, Senftenberg, and Alachua isolated from food animal sources in the United States. *PLoS ONE* **2024**, *19*, e0299354. DOI:10.1371/journal.pone.0299354
 101. EFSA (European Food Safety Authority), ECDC (European Centre for Disease Prevention and Control). The European Union Summary Report on Antimicrobial Resistance in zoonotic and indicator bacteria from humans, animals and food in 2019–2020. *EFSA J.* **2022**, *20*, 7209. DOI:10.2903/j.efsa.2022.7209
 102. Mattock J, Chattaway MA, Hartman H, Dallman TJ, Smith AM, Keddy K, et al. A One Health Perspective on *Salmonella enterica* Serovar Infantis, an Emerging Human Multidrug-Resistant Pathogen. *Emerg. Infect. Dis.* **2024**, *30*, 701–710. DOI:10.3201/eid3004.231031
 103. Medina-Santana JL, Ortega-Paredes D, de Janon S, Burnett E, Ishida M, Sauders B, et al. Investigating the dynamics of *Salmonella* contamination in integrated poultry companies using a whole genome sequencing approach. *Poultry Sci.* **2022**, *101*, 101611. DOI:10.1016/j.psj.2021.101611
 104. dos Santos AMP, Panzenhagen P, Ferrari RG, Conte-Junior CA. Large-scale genomic analysis reveals the pESI-like megaplasmid presence in *Salmonella* Agona, Muenchen, Schwarzengrund, and Seftenberg. *Food Microbiol.* **2022**, *108*, 104112. DOI:10.1016/j.fm.2022.104112
 105. Valderrama M, Quevedo M, Pastor J, Mantilla Y, Ortiz M. Estudio de la prevalencia de serovares de *Salmonella* en granjas avícolas tecnificadas de Perú. *SENASA* **2014**, *31*. Available online: <https://www.scribd.com/document/264956147/Estudio-sobre-la-salmonella-en-el-Peru> (accessed on 10 August 2026).
 106. Vinuesa-Burgos C, Cevallos M, Ron-Garrido L, Bertrand S, De Zutter L. Prevalence and Diversity of *Salmonella* Serotypes in Ecuadorian Broilers at Slaughter Age. *PLoS ONE* **2016**, *11*, e0159567. DOI:10.1371/journal.pone.0159567
 107. Quino W, Hurtado CV, Escalante-Maldonado O, Flores-León D, Mestanza O, Vences-Rosales F, et al. Multidrug resistance of *Salmonella* Infantis in Peru: A study through next generation sequencing. *Rev. Peru. Med. Exp. Salud Publica* **2019**, *36*, 37–45. DOI:10.17843/rpmesp.2019.361.3934
 108. Mejía L, Medina JL, Bayas R, Salazar CS, Villavicencio F, Zapata S, et al. Genomic Epidemiology of *Salmonella* Infantis in Ecuador: From Poultry Farms to Human Infections. *Front. Vet. Sci.* **2020**, *7*, 547891. DOI:10.3389/fvets.2020.547891
 109. Pardo-Esté C, Lorca D, Castro-Severyn J, Krüger G, Alvarez-Thon L, Zepeda P, et al. Genetic Characterization of *Salmonella* Infantis with Multiple Drug Resistance Profiles Isolated from a Poultry-Farm in Chile. *Microorganisms* **2021**, *9*, 2370. DOI:10.3390/microorganisms9112370
 110. Bogomazova AN, Gordeeva VD, Krylova EV, Soltynskaya IV, Davydova EE, Ivanova OE, et al. Mega-plasmid found worldwide confers multiple antimicrobial resistance in *Salmonella* Infantis of broiler origin in Russia. *Int. J. Food Microbiol.* **2020**, *319*, 108497. DOI:10.1016/j.ijfoodmicro.2019.108497
 111. Di Marcantonio L, Romantini R, Marotta F, Chiaverini A, Zilli K, Abass A, et al. The Current Landscape of Antibiotic Resistance of *Salmonella* Infantis in Italy: The Expansion of Extended-Spectrum Beta-Lactamase Producers on a Local Scale. *Front. Microbiol.* **2022**, *13*, 812481. DOI:10.3389/fmicb.2022.812481
 112. Papić B, Kušar D, Mićunović J, Pirš M, Ocepek M, Avberšek J. Clonal Spread of pESI-Positive Multidrug-Resistant ST32 *Salmonella enterica* Serovar Infantis Isolates among Broilers and Humans in Slovenia. *Microbiol. Spectr.* **2022**, *10*, e02481-22. DOI:10.1128/spectrum.02481-22
 113. Srednik ME, Morningstar-Shaw BR, Hicks JA, Tong C, Mackie TA, Schlater LK. Whole-genome sequencing and phylogenetic analysis capture the emergence of a multi-drug resistant *Salmonella enterica* serovar Infantis clone from diagnostic animal samples in the United States. *Front. Microbiol.* **2023**, *14*, 1166908. DOI:10.3389/fmicb.2023.1166908
 114. Guzinski J, Potter J, Tang Y, Davies R, Teale C, Petrovska L. Geographical and temporal distribution of multidrug resistant *Salmonella* Infantis in Europe and the Americas. *Front. Microbiol.* **2024**, *14*, 1244533. DOI:10.3389/fmicb.2023.1244533

115. Sanguankiat A, Pinniam N, Tulayakul P. Surveillance of antimicrobial resistance, phenotypic, and genotypic patterns of *Salmonella enterica* isolated from animal feedstuffs: Annual Study. *Vet. World* **2023**, *16*, 939–945. DOI:10.14202/vetworld.2023.939-945
116. Li X, Bethune LA, Jia Y, Lovell RA, Proescholdt TA, Benz SA, et al. Surveillance of *Salmonella* Prevalence in Animal Feeds and Characterization of the *Salmonella* Isolates by Serotyping and Antimicrobial Susceptibility. *Foodborne Pathog. Dis.* **2012**, *9*, 692–698. DOI:10.1089/fpd.2011.1083
117. Hsieh Y-C, Poole TL, Runyon M, Hume M, Herrman TJ. Prevalence of Nontyphoidal *Salmonella* and *Salmonella* Strains with Conjugative Antimicrobial-Resistant Serovars Contaminating Animal Feed in Texas. *J. Food Protect.* **2016**, *79*, 194–204. DOI:10.4315/0362-028X.JFP-15-163
118. Parker EM, Valcanis M, Edwards LJ, Anderson P, Mollenkopf DF, Wittum TE. Antimicrobial-resistant *Salmonella* is detected more frequently in feed milling equipment than in raw feed components or processed animal feed. *Aust. Vet. J.* **2022**, *100*, 213–219. DOI:10.1111/avj.13146
119. Carresi C, Marabelli R, Roncada P, Britti D. Is the Use of Monensin Another Trojan Horse for the Spread of Antimicrobial Resistance? *Antibiotics* **2024**, *13*, 129. DOI:10.3390/antibiotics13020129
120. Oh H, Choi Y, Lee J. Antibiotic-Resistant *Salmonella* in Animal Products Jeopardize Human Health. *Food Sci. Anim. Resour.* **2025**, *45*, 409–428. DOI:10.5851/kosfa.2025.e4
121. Molina A, Granados-Chinchilla F, Jiménez M, Acuña-Calvo MT, Alfaro M, Chavarria G. Vigilance for *Salmonella* in Feedstuffs Available in Costa Rica: Prevalence, Serotyping and Tetracycline Resistance of Isolates Obtained from 2009 to 2014. *Foodborne Pathog. Dis.* **2016**, *13*, 119–127. DOI:10.1089/fpd.2015.2050
122. Mdemu S, Mathara JM, Makondo ZE. Antibiotic resistance of *Salmonella* isolated from commercial chicken feeds in Dar es Salaam, Tanzania. *Tanzania Vet. Assoc. Proceed.* **2018**, *35*, 63–69. Available online: <http://tva.or.tz/images/THE%2035%20TVA%20PROCEEDINGS%202017%20PUBLISHED.pdf#page=64> (accessed on 10 August 2026).
123. Ngai DG, Nyamache AK, Ombori O. Prevalence and antimicrobial resistance profiles of *Salmonella* species and *Escherichia coli* isolates from poultry feeds in Ruiru Sub-County, Kenya. *BMC Res. Notes* **2021**, *14*, 41. DOI:10.1186/s13104-021-05456-4
124. Paredes R, Damme M, Mantilla J, Castellanos LR, Clavijo V, Celis Y, et al. Prevalence and antimicrobial resistance of *Escherichia coli* and *Salmonella* spp. in animal feed in Colombia. *Rev. Panam. Salud Publica* **2023**, *47*, e57. DOI:10.26633/RPSP.2023.57
125. Rojas-Sánchez E, Jiménez-Soto M, Barquero-Calvo E, Duarte-Martínez F, Mollenkopf DF, Wittum TE, et al. Prevalence Estimation, Antimicrobial Susceptibility, and Serotyping of *Salmonella enterica* Recovered from New World Non-Human Primates (*Platyrrhini*), Feed, and Environmental Surfaces from Wildlife Centers in Costa Rica. *Antibiotics* **2023**, *12*, 844. DOI:10.3390/antibiotics12050844
126. Lertworapreecha M, Sutthimusik S, Tontikapong K. Antimicrobial Resistance in *Salmonella enterica* Isolated From Pork, Chicken, and Vegetables in Southern Thailand. *Jundishapur J. Microbiol.* **2013**, *6*, 36–41. DOI:10.5812/jjm.4312
127. Taban BM, Aytac SA, Akkoc N, Akcelik M. Characterization of antibiotic resistance in *Salmonella enterica* isolates determined from ready-to-eat (RTE) salad vegetables. *Braz. J. Microbiol.* **2013**, *44*, 385–391. DOI:10.1590/S1517-83822013005000047
128. Umeh SI, Enwuru CP. Antimicrobial Resistance Profile of *Salmonella* Isolates from Livestock. *Open J. Med. Microbiol.* **2014**, *4*, 242–248. DOI:10.4236/ojmm.2014.44027
129. Abakpa GO, Umoh VJ, Ameh JB, Yakubu SE, Kwaga JKP, Kamaruzaman S. Diversity and antimicrobial resistance of *Salmonella enterica* isolated from fresh produce and environmental samples. *Environ. Nanotechnol. Monitor. Manag.* **2015**, *3*, 38–46. DOI:10.1016/j.enmm.2014.11.004
130. Mąka Ł, Maćkiw E, Ścieżyńska H, Popowska M. Occurrence and antimicrobial resistance of *Salmonella* spp. isolated from food other than meat in Poland. *Ann. Agric. Environ. Med.* **2015**, *22*, 405–408. DOI:10.5604/12321966.1167701
131. Najwa MS, Rukayadi Y, Ubong A, Loo YY, Chang WS, Lye YL, et al. Quantification and antibiotic susceptibility of *Salmonella* spp., *Salmonella* Enteritidis and *Salmonella* Typhimurium in raw vegetables (*ulam*). *Int. Food Res. J.* **2015**, *22*, 1761–1769. Available online: <https://research.monash.edu/en/publications/quantification-and-antibiotic-susceptibility-of-salmonella-spp-sa/> (accessed on 10 August 2026).
132. Andoh LA, Dalsgaard A, Obiri-Danso K, Newman MJ, Barco L, Olsen JE. Prevalence and antimicrobial resistance of *Salmonella* serovars isolated from poultry in Ghana. *Epidemiol. Infect.* **2016**, *144*, 3288–3299. DOI:10.1017/S0950268816001126
133. Okorie-Kanu OJ, Ezenduka EV, Okorie-Kanu CO, Ugwu LC, Nnamani UJ. Occurrence and antimicrobial resistance of pathogenic *Escherichia coli* and *Salmonella* spp. in retail raw table eggs sold for human consumption in Enugu state,

- Nigeria. *Vet. World* **2016**, *9*, 1312–1319. DOI:10.14202/vetworld.2016.1312-1319
134. Zishiri OT, Mkhize N, Mukaratirwa S. Prevalence of virulence and antimicrobial resistance genes in *Salmonella* spp. isolated from commercial chickens and human clinical isolates from South Africa and Brazil. *Onderstepoort J. Vet. Res.* **2016**, *83*, a1067. DOI:10.4102/ojvr.v83i1.1067
 135. Deli RA, Adzitey F. Prevalence and antibiotic resistance of *Salmonella* serovars isolated from spent hens and its environmental samples in Penang and Kedah, Malaysia. *J. Trop. Agric. Food Sci.* **2017**, *45*, 37–50. DOI:10.5555/20173282271
 136. Terentjeva M, Avsejenko J, Streikiša M, Utināne A, Kovaļenko K, Bērziņš A. Prevalence and antimicrobial resistance of *Salmonella* in meat and meat products in Latvia. *Ann. Agric. Environ. Med.* **2017**, *24*, 317–321. DOI:10.5604/12321966.1235180
 137. Adzitey F. Antibiotic resistance of *Escherichia coli* and *Salmonella enterica* isolated from cabbage and lettuce samples in Tamale metropolis of Ghana. *Int. J. Food Contam.* **2018**, *5*, 7. DOI:10.1186/s40550-018-0068-z
 138. An HTT, Khai LTL. The prevalence and antibiotic resistance of *Salmonella* spp. isolated from pigs and farm environments in Vinh Long province. *CTU J. Innov. Sustain. Dev.* **2018**, *54*, 26–33. DOI:10.22144/ctu.jsi.2018.091
 139. Hassan MM, Begum S, Al Faruq A, Alam M, Mahmud T, Islam A. Multidrug Resistant *Salmonella* Isolated from Street Foods in Chittagong, Bangladesh. *Microb. Res. J. Int.* **2018**, *26*, 1–8. DOI:10.9734/MRJI/2018/v26i630083
 140. Ketema L, Ketema Z, Kiflu B, Alemayehu H, Terefe Y, Ibrahim M, et al. Prevalence and Antimicrobial Susceptibility Profile of *Salmonella* Serovars Isolated from Slaughtered Cattle in Addis Ababa, Ethiopia. *BioMed Res. Int.* **2018**, *2018*, 9794869. DOI:10.1155/2018/9794869
 141. Pan H, Paudyal N, Li X, Fang W, Yue M. Multiple Food-Animal-Borne Route in Transmission of Antibiotic-Resistant *Salmonella* Newport to Humans. *Front. Microbiol.* **2018**, *9*, 23. DOI:10.3389/fmicb.2018.00023
 142. Akinola SA, Mwanza M, Ateba CN. Occurrence, Genetic Diversities And Antibiotic Resistance Profiles of *Salmonella* Serovars Isolated From Chickens. *Infect. Drug Resist.* **2019**, *12*, 3327–3342. DOI:10.2147/IDR.S217421
 143. Beshiru A, Igbinosa IH, Igbinosa EO. Prevalence of Antimicrobial Resistance and Virulence Gene Elements of *Salmonella* Serovars From Ready-to-Eat (RTE) Shrimps. *Front. Microbiol.* **2019**, *10*, 1613. DOI:10.3389/fmicb.2019.01613
 144. Chen T, Jiang J, Ye C, Xie J, Chen X, Xu D, et al. Genotypic characterization and antimicrobial resistance profile of *Salmonella* isolated from chicken, pork and the environment at abattoirs and supermarkets in Chongqing, China. *BMC Vet. Res.* **2019**, *15*, 456. DOI:10.1186/s12917-019-2202-4
 145. Rattanadilok na Phuket N, Siripornadulsil S, Siripornadulsil W. Prevalence of Antibiotic-resistant *Salmonella* in Vegetables and Fermented Foods and their Control by Lactic Acid Bacteria. *J. Pure Appl. Microbiol.* **2019**, *13*, 1929–1939. DOI:10.22207/JPAM.13.4.05
 146. Wang X, Biswas S, Paudyal N, Pan H, Li X, Fang W, et al. Antibiotic Resistance in *Salmonella* Typhimurium Isolates Recovered From the Food Chain Through National Antimicrobial Resistance Monitoring System Between 1996 and 2016. *Front. Microbiol.* **2019**, *10*, 985. DOI:10.3389/fmicb.2019.00985
 147. Yang X, Wu Q, Zhang J, Huang J, Chen L, Wu S, et al. Prevalence, Bacterial Load, and Antimicrobial Resistance of *Salmonella* Serovars Isolated From Retail Meat and Meat Products in China. *Front. Microbiol.* **2019**, *10*, 2121. DOI:10.3389/fmicb.2019.02121
 148. Benevides V, Rubio M, Alves L, Barbosa F, Souza A, Almeida A, et al. Antimicrobial Resistance in *Salmonella* Serovars Isolated From an Egg-Producing Region in Brazil. *Braz. J. Poult. Sci.* **2020**, *22*, eRBCA-2020. DOI:10.1590/1806-9061-2020-1259
 149. Kapena MS, Muma JB, Mubita CM, Munyeme M. Antimicrobial resistance of *Escherichia coli* and *Salmonella* in raw retail table eggs in Lusaka, Zambia. *Vet. World* **2020**, *13*, 2528–2533. DOI:10.14202/vetworld.2020.2528-2533
 150. Mejía L, Vela G, Zapata S. High Occurrence of Multiresistant *Salmonella* Infantis in Retail Meat in Ecuador. *Foodborne Pathog. Dis.* **2021**, *18*, 41–48. DOI:10.1089/fpd.2020.2808
 151. Manishimwe R, Moncada PM, Musanayire V, Shyaka A, Scott HM, Loneragan GH. Antibiotic-Resistant *Escherichia coli* and *Salmonella* from the Feces of Food Animals in the East Province of Rwanda. *Animals* **2021**, *11*, 1013. DOI:10.3390/ani11041013
 152. Nguyen TK, Nguyen LT, Chau TTH, Nguyen TT, Tran BN, Taniguchi T, et al. Prevalence and antibiotic resistance of *Salmonella* isolated from poultry and its environment in the Mekong Delta, Vietnam. *Vet. World* **2021**, *14*, 3216–3223. DOI:10.14202/vetworld.2021.3216-3223
 153. Alessiani A, Goffredo E, Mancini M, Occhiochiuso G, Faleo S, Didonna A, et al. Evaluation of Antimicrobial Resistance in *Salmonella* Strains Isolated from Food, Animal and Human Samples between 2017 and 2021 in Southern Italy. *Microorganisms* **2022**, *10*, 812. DOI:10.3390/microorganisms10040812
 154. Lauteri C, Festino AR, Conter C, Vergara A. Prevalence and antimicrobial resistance profile in *Salmonella* spp. isolates

- from swine food chain. *Ital. J. Food Saf.* **2022**, *11*, 9980. DOI:10.4081/ijfs.2022.9980
155. Mir R, Salari S, Najimi M, Rashki A. Determination of frequency, multiple antibiotic resistance index and resistotype of *Salmonella* spp. in chicken meat collected from southeast of Iran. *Vet. Med. Sci.* **2022**, *8*, 229–236. DOI:10.1002/vms3.647
 156. Rakitin AL, Yushina YK, Zaiko EV, Bataeva DS, Kuznetsova OA, Semenova AA, et al. Evaluation of Antibiotic Resistance of *Salmonella* Serotypes and Whole-Genome Sequencing of Multiresistant Strains Isolated from Food Products in Russia. *Antibiotics* **2022**, *11*, 1. DOI:10.3390/antibiotics11010001
 157. Retamal P, Gaspar J, Benavides MB, Saenz L, Galarce N, Aravena T, et al. Virulence and antimicrobial resistance factors in *Salmonella enterica* serotypes isolated from pigs and chickens in central Chile. *Front. Vet. Sci.* **2022**, *9*, 971246. DOI:10.3389/fvets.2022.971246
 158. Russo I, Bencardino D, Napoleoni M, Andreoni F, Schiavano GF, Baldelli G, et al. Prevalence, Antibiotic-Resistance, and Replicon-Typing of *Salmonella* Strains among Serovars Mainly Isolated from Food Chain in Marche Region, Italy. *Antibiotics* **2022**, *11*, 725. DOI:10.3390/antibiotics11060725
 159. Siddiky NA, Sarker S, Khan SR, Rahman T, Kafi A, Samad MA. Virulence and antimicrobial resistance profile of non-typhoidal *Salmonella enterica* serovars recovered from poultry processing environments at wet markets in Dhaka, Bangladesh. *PLoS ONE* **2022**, *17*, e0254465. DOI:10.1371/journal.pone.0254465
 160. Barrera S, Vázquez-Flores S, Needle D, Rodríguez-Medina N, Iglesias D, Sevigny JL, et al. Serovars, Virulence and Antimicrobial Resistance Genes of Non-Typhoidal *Salmonella* Strains from Dairy Systems in Mexico. *Antibiotics* **2023**, *12*, 1662. DOI:10.3390/antibiotics12121662
 161. Ganjeer T, Patyal A, Shakya S, Chandrakar C, Verma SK, Gade NE, et al. Antibiotic Resistance Pattern and Distribution of Resistance Genes in *Salmonella* Isolated from Chicken and Duck Eggs in Chhattisgarh, India. *Asian J. Dairy Food Res.* **2023**, *1*, 1–8. DOI:10.18805/ajdrf.DR-2033
 162. Huong CTT, Ngoc PT, Thai TH. Prevalence and antibiotic resistance of *Salmonella* spp. and *Escherichia coli* isolated from poultry farms in the North Vietnam. *J. Dairy Vet. Anim. Res.* **2023**, *12*, 70–75. DOI:10.15406/jdvar.2023.12.00326
 163. Mumbo MT, Nyaboga EN, Kinyua JK, Muge EK, Mathenge SGK, Rotich H, et al. Antimicrobial resistance profiles of *Salmonella* spp. and *Escherichia coli* isolated from fresh Nile tilapia (*Oreochromis niloticus*) fish marketed for human consumption. *BMC Microbiol.* **2023**, *23*, 306. DOI:10.1186/s12866-023-03049-8
 164. Penjor K, Guring M, Islam K, Magalhaes RJS. Prevalence and Antibiotic Resistance Profile of *Salmonella* spp. in Broiler Carcasses from Dominant Poultry Production Areas in Bhutan. *J. Infect. Dis. Microbiol.* **2023**, *1*, 1–11. DOI:10.37191/Mapsci-JIDM-1(3)-013
 165. Sabri C, Karraouan B, Berradi T, Bouchrif B, El Mellouli F, Ziyate N, et al. Prevalence and antimicrobial susceptibility profile of *Salmonella* isolated from food products in the region of Casablanca, Morocco. *J. Infect. Dev. Ctries.* **2023**, *17*, 388–396. DOI:10.3855/jidc.17236
 166. Vázquez X, García-Fierro R, Fernández J, Bances M, Herrero-Fresno A, Olsen JE, et al. Incidence and Genomic Background of Antibiotic Resistance in Food-Borne and Clinical Isolates of *Salmonella enterica* Serovar Derby from Spain. *Antibiotics* **2023**, *12*, 1204. DOI:10.3390/antibiotics12071204
 167. Xiao Q, Yu Y, Xu B, Fang Z, Chen W, Feng J, et al. Serotype and antimicrobial resistance of *Salmonella* from poultry meats in 2021 in Shanghai, China. *Food Agric. Immunol.* **2023**, *34*, 2220568. DOI:10.1080/09540105.2023.2220568
 168. Walsh D, Duffy G, Sheridan JJ, Blair IS, McDowell DA. Antibiotic resistance among *Listeria*, including *Listeria monocytogenes*, in retail foods. *J. Appl. Microbiol.* **2001**, *90*, 517–522. DOI:10.1046/j.1365-2672.2001.01273.x
 169. Getaneh A, Berhanu L, Gume B, Deneke Y, Kassa T, Dadi LS, et al. Occurrences and antibiotic susceptibility patterns of *Listeria monocytogenes* in raw meat samples from abattoir and butcher shops in Jimma Town, Southwest Ethiopia. *Heliyon* **2025**, *11*, e42589. DOI:10.1016/j.heliyon.2025.e42589
 170. Lungu B, O'Bryan CA, Muthaiyan A, Milillo SR, Johnson MG, Crandall PG, et al. *Listeria monocytogenes*: Antibiotic Resistance in Food Production. *Foodborne Pathog. Dis.* **2011**, *8*, 569–578. DOI:10.1089/fpd.2010.0718
 171. Byahut A, Das M, Chowdhury G, Mukhopadhyay AK, Khatri R, Das S, et al. Exploring the prevalence and antibiotic resistance of *Listeria monocytogenes* in diverse food commodities across Sikkim, India. *Braz. J. Microbiol.* **2025**, *57*, 23. DOI:10.1007/s42770-025-01853-0
 172. Anupama A, Pattapulavar V, Christopher JG. The past, present, future of *Listeria monocytogenes*: Understanding the molecular pathways, antibiotic resistance, and public health implications. *Med. Macroecol.* **2025**, *25*, 100127. DOI:10.1016/j.medmic.2025.100127
 173. Luque-Sastre L, Arroyo C, Fox EM, McMahon BJ, Bai L, Li F, et al. Antimicrobial Resistance in *Listeria* species. *Microbiol. Spectr.* **2018**, *6*, ARBA-0031-2017. DOI:10.1128/microbiolspec.ARBA-0031-2017
 174. Hyden P, Pietzka A, Lennkh A, Murer A, Springer B, Blaschitz M, et al. Whole genome sequence based serogrouping of *Listeria monocytogenes* isolates. *J. Biotechnol.* **2016**, *235*, 181–186. DOI:10.1016/j.jbiotec.2016.06.005

175. Gorski L, Aviles Noriega A, Carter MQ. *Listeria monocytogenes* serotype 4b strains demonstrate a fitness advantage over strains of serotypes 1/2a and 4b-v1 on cantaloupe rind and flesh at different temperatures. *Microbiol. Spectr.* **2025**, *13*, e01252-25. DOI:10.1128/spectrum.01252-25
176. Guel-García P, García De León FJ, Aguilera-Arreola G, Mandujano A, Mireles-Martínez M, Oliva-Hernández A, et al. Prevalence and Antimicrobial Resistance of *Listeria monocytogenes* in Different Raw Food from Reynosa, Tamaulipas, Mexico. *Foods* **2024**, *13*, 1656. DOI:10.3390/foods13111656
177. TiRziu E, Herman V, Nichita I, Morar A, Imre M, Ban-Cucerzan A, et al. Diversity and Antibiotic Resistance Profiles of *Listeria monocytogenes* Serogroups in Different Food Products from the Transylvania Region of Central Romania. *J. Food Prot.* **2022**, *85*, 54–59. DOI:10.4315/JFP-21-172
178. Duma MN, Ciupescu LM, Dan SD, Crisan-Reget OL, Tabaran A. Virulence and Antimicrobial Resistance of *Listeria monocytogenes* Isolated from Ready-to-Eat Food Products in Romania. *Microorganisms* **2024**, *12*, 954. DOI:10.3390/microorganisms12050954
179. Rippa A, Bilei S, Peruzzy MF, Marrocco MG, Leggeri P, Bossù T, et al. Antimicrobial Resistance of *Listeria monocytogenes* Strains Isolated in Food and Food-Processing Environments in Italy. *Antibiotics* **2024**, *13*, 525. DOI:10.3390/antibiotics13060525
180. Altuntas EG, Kocan D, Cosansu S, Ayhan K, Juneja VK, Materon L. Antibiotic and Bacteriocin Sensitivity of *Listeria monocytogenes* Strains Isolated from Different Foods. *Food Nutr. Sci.* **2012**, *3*, 363–368. DOI:10.4236/fns.2012.33052
181. Moura A, Leclercq A, Vales G, Tessaud-Rita N, Bracq-Dieye H, Thouvenot P, et al. Phenotypic and genotypic antimicrobial resistance of *Listeria monocytogenes*: An observational study in France. *Lancet Reg. Health Eur.* **2024**, *37*, 100800. DOI:10.1016/j.lanepe.2023.100800
182. Hanes RM, Huang Z. Investigation of Antimicrobial Resistance Genes in *Listeria monocytogenes* from 2010 through to 2021. *Int. J. Environ. Res. Public Health* **2022**, *19*, 5506. DOI:10.3390/ijerph19095506
183. Parra-Flores J, Holý O, Bustamante F, Lepuschitz S, Pietzka A, Contreras-Fernández A, et al. Virulence and Antibiotic Resistance Genes in *Listeria monocytogenes* Strains Isolated From Ready-to-Eat Foods in Chile. *Front. Microbiol.* **2022**, *12*, 796040. DOI:10.3389/fmicb.2021.796040
184. Wiśniewski P, Adamski P, Trymers M, Chajęcka-Wierzchowska W, Zadernowska A. Predicting Antibiotic Resistance in *Listeria monocytogenes* from Food and Food-Processing Environments Using Next-Generation Sequencing: A Systematic Review. *Int. J. Mol. Sci.* **2025**, *26*, 10112. DOI:10.3390/ijms262010112
185. Anne du Toit S, Rip D. Exploring the genetic variability, virulence factors, and antibiotic resistance of *Listeria monocytogenes* from fresh produce, ready-to-eat hummus, and food-processing environments. *J. Food Sci.* **2024**, *89*, 6916–6945. DOI:10.1111/1750-3841.17399
186. Wang H, Feng M, Anwar TM, Chai W, Ed-Dra A, Kang X, et al. Change in antimicrobial susceptibility of *Listeria* spp. in response to stress conditions. *Front. Sustain. Food Syst.* **2023**, *7*, 1179835. DOI:10.3389/fsufs.2023.1179835
187. Marmion M, Macori G, Ferone M, Whyte P, Scannell AGM. Survive and thrive: Control mechanisms that facilitate bacterial adaptation to survive manufacturing-related stress. *Int. J. Food Microbiol.* **2022**, *368*, 109612. DOI:10.1016/j.ijfoodmicro.2022.109612
188. Sakaridis I, Soultos N, Iossifidou E, Papa A, Ambrosiadis I, Koidis P. Prevalence and antimicrobial resistance of *Listeria monocytogenes* isolated in chicken slaughterhouses in Northern Greece. *J. Food Prot.* **2011**, *74*, 1017–1021. DOI:10.4315/0362-028X.JFP-10-545
189. Keet R, Rip D. *Listeria monocytogenes* isolates from Western Cape, South Africa exhibit resistance to multiple antibiotics and contradicts certain global resistance patterns. *AIMS Microbiol.* **2021**, *7*, 40–58. DOI:10.3934/microbiol.2021004
190. Ramatla T, Khasapane G, Achilonu CC, Tshikantwa TS, Nkhebenyane J, Lekota KE, et al. “One Health” perspective on the prevalence of *Listeria monocytogenes* and its antibiotic resistance profiles in South Africa: A comprehensive systematic review and meta-analysis. *Sci. Rep.* **2026**, *16*, 7680. DOI:10.1038/s41598-025-34005-3
191. Sarr M, Alou MT, Padane A, Diouf FS, Beye M, Sokhna C, et al. A review of the literature of *Listeria monocytogenes* in Africa highlights breast milk as an overlooked human source. *Front. Microbiol.* **2023**, *14*, 1213953. DOI:10.3389/fmicb.2023.1213953
192. Threlfall EJ, Skinner JA, McLauchlin J. Antimicrobial resistance in *Listeria monocytogenes* from humans and food in the UK, 1967–1996. *Clin. Microbiol. Infect.* **1998**, *4*, 410–412. DOI:10.1111/j.1469-0691.1998.tb00087.x
193. Escolar C, Gómez D, del Carmen Rota García M, Conchello P, Herrera A. Antimicrobial Resistance Profiles of *Listeria monocytogenes* and *Listeria innocua* Isolated from Ready-to-Eat Products of Animal Origin in Spain. *Foodborne Pathog. Dis.* **2017**, *14*, 357–363. DOI:10.1089/fpd.2016.2248
194. Kayode AJ, Okoh AI. Antibiotic Resistance Profile of *Listeria monocytogenes* Recovered from Ready-to-Eat Foods Surveyed in South Africa. *J. Food Prot.* **2022**, *85*, 1807–1814. DOI:10.4315/JFP-22-090

195. Gana J, Gcebe N, Moerane R, Ngoshe Y, Tshuma T, Moabelo K, et al. Antimicrobial Resistance Profiles of *Listeria* Species Recovered from Retail Outlets in Gauteng Province, South Africa. *J. Food Prot.* **2024**, *87*, 100322. DOI:10.1016/j.jfp.2024.100322
196. Hailu T, Gugsu G, Tsegaye Y, Ahmed M, Awol N. Occurrence and Antimicrobial Resistance Pattern of *Listeria monocytogenes* Isolated From Bovine's Milk and Meat in Mekelle City, Ethiopia. *Adv. Public Health* **2024**, *2024*, 8813787. DOI:10.1155/2024/8813787
197. Panera-Martínez S, Capita R, Pedriza-González Á, Diez-Moura M, Riesco-Peláez F, Alonso-Calleja C. Occurrence, Antibiotic Resistance and Biofilm-Forming Ability of *Listeria monocytogenes* in Chicken Carcasses and Cuts. *Foods* **2024**, *13*, 3822. DOI:10.3390/foods13233822
198. Mtshali NC, Gcebe N, Moerane R, Adesiyun AA. Prevalence, Molecular Profile and Antibiotic Resistance of *Listeria Species* in Retail Beef Products in North-West Province, South Africa: A Cross-Sectional Analysis. *Vet. Med. Sci.* **2025**, *11*, e70680. DOI:10.1002/vms3.70680
199. Nikolaou FG, Colobatiu LM, Ciupescu LM, Tabaran A, Hategan AR, Mihaiu R, et al. Prevalence and Antimicrobial Resistance of *Listeria monocytogenes* Isolated from Dairy Products in Romania. *Antibiotics* **2025**, *14*, 482. DOI:10.3390/antibiotics14050482
200. Sołtysiuk M, Przyborowska P, Wiszniewska-Łaszczych A, Tobolski D. Prevalence and antimicrobial resistance profile of *Listeria* spp. isolated from raw fish. *BMC Vet. Res.* **2025**, *21*, 333. DOI:10.1186/s12917-025-04792-y
201. Kaakoush NO, Castaño-Rodríguez N, Mitchell HM, Man SM. Global Epidemiology of *Campylobacter* Infection. *Clin. Microbiol. Rev.* **2015**, *28*, 687–704. DOI:10.1128/CMR.00006-15
202. Igwaran A, Okoh AI. Human campylobacteriosis: A public health concern of global importance. *Heliyon* **2019**, *5*, e02814. DOI:10.1016/j.heliyon.2019.e02814
203. Sahin O, Shen Z, Zhang Q. Methods to Study Antimicrobial Resistance in *Campylobacter jejuni*. In *Methods in Molecular Biology*; Butcher J, Stintzi A, Eds.; Humana Press: New York, NY, USA, 2017; Volume 1512; pp. 29–42. DOI:10.1007/978-1-4939-6536-6_4
204. Iovine NM. Resistance Mechanisms in *Campylobacter jejuni*. *Virulence* **2013**, *4*, 230–240. DOI:10.4161/viru.23753
205. Shen Z, Wang Y, Zhang Q, Shen J. Antimicrobial Resistance in *Campylobacter* spp. *Microbiol. Spectr.* **2017**, *6*, 10-1128. DOI:10.1128/microbiolspec.ARBA-0013-2017
206. Tang Y, Fang L, Xu C, Zhang Q. Antibiotic resistance trends and mechanisms in the foodborne pathogen, *Campylobacter*. *Anim. Health Res. Rev.* **2017**, *18*, 87–98. DOI:10.1017/S1466252317000135
207. Aleksić E, Miljković-Selimović B, Tambur Z, Aleksić N, Biočanin V, Avramov S. Resistance to Antibiotics in Thermophilic *Campylobacters*. *Front. Med.* **2021**, *8*, 763434. DOI:10.3389/fmed.2021.763434
208. Schreyer ME, Olivero CR, Rossler E, Soto LP, Frizzo LS, Zimmermann JA, et al. Prevalence and antimicrobial resistance of *Campylobacter jejuni* and *C. coli* identified in a slaughterhouse in Argentina. *Curr. Res. Food Sci.* **2022**, *5*, 590–597. DOI:10.1016/j.crfs.2022.03.005
209. Maćkiw E, Korsak D, Rzewuska K, Tomczuk K, Rozynek E. Antibiotic resistance in *Campylobacter jejuni* and *Campylobacter coli* isolated from food in Poland. *Food Control* **2012**, *23*, 297–301. DOI:10.1016/j.foodcont.2011.08.022
210. Paintsil EK, Ofori LA, Akenten CW, Zautner AE, Mbwana J, Jaeger A, et al. Antibiotic-resistant *Campylobacter coli* and *Campylobacter jejuni* in commercial and smallholder farm animals in the Asante Akim North Municipality of Ghana. *Front. Microbiol.* **2022**, *13*, 983047. DOI:10.3389/fmicb.2022.983047
211. Chen S. The culmination of multidrug-resistant efflux pumps vs. meager antibiotic arsenal era: Urgent need for an improved new generation of EPIs. *Front. Microbiol.* **2023**, *14*, 1149418. DOI:10.3389/fmicb.2023.1149418
212. Oh E, Jeon B. Contribution of surface polysaccharides to the resistance of *Campylobacter jejuni* to antimicrobial phenolic compounds. *J. Antibiot.* **2015**, *68*, 591–593. DOI:10.1038/ja.2015.26
213. Efimochkina NR, Stetsenko VV, Sheveleva SA. Formation of the Resistance of *Campylobacter jejuni* to Macrolide Antibiotics. *Bull. Exp. Biol. Med.* **2020**, *169*, 351–356. DOI:10.1007/s10517-020-04885-8
214. Davin-Regli A, Pages JM, Ferrand A. Clinical Status of Efflux Resistance Mechanisms in Gram-Negative Bacteria. *Antibiotics* **2021**, *10*, 1117. DOI:10.3390/antibiotics10091117
215. Shrif S, Bakhshi B, Najar-Peerayeh S. Significant contribution of the CmeABC Efflux pump in high-level resistance to ciprofloxacin and tetracycline in *Campylobacter jejuni* and *Campylobacter coli* clinical isolates. *Ann. Clin. Microbiol. Antimicrob.* **2021**, *20*, 36. DOI:10.1186/s12941-021-00439-6
216. Narvaez-Bravo C, Taboada EN, Mutschall SK, Aslam M. Epidemiology of antimicrobial resistant *Campylobacter* spp. isolated from retail meats in Canada. *Int. J. Food Microbiol.* **2017**, *253*, 43–47. DOI:10.1016/j.ijfoodmicro.2017.04.019
217. Paravisi M, Laviniki V, Bassani J, Kunert Filho H, Carvalho D, Wilsman D, et al. Antimicrobial Resistance in *Campylobacter jejuni* Isolated from Brazilian Poultry Slaughterhouses. *Braz. J. Poultry Sci.* **2020**, *22*, eRBCA-2020.

DOI:10.1590/1806-9061-2020-1262

218. Iovine NM, Blasser MJ. Antibiotics in Animal Feed and Spread of Resistant *Campylobacter* from Poultry to Humans. *Emerg. Infect. Dis.* **2004**, *10*, 1158–1159. DOI:10.3201/eid1006.040403
219. Yang Y, Feye KM, Shi Z, Pavlidis HO, Kogut M, J. Ashworth A, et al. A Historical Review on Antibiotic Resistance of Foodborne *Campylobacter*. *Front. Microbiol.* **2019**, *10*, 1509. DOI:10.3389/fmicb.2019.01509
220. Kiran A, Alvarez MJR, Han KSS, Kumar A, Chakraborty D, Magisha L, et al. The notorious case of *Campylobacter* spp.: An increasingly antimicrobial resistant foodborne pathogen. *AFFIDIA J. Food Diagn.* **2021**, *3*, 32–41. Available online: <https://www.amr-insights.eu/wp-content/uploads/2021/10/AFFIDIA-Vol3-Nr2-AMRITANJALI-et-al.pdf> (accessed on 10 August 2026).
221. Zhang Q, Beyi AF, Yin Y. Zoonotic and antibiotic-resistant *Campylobacter*: A view through the One Health lens. *One Health Adv.* **2023**, *1*, 4. DOI:10.1186/s44280-023-00003-1
222. Agbankpe AJ, Koughblenou SD, Dougnon TV, Oussou A, Gbotche E, Koudokpon CH, et al. Prevalence and Antimicrobial Resistance of *Campylobacter coli* and *Campylobacter jejuni* Isolated from Pig Guts, Pig Feces, and Surface Swabs from the Cutting Tables at Slaughterhouse and Taverns in Southern Benin. *Int. J. Microbiol.* **2022**, *2022*, 5120678. DOI:10.1155/2022/5120678
223. Moradi F, Akbari M, Zandi H, Jahromi RR. Prevalence and Antimicrobial Resistance of *Campylobacter coli* and *Campylobacter jejuni* in the Animals, Food Products, and Human Clinical Specimens in Iran During 2004–2017: A Review Study. *Jundishapur J. Health Sci.* **2020**, *12*, e108609. DOI:10.5812/jjhs.108609
224. European Food Safety Authority [EFSA]. The European Union summary report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2022–2023. *EFSA J.* **2025**, *23*, e9237. DOI:10.2903/j.efsa.2025.9237
225. Portes AB, Panzenhagen P, Pereira dos Santos AM, Conte Junior CA. Antibiotic Resistance in *Campylobacter*: A Systematic Review of South American Isolates. *Antibiotics* **2023**, *12*, 548. DOI:10.3390/antibiotics12030548
226. Hlashwayo DF, Sigaúque B, Bila CG. Epidemiology and antimicrobial resistance of *Campylobacter* spp. in animals in Sub-Saharan Africa: A systematic review. *Heliyon* **2020**, *6*, e03537. DOI:10.1016/j.heliyon.2020.e03537
227. Wanja DW, Mbuthia PG, Bebora LC, Aboge GO, Ogoti B. Antimicrobial Usage, Susceptibility Profiles, and Resistance Genes in *Campylobacter* Isolated from Cattle, Chicken, and Water Samples in Kajiado County, Kenya. *Int. J. Microbiol.* **2023**, *2023*, 8394605. DOI:10.1155/2023/8394605
228. Granados-Chinchilla F, Rodríguez C. Tetracyclines in Food and Feedingstuffs: From Regulation to Analytical Methods, Bacterial Resistance, and Environmental and Health Implications. *J. Anal. Methods Chem.* **2017**, *2017*, 1315497. DOI:10.1155/2017/1315497
229. di Cerbo A, Pezzuto F, Guindetti G, Canello S, Corsi L. Tetracyclines: Insights and Updates of their Use in Human and Animal Pathology and their Potential Toxicity. *Open Biochem. J.* **2019**, *13*, 1–12. DOI:10.2174/1874091X01913010001
230. Przeniosło-Siwczyńska M, Patyra E, Grelik A, Chylek-Purchała M, Kozak B, Kwiatek K. Contamination of Animal Feed with Undeclared Tetracyclines—Confirmatory Analysis by Liquid Chromatography–Mass Spectrometry after Microbiological Plate Test. *Molecules* **2020**, *25*, 2162. DOI:10.3390/molecules25092162
231. Morello S, Pederiva S, Avolio R, Amato G, Zoppi S, Di Blasio A, et al. Tetracyclines in Processed Animal Proteins: A Monitoring Study on Their Occurrence and Antimicrobial Activity. *Foods* **2021**, *10*, 696. DOI:10.3390/foods10040696
232. Shrestha RD, Agunos A, Gow SP, Varga C. Assessing antimicrobial resistance in *Campylobacter jejuni* and *Campylobacter coli* and its association with antimicrobial use in Canadian turkey flocks. *Epidemiol. Infect.* **2023**, *151*, e152. DOI:10.1017/S0950268823001462
233. Gradmann C. Re-Inventing Infectious Disease: Antibiotic Resistance and Drug Development at the Bayer Company 1945–1980. *Med. His.* **2016**, *60*, 155–180. DOI:10.1017/mdh.2016.2
234. Gharbi M, Kamoun S, Hkimi C, Ghedira K, Béjaoui A, Maarroufi A. Relationships between Virulence Genes and Antibiotic Resistance Phenotypes/Genotypes in *Campylobacter* spp. Isolated from Layer Hens and Eggs in the North of Tunisia: Statistical and Computational Insights. *Foods* **2022**, *11*, 3554. DOI:10.3390/foods11223554
235. Byambajav Z, Bulgan E, Hirai Y, Nakayama M, Tanaka M, Nitta Y, et al. Research Note: Antimicrobial resistance of *Campylobacter* species isolated from chickens near Ulaanbaatar city, Mongolia. *Poultry Sci.* **2021**, *100*, 100916. DOI:10.1016/j.psj.2020.11.079
236. Elhadidy M, Ali MM, El-Shibiny A, Miller WG, Elkatib WF, Botteldoorn N, et al. Antimicrobial resistance patterns and molecular resistance markers of *Campylobacter jejuni* isolates from human diarrheal cases. *PLoS ONE* **2020**, *15*, e0227833. DOI:10.1371/journal.pone.0227833
237. Rodrigues JA, Blankenship HM, Cha W, Mukherjee S, Sloup RE, Rudrik JT, et al. Pangenomic analyses of antibiotic-resistant *Campylobacter jejuni* reveal unique lineage distributions and epidemiological associations. *Microb. Genom.* **2023**, *9*, 001073. DOI:10.1099/mgen.0.001073

238. Zachariah OH, Lizzy MA, Rose K, Angela MM. Multiple drug resistance of *Campylobacter jejuni* and *Shigella* isolated from diarrhoeic children at Kapsabet County referral hospital, Kenya. *BMC Infect. Dis.* **2021**, *21*, 109. DOI:10.1186/s12879-021-05788-3
239. Wiczorek K, Szewczyk R, Osek J. Prevalence, antimicrobial resistance, and molecular characterization of *Campylobacter jejuni* and *C. coli* isolated from retail raw meat in Poland. *Vet. Med.* **2012**, *57*, 293–299. DOI:10.17221/6016-VETMED
240. Guyard-Nicodème M, Rivoal K, Houard E, Rose V, Quesne S, Mourand G, et al. Prevalence and characterization of *Campylobacter jejuni* from chicken meat sold in French retail outlets. *Int. J. Food Microbiol.* **2015**, *203*, 8–14. DOI:10.1016/j.ijfoodmicro.2015.02.013
241. Olatoye O, Ogunsemoyin O. Prevalence and Antibiotics Resistance of *Campylobacter jejuni* in Retail Chickens in Oyo State, Nigeria. *Food Sci. Qual. Manag.* **2016**, *48*, 7–11. Available online: <https://www.iiste.org/Journals/index.php/FSQM/article/download/28672/29434> (accessed on 10 August 2026).
242. Jorgensen F, Kesby M, Aranega-Bou P, Willis C, Airo H, Lai S, et al. A Survey of Microbiological and Antimicrobial Resistance (AMR) Contamination Found in Raw Dog and Cat Food on Retail Sale in the UK. Available online: <https://science.food.gov.uk/article/155340-a-survey-of-microbiological-and-antimicrobial-resistance-amr-contamination-found-in-raw-dog-and-cat-food-on-retail-sale-in-the-uk> (accessed on 10 February 2026).
243. Salem LMA, Nashwa OK, Barakat AMA, Mona SA. Antimicrobial resistance of *Campylobacter jejuni* isolated from chicken, some animal products and human in Kalyoubia, Egypt, with special reference to its viability. *Benha Vet. Med. J.* **2019**, *36*, 156–163. DOI:10.21608/bvmj.2019.103407
244. Varga C, Guerin MT, Brash ML, Slavic D, Boerlin P, Susta L. Antimicrobial resistance in *Campylobacter jejuni* and *Campylobacter coli* isolated from small poultry flocks in Ontario, Canada: A two-year surveillance study. *PLoS ONE* **2019**, *14*, e221429. DOI:10.1371/journal.pone.0221429
245. Linn KZ, Furuta M, Nakayama M, Masuda Y, Honjoh K-I, Miyamoto T. Characterization and antimicrobial resistance of *Campylobacter jejuni* and *Campylobacter coli* isolated from chicken and pork. *Int. J. Food Microbiol.* **2021**, *360*, 109440. DOI:10.1016/j.ijfoodmicro.2021.109440
246. Conesa A, Garofolo G, Di Pasquale A, Camma C. Monitoring AMR in *Campylobacter jejuni* from Italy in the last 10 years (2011–2021): Microbiological and WGS data risk assessment. *EFSA J.* **2022**, *20*, e200406. DOI:10.2903/j.efsa.2022.e20
247. Mohamed M-YI, Abu J, Aziz SA, Zakaria Z, Khan AR, Habib I. Occurrence of antibiotic resistant *C. jejuni* and *E. coli* in wild birds, chickens, humans, and the environment in Malay villages, Kedah, Malaysia. *Vet. Med.* **2022**, *67*, 298–308. DOI:10.17221/102/2021-VETMED
248. Tedersoo T, Roasto M, Mäesaar M, Häkkinen L, Kisand V, Ivanova M, et al. Antibiotic Resistance in *Campylobacter* spp. Isolated from Broiler Chicken Meat and Human Patients in Estonia. *Microorganisms* **2022**, *10*, 1067. DOI:10.3390/microorganisms10051067
249. Amjad M, Hassan MM, Raza S, Nisar M, Ahmad M, Zia U-U-R. Prevalence, Risk Factors Analysis and Antimicrobial Resistance Profile of *Campylobacter* Species Isolated from Food Animals in District Okara, Pakistan. *Eur. J. Agric Food Sci.* **2023**, *5*, 61–65. DOI:10.24018/ejfood.2023.5.1.643
250. Bendahou A, Lebbadi M, Ennane L, Essadqui FZ, Abid M. Characterization of *Staphylococcus* species isolated from raw milk and milk products (Iben and jben) in North Morocco. *J. Infect. Dev. Ctries.* **2008**, *2*, 218–225. DOI:10.3855/jidc.266
251. Osman KM, Amer AM, Badr JM, Helmy NM, Elhelw RA, Orabi A, et al. Antimicrobial Resistance, Biofilm Formation and *mecA* Characterization of Methicillin-Susceptible *S. aureus* and Non-*S. aureus* of Beef Meat Origin in Egypt. *Front. Microbiol.* **2016**, *7*, 222. DOI:10.3389/fmicb.2016.00222
252. Ahmad-Mansour N, Loubet P, Pouget C, Dunyach-Remy C, Sotto A, Lavigne J-P, et al. *Staphylococcus aureus* Toxins: An Update on Their Pathogenic Properties and Potential Treatments. *Toxins* **2021**, *13*, 677. DOI:10.3390/toxins13100677
253. da Silva AC, Rodrigues MX, Silva NCC. Methicillin-resistant *Staphylococcus aureus* in food and the prevalence in Brazil: A review. *Braz. J. Microbiol.* **2020**, *51*, 347–356. DOI:10.1007/s42770-019-00168-1
254. Nunes Nascimento JC, Alvarenga VO, Paulino BN, de Cerqueira e Silva AB, Matos JR, Rosário ILDS, et al. *Staphylococcus aureus* in food safety: Antimicrobial resistance, detection technologies, and future perspectives. *Crit. Rev. Food Sci. Nutr.* **2025**, *66*, 607–630. DOI:10.1080/10408398.2025.2528165
255. Castro A, Silva J, Teixeira P. *Staphylococcus aureus*, a Food Pathogen: Virulence Factors and Antibiotic Resistance. In *Foodborne Diseases*; Academic Press: Cambridge, MA, USA, 2018. DOI:10.1016/B978-0-12-811444-5.00008-7
256. Jia K, Fang T, Wang X, Liu Y, Sun W, Wang Y, et al. Antibiotic Resistance Patterns of *Staphylococcus aureus* Isolates from Retail Foods in Mainland China: A Meta-Analysis. *Foodborne Pathog. Dis.* **2020**, *17*, 296–307. DOI:10.1089/fpd.2019.2686
257. Chieffi D, Fanelli F, Fusco V. Antimicrobial and biocide resistance in *Staphylococcus aureus*: Genomic features, decontamination strategies, and the role of *S. aureus* complex-related species, with a focus on ready-to-eat food and food-

- contact surfaces. *Front. Food. Sci. Technol.* **2023**, *3*, 1165871. DOI:10.3389/frfst.2023.1165871
258. Di Bella S, Marini B, Stroffolini G, Geremia N, Giacobbe DR, Campanile F, et al. The virulence toolkit of *Staphylococcus aureus*: A comprehensive review of toxin diversity, molecular mechanisms, and clinical implications. *Eur. J. Clin. Microbiol. Infect. Dis.* **2025**, *44*, 1797–1816. DOI:10.1007/s10096-025-05148-y
 259. Viana GGF, Cardozo MV, Pereira JG, Rossi GAM. Antimicrobial Resistant *Staphylococcus* spp., *Escherichia coli*, and *Salmonella* spp. in Food Handlers: A Global Review of Persistence, Transmission, and Mitigation Challenges. *Pathogens* **2025**, *14*, 496. DOI:10.3390/pathogens14050496
 260. Castro A, Komora N, Ferreira V, Mota M, Silva J, Teixeira P. Prevalence of *Staphylococcus aureus* from nares and hands on health care professionals in a Portuguese Hospital. *J. Appl. Microbiol.* **2016**, *121*, 831–839. DOI:10.1111/jam.13186
 261. Mekhloufi OA, Chieffi D, Hammoudi A, Bensefia SA, Fanelli F, Fusco V. Prevalence, Enterotoxigenic Potential and Antimicrobial Resistance of *Staphylococcus aureus* and Methicillin-Resistant *Staphylococcus aureus* (MRSA) Isolated from Algerian Ready to Eat Foods. *Toxins* **2021**, *13*, 835. DOI:10.3390/toxins13120835
 262. Elsalkh SA, Zakaria AI, Abd-Elghany SM, Imre K, Morar A, Sallam KI. Prevalence and Characterization of the Antimicrobial Resistance and Virulence Profiles of *Staphylococcus aureus* in Ready-to-Eat (Meat, Chicken, and Tuna) Pizzas in Mansoura City, Egypt. *Antibiotics* **2025**, *14*, 817. DOI:10.3390/antibiotics14080817
 263. Ohlsen K, Ziebuhr W, Koller KP, Hell W, Wichelhaus TA, Hacker J. Effects of subinhibitory concentrations of antibiotics on alpha-toxin (*hla*) gene expression of methicillin-sensitive and methicillin-resistant *Staphylococcus aureus* isolates. *Antimicrob. Agents Chemother.* **1998**, *42*, 2817–2823. DOI:10.1128/AAC.42.11.2817
 264. Pereira GDN, Rosa RDS, Dias AA, Gonçalves DJS, Seribelli AA, Pinheiro-Hubinger L, et al. Characterization of the virulence, *agr* typing and antimicrobial resistance profile of *Staphylococcus aureus* strains isolated from food handlers in Brazil. *Braz. J. Infect. Dis.* **2022**, *26*, 102698. DOI:10.1016/j.bjid.2022.102698
 265. Uehara Y. Current Status of Staphylococcal Cassette Chromosome *mec* (SCC*mec*). *Antibiotics* **2022**, *11*, 86. DOI:10.3390/antibiotics11010086
 266. Achek R, Hotzel H, Cantekin Z, Nabi I, Hamdi TM, Neubauer H, et al. Emerging of antimicrobial resistance in staphylococci isolated from clinical and food samples in Algeria. *BMC Res. Notes* **2018**, *11*, 663. DOI:10.1186/s13104-018-3762-2
 267. Anas M, Lone SA, Malik A, Ahmad J. Antimicrobial Resistance and Public Health Risks Associated with Staphylococci Isolated from Raw and Processed Meat Products. *Foodborne Pathog. Dis.* **2025**, *22*, 39–50. DOI:10.1089/fpd.2024.0081
 268. Urimi MR, Ansari WK, Islam MS, Sobur MA, Rahman M, Rahman MT. Antibiotic resistance patterns of *Staphylococcus* spp. isolated from fast foods sold in different restaurants of Mymensingh, Bangladesh. *J. Adv. Vet. Anim. Res.* **2021**, *8*, 274–281. DOI:10.5455/javar.2021.h512
 269. Argudín MA, Mendoza MC, González-Hevia MA, Bances M, Guerra B, Rodicio MR. Genotypes, Exotoxin Gene Content, and Antimicrobial Resistance of *Staphylococcus aureus* Strains Recovered from Foods and Food Handlers. *Appl. Environ. Microbiol.* **2012**, *78*, 2930–2935. DOI:10.1128/AEM.07487-11
 270. Taddese D, Abdurahaman M, Debelo M, Shumi E, Urgessa G, Kefyalew D, et al. Antimicrobial resistance characterization of *Staphylococcus aureus* from different animal food origins in Jimma, South Western Ethiopia. *Discov. Food* **2025**, *5*, 27. DOI:10.1007/s44187-025-00300-1
 271. Seow WL, Mahyudin NA, Amin-Nordin S, Radu S, Abdul-Mutalib NA. Antimicrobial resistance of *Staphylococcus aureus* among cooked food and food handlers associated with their occupational information in Klang Valley, Malaysia. *Food Control* **2021**, *124*, 107872. DOI:10.1016/j.foodcont.2021.107872
 272. Phan A, Mijar S, Harvey C, Biswas D. *Staphylococcus aureus* in Foodborne Diseases and Alternative Intervention Strategies to Overcome Antibiotic Resistance by Using Natural Antimicrobials. *Microorganisms* **2025**, *13*, 1732. DOI:10.3390/microorganisms13081732
 273. Arafa AA, Ibrahim ES, Fourad EA, Garber ES. Antibiotic Resistance of Staphylococci Concerning Strains Included in Food Industry in Egypt. *Int. J. Pharm. Clin. Res.* **2016**, *8*, 1583–1589. Available online: <https://impactfactor.org/PDF/IJPCR/8/IJPCR,Vol8,Issue12,Article6.pdf> (accessed on 10 August 2026).
 274. Kayili E, Sanlibaba P. Prevalence, characterization and antibiotic resistance of *Staphylococcus aureus* isolated from traditional cheeses in Turkey. *Int. J. Food Prop.* **2020**, *23*, 1441–1451. DOI:10.1080/10942912.2020.1814323
 275. Touimi GB, Bennani L, Berrada S, Mousa B, Bennani B. Prevalence and antibiotic resistance profiles of *Staphylococcus* sp. isolated from food, food contact surfaces and food handlers in a Moroccan hospital kitchen. *Lett. Appl. Microbiol.* **2020**, *70*, 241–251. DOI:10.1111/lam.13278
 276. Moges M, Rodland EK, Legesse T, Argaw A. Antibiotic resistance patterns of *Staphylococcus aureus* and Enterobacteriaceae isolated from street foods in selected towns of Ethiopia. *BMC Infect. Dis.* **2024**, *24*, 267. DOI:10.1186/s12879-024-09266-4

277. Byczkowska-Rostkowska Z, Gajewska J, Chajęcka-Wierzchowska W. Whole genome analysis and antimicrobial resistance assessment of *Staphylococcus epidermidis* isolated from food sources. *Sci. Total Environ.* **2025**, 993, 179999. DOI:10.1016/j.scitotenv.2025.179999
278. Letuka P, Nkhebenyane SJ, Ramatla T, Zikhona T-N, Lekota KE, Khasapane NG. Antimicrobial resistance and genomic characterization of *Staphylococcus aureus* in ready-to-eat foods from Mangaung Metro Municipality. *Front. Microbiol.* **2025**, 16, 1669035. DOI:10.3389/fmicb.2025.1669035
279. Babines-Orozco L, Balbuena-Alonso MG, Barrios-Villa E, Lozano-Zarain P, Martínez-Laguna Y, Rocha-Gracia RD, et al. Antimicrobial resistance in food-associated *Escherichia coli* in Mexico and Latin America. *Biosci. Microbiota Food Health* **2024**, 43, 4–12. DOI:10.12938/bmfh.2023-022
280. Silva A, Silva V, Pereira JE, Maltez L, Igrejas G, Valentão P, et al. Antimicrobial Resistance and Clonal Lineages of *Escherichia coli* from Food-Producing Animals. *Antibiotics* **2023**, 12, 1061. DOI:10.3390/antibiotics12061061
281. Naidoo N, Zishiri OT. Presence, Pathogenicity, Antibiotic Resistance, and Virulence Factors of *Escherichia coli*: A Review. *Bacteria* **2025**, 4, 16. DOI:10.3390/bacteria4010016
282. Mora-Coto D, Moreno-Vélez P, Luna-Muñoz J, Moreno-Campuzano S, Ontiveros-Torres MA. Intestinal and Extraintestinal Pathotypes of *Escherichia coli* Are Prevalent in Food Prepared and Marketed on the Streets from the Central Zone of Mexico and Exhibit a Differential Phenotype of Resistance Against Antibiotics. *Antibiotics* **2025**, 14, 406. DOI:10.3390/antibiotics14040406
283. Bakshi P, Bhowmik A, Ahsan S, Alim SR. Identification of antibiotic-resistant pathogens and virulence genes in *Escherichia coli* isolates from food samples in the Dhaka University campus of Bangladesh. *Food Sci. Nutr.* **2024**, 12, 1995–2002. DOI:10.1002/fsn3.3896
284. Li Y, Zhang M, Luo J, Chen J, Wang Q, Lu S, et al. Antimicrobial resistance of *Escherichia coli* isolated from retail foods in northern Xinjiang, China. *Food Sci. Nutr.* **2020**, 8, 2035–2051. DOI:10.1002/fsn3.1491
285. Ronald C, Matofari JW, Nduko JM. Antimicrobial resistance of *E. coli* strains in ready-to-eat red meat products in Nakuru County, Kenya. *Microbe* **2023**, 1, 100022. DOI:10.1016/j.microb.2023.100022
286. Liu C, Sun S, Sun Y, Li X, Gu W, Luo Y, et al. Antibiotic resistance of *Escherichia coli* isolated from food and clinical environment in China from 2001 to 2020. *Sci. Total Environ.* **2024**, 939, 173498. DOI:10.1016/j.scitotenv.2024.173498
287. Ramatla T, Jane N, Dineo M, Mpho T, Tshagofatso M, Khasapane NG. The Global Prevalence of Antibiotic Resistance and Shiga Toxin-Producing *Escherichia coli* in Chickens: A Systematic Review and Meta-Analysis (2011–2024). *Antibiotics* **2025**, 14, 568. DOI:10.3390/antibiotics14060568
288. Hinson C, Sessou P, Azokpota P, Daube G, Korsak N. Antibiotic Resistance of *Escherichia coli* and *Salmonella* Isolates from Table Eggs, Poultry Sausages, and Clinical Samples in Southwest Benin. *Hygiene* **2026**, 6, 9. DOI:10.3390/hygiene6010009
289. Gruel G, Sellin A, Riveiro H, Pot M, Breurec S, Guyomard-Rabenirina S, et al. Antimicrobial use and resistance in *Escherichia coli* from healthy food-producing animals in Guadeloupe. *BMC Vet. Res.* **2021**, 17, 116. DOI:10.1186/s12917-021-02810-3
290. Alexandre LA, da Silva AC, Nascimento FLDA, de Melo APZ, Verruck S. Antimicrobial resistance of *Escherichia coli* isolated from fish and aquaculture water: An emerging concern for consumers. *Lett. Appl. Microbiol.* **2025**, 78, ovaf099. DOI:10.1093/lambio/ovaf099
291. van Hamerlsveld S, Kurenbach B, Paull DJ, Godsoe WA, Ferguson GC, Heinemann JA. Indigenous food sources as vectors of *Escherichia coli* and antibiotic resistance. *Environ. Pollut.* **2023**, 334, 122155. DOI:10.1016/j.envpol.2023.122155
292. Osman KM, Kappell AD, Elhadidy M, ElMougy F, Abd El-Ghany WA, Orabi A, et al. Poultry hatcheries as potential reservoirs for antimicrobial resistant *Escherichia coli*: A risk to public health and food safety. *Sci. Rep.* **2018**, 8, 5859. DOI:10.1038/s41598-018-23962-7
293. Nesse LL, Osland AM, Asal B, Mo SS. Evolution of antimicrobial resistance in *E. coli* biofilm treated with high doses of ciprofloxacin. *Front. Microbiol.* **2023**, 14, 1246895. DOI:10.3389/fmicb.2023.1246895
294. Muloi D, Ward MJ, Pedersen AB, Fèvre EM, Woolhouse MEJ, van Bunnik BAD. Are Food Animals Responsible for Transfer of Antimicrobial-Resistant *Escherichia coli* or Their Resistance Determinants to Human Populations? A Systematic Review. *Foodborne Pathog. Dis.* **2018**, 15, 467–474. DOI:10.1089/fpd.2017.2411
295. Baloch AB, Yang H, Feng Y, Xi M, Wu Q, Yang Q, et al. Presence and Antimicrobial Resistance of *Escherichia coli* in Ready-to-Eat Foods in Shaanxi, China. *J. Food Prot.* **2017**, 80, 420–424. DOI:10.4315/0362-028X.JFP-16-175
296. Davis GS, Waits K, Nordstrom L, Grande H, Weaver B, Papp K, et al. Antibiotic-resistant *Escherichia coli* from retail poultry meat with different antibiotic use claims. *BMC Microbiol.* **2018**, 18, 174. DOI:10.1186/s12866-018-1322-5
297. Eltai NO, Yassine HM, Al Thani AA, Madi MAA, Ismail A, Ibrahim E, et al. Prevalence of antibiotic resistant *Escherichia coli* isolates from fecal samples of food handlers in Qatar. *Antimicrob. Resist. Infect. Control.* **2018**, 7, 78.

DOI:10.1186/s13756-018-0369-2

298. Rezaei N, Rezaei A, Shahmoradi S, Bashtin A. Evaluation of Antibiotic Resistance in *Escherichia coli* Strains Isolated from Pastry Cream in Hamadan, Iran. *Int. J. Health Life Sci.* **2019**, *5*, e87103. DOI:10.5812/ijhls.87103
299. Majewski M, Józefiak A, Kimsa-Furdzik M, Dziubdziela L, Hudak-Nowak M, Wilczyński J, et al. Antimicrobial resistance of *Escherichia coli* and *Klebsiella* spp. conventionally sampled from factory-farmed chickens—clinical submissions. *Ann. Agric. Environ. Med.* **2021**, *28*, 271–276. DOI:10.26444/aaem/120927
300. Abdelwahab GE, Ishag HZA, Al Hammadi ZM, Al Yammahi SMS, Mohd Yusof MFB, Al Yassi MSY, et al. Antibiotics Resistance in *Escherichia coli* Isolated from Livestock in the Emirate of Abu Dhabi, UAE, 2014–2019. *Int. J. Microbiol.* **2022**, *2022*, 3411560. DOI:10.1155/2022/3411560
301. Abebe E, Gugsu G, Ahmed M, Awol N, Tefera Y, Abegaz S, et al. Occurrence and antimicrobial resistance pattern of *E. coli* O157:H7 isolated from foods of Bovine origin in Dessie and Kombolcha towns, Ethiopia. *PLoS Negl. Trop. Dis.* **2023**, *17*, e0010706. DOI:10.1371/journal.pntd.0010706
302. Mansaray AHD, Yankson DPY, Johnson RAB, Moses FL, Kanu JS, Kamara IF, et al. Bacterial Isolates and Antibiotic Resistance of *Escherichia coli* Isolated from Fresh Poultry Excreta Used for Vegetable Farming in Freetown, Sierra Leone. *Int. J. Environ. Res. Public Health* **2022**, *19*, 5405. DOI:10.3390/ijerph19095405
303. Runa N, Yesmin S, Husna A, Runa N, Islam M, Lovelu M, et al. Prevalence of multidrug-resistant ESBL-producing *Escherichia coli* isolated from beef and sheep meat in Sylhet, Bangladesh. *J. Adv. Biotechnol. Exp. Ther.* **2024**, *7*, 520–529. DOI:10.5455/jabet.2024.d45
304. Iribagiza A, Gwimile ES, Nganga PM, Nkundwanayo C, Ntakirutimana D, Ntirandekura JB, et al. Qualitative analysis of antibiotic residue and antibiotic resistance testing of *E. coli* isolated from traditionally heat-treated fresh milk sold in local restaurants of Bujumbura City, Burundi. *ISAR J. Agric. Biol.* **2024**, *2*, 5–9. DOI:10.5281/zenodo.14533215
305. Amare AD, Kurra FA, Gari FR. Prevalence and Antimicrobial Susceptibility Profile of *E. coli* O157:H7 in Fish Collected From Lake Ziway, Ethiopia. *BioMed. Res. Int* **2026**, *2026*, 5097078. DOI:10.1155/bmri/5097078
306. Guimarães Carvalho G, Calarga AP, Zorgi NE, Astudillo-Trujillo CA, Pardini Gontijo MT, Brocchi M, et al. Virulence and DNA sequence analysis of *Cronobacter* spp. isolated from infant cereals. *Int. J. Food Microbiol.* **2022**, *376*, 109745. DOI:10.1016/j.ijfoodmicro.2022.109745
307. Jang H, Gopinath GR, Eshwar A, Srikumar S, Nguyen S, Gangiredla J, et al. The Secretion of Toxins and Other Exoproteins of *Cronobacter*: Role in Virulence, Adaption, and Persistence. *Microorganisms* **2020**, *8*, 229. DOI:10.3390/microorganisms8020229
308. Sani NA, Odeyemi OA. Occurrence and prevalence of *Cronobacter* spp. in plant and animal derived food sources: A systematic review and meta-analysis. *SpringerPlus* **2015**, *4*, 545. DOI:10.1186/s40064-015-1324-9
309. Kumar A, Vaiyapuri M, Mothadaka MP. Antimicrobial Resistance in *Cronobacter sakazakii*. In *Handbook on Antimicrobial Resistance*, 1st ed.; Mothadaka MP, Vaiyapuri M, Rao Badireddy M, Nagarajrao Ravishankar C, Bhatia R, Jena J, Eds.; Springer: Singapore, 2023; pp. 541–559. DOI:10.1007/978-981-19-9279-7_26
310. Fei P, Jiang Y, Feng J, Forsythe SJ, Li R, Zhou Y, et al. Antibiotic and Desiccation Resistance of *Cronobacter sakazakii* and *C. malonaticus* Isolates from Powdered Infant Formula and Processing Environments. *Front. Microbiol.* **2017**, *8*, 316. DOI:10.3389/fmicb.2017.00316
311. Csorba C, Pajić M, Blagojević B, Forsythe S, Radinović M, Velebit B. Prevalence, characterization, and antibiotic susceptibility of *Cronobacter* spp. in a milk powder processing environment: The first reported case in Serbia. *Food Sci. Nutr.* **2022**, *10*, 2254–2563. DOI:10.1002/fsn3.2681
312. Fakruddin M, Rahaman M, Ahmed MM, Hoque MM. Stress tolerant virulent strains of *Cronobacter sakazakii* from food. *Biol. Res.* **2014**, *47*, 63. DOI:10.1186/0717-6287-47-63
313. Wang L, Forsythe SJ, Yang X, Fu S, Man C, Jiang Y. Invited review: Stress resistance of *Cronobacter* spp. affecting control of its growth during food production. *J. Dairy Sci.* **2021**, *104*, 11348–11367. DOI:10.3168/jds.2021-20591
314. Jaradat ZW, Al-Mousa WA, Elbetieha AM, Ababneh QO, Al-Nabulsi AA, Jang H, et al. Virulence, antimicrobial susceptibility and phylogenetic analysis of *Cronobacter sakazakii* isolates of food origins from Jordan. *J. Appl. Microbiol.* **2022**, *133*, 2528–2546. DOI:10.1111/jam.15723
315. Pakbin B, Brück WM, Allahyari S, Rossen JWA, Mahmoudi R. Antibiotic Resistance and Molecular Characterization of *Cronobacter sakazakii* Strains Isolated from Powdered Infant Formula Milk. *Foods* **2022**, *11*, 1093. DOI:10.3390/foods11081093
316. Pakbin B, Mahmoudi R, Mousavi S, Allahyari S, Amani Z, Peymani A, et al. Genotypic and antimicrobial resistance characterizations of *Cronobacter sakazakii* isolated from powdered milk infant formula: A comparison between domestic and imported products. *Food Sci. Nutr.* **2020**, *8*, 6708–6717. DOI:10.1002/fsn3.1965
317. Abbasi Bafetrat S, Goli M, Momtaz H. The Study on antibiotic resistance pattern of *Cronobacter sakazakii* strains isolated

- from infant formula and baby food types. *J. Food Microbiol.* **2017**, *3*, 39–50. Available online: <https://www.sid.ir/paper/250763/en> (accessed on 10 August 2026).
318. Gan X, Li M, Xu J, Yan S, Wang W, Li F. Emerging of Multidrug-Resistant *Cronobacter sakazakii* Isolated from Infant Supplementary Food in China. *Microbiol. Spectr.* **2022**, *10*, e01197–22. DOI:10.1128/spectrum.01197-22
 319. Abebe GM. *Cronobacter sakazakii* in Infant Food Contamination and Its Survival Strategies in Hostile Conditions. *Int. J. Pediatr. Res.* **2020**, *6*, 067. DOI:10.23937/2469-5769/1510067
 320. Dostert M, Trimble MJ, Hancock REW. Antibiofilm peptides: Overcoming biofilm-related treatment failure. *RSC Adv.* **2021**, *11*, 2718–2728. DOI:10.1039/D0RA09739J
 321. Holý O, Parra-Flores J, Bzdil J, Cabal-Rosel A, Diaz-Prieto B, Cruz-Córdoba A, et al. Screening of Antibiotic and Virulence Genes from Whole Genome Sequenced *k* Isolated from Food and Milk-Producing Environments. *Antibiotics* **2023**, *12*, 851. DOI:10.3390/antibiotics12050851
 322. Lee IPA, Andam CP. Pan-genome diversification and recombination in *Cronobacter sakazakii*, an opportunistic pathogen in neonates, and insights to its xerotolerant lifestyle. *BMC Microbiol.* **2019**, *19*, 306. DOI:10.1186/s12866-019-1664-7
 323. Guimarães Carvalho G, Parolin Calarga A, Teodoro JR, Mariz Queiroz M, Astudillo-Trujillo CA, Levy CE, et al. Isolation, comparison of identification methods and antibiotic resistance of *Cronobacter* spp. in infant foods. *Food Res. Int.* **2020**, *137*, 109643. DOI:10.1016/j.foodres.2020.109643
 324. Parra-Flores J, Aguirre J, Junja V, Jackson EE, Cruz-Córdoba A, Silva-Sánchez J, et al. Virulence and Antibiotic Resistance Profiles of *Cronobacter sakazakii* and *Enterobacter* spp. Involved in the Diarrheic Hemorrhagic Outbreak in Mexico. *Front. Microbiol.* **2018**, *9*, 2206. DOI:10.3389/fmicb.2018.02206
 325. Fei P, Xing M, Feng Y, Liu S, Chang Y, Wang Y, et al. Occurrence, Molecular Characterization, and Antibiotic Resistance of *Cronobacter sakazakii* in Goat Milk-Based Infant Formula from Shaanxi Province, China. *Foodborne Path. Dis.* **2022**, *19*, 304–310. DOI:10.1089/fpd.2021.0095
 326. Mardaneh J, Dallal MMS. Study of *Cronobacter sakazakii* Strains Isolated from Powdered Milk Infant Formula by Phenotypic and Molecular Methods in Iran. *Arch. Pediatr. Infect. Dis.* **2017**, *5*, e38867. DOI:10.5812/pedinf.38867
 327. Song D, Jia A, Liu B, Liu S, Dong K, Man C, et al. Whole-transcriptome analysis after the acquisition of antibiotic resistance of *Cronobacter sakazakii*: Mechanisms of antibiotic resistance and virulence changes. *Food Res. Int.* **2023**, *174*, 113664. DOI:10.1016/j.foodres.2023.113664
 328. Garbowska M, Berthold-Pluta A, Stasiak-Różańska L, Pluta A, Fortsythe S, Stefańska I. The Genotyping Diversity and Hemolytic Activity of *Cronobacter* spp. Isolated from Plant-Based Food Products in Poland. *Foods* **2023**, *12*, 3873. DOI:10.3390/foods12203873
 329. Oloninefa SD, Daniyan SY, Abalaka ME. Antibiotic Susceptibility Profile of *Cronobacter sakazakii*. *Anchor Univ. J. Sci. Technol. (AUJST)* **2020**, *1*, 10–15. Available online: <https://www.ajol.info/index.php/aujst/article/view/222124/209605> (accessed on 10 August 2026)
 330. Capita R, Vicente-Velasco M, Rodríguez-Melcón C, García-Fernández C, Carballo J, Alonso-Calleja C. Effect of low doses of biocides on the antimicrobial resistance and the biofilms of *Cronobacter sakazakii* and *Yersinia enterocolitica*. *Sci. Rep.* **2019**, *9*, 15905. DOI:10.1038/s41598-019-51907-1
 331. Wirtanen G, Salo S. Disinfection in food processing—Efficacy testing of disinfectants. *Rev. Environ. Sci. Bio/Technol.* **2003**, *2*, 293–306. DOI:10.1023/B:RESB.0000040471.15700.03
 332. Aryal M, Muriana PM. Efficacy of Commercial Sanitizers Used in Food Processing Facilities for Inactivation of *Listeria monocytogenes* E. coli O157:H7, and *Salmonella* Biofilms. *Foods* **2019**, *8*, 639. DOI:10.3390/foods8120639
 333. Song D, Qi X, Huang Y, Jia A, Liang Y, Man C, et al. Comparative proteomics reveals the antibiotic resistance and virulence of *Cronobacter* isolated from powdered infant formula and its processing environment. *Int. J. Food. Microbiol.* **2023**, *407*, 110374. DOI:10.1016/j.ijfoodmicro.2023.110374
 334. Cui JH, Yu B, Xiang Y, Zhang Z, Zhang T, Zeng YC, et al. Two Cases of Multi-antibiotic Resistant *Cronobacter* spp. Infections of Infants in China. *Biomed. Environ. Sci.* **2017**, *30*, 601–605. DOI:10.3967/bes2017.079
 335. Shi L, Liang Q, Zhan Z, Feng J, Zhao Y, Chen Y, et al. Co-occurrence of 3 different resistance plasmids in a multi-drug resistant *Cronobacter sakazakii* isolate causing neonatal infections. *Virulence* **2018**, *9*, 110–120. DOI:10.1080/21505594.2017.1356537
 336. Zeng H, Lei T, He W, Zhang J, Liang B, Li C, et al. Novel Multidrug-Resistant *Cronobacter sakazakii* Causing Meningitis in Neonate, China, 2015. *Emerg. Infect. Dis.* **2018**, *24*, 2121–2123. DOI:10.3201/eid2411.180718
 337. Ye Y, Wu Q, Xu X, Yang X, Dong X, Zhang J. The phenotypic and genotypic characterization of *Enterobacter sakazakii* strains from infant formula milk. *J. Dairy Sci.* **2010**, *93*, 2315–2320. DOI:10.3168/jds.2009-2662
 338. Kilonzo-Nthenge A, Rotich E, Godwin S, Nahashon S, Chen F. Prevalence and Antimicrobial Resistance of *Cronobacter sakazakii* Isolated from Domestic Kitchens in Middle Tennessee, United States. *J. Food Prot.* **2012**, *75*, 1512–1517.

DOI:10.4315/0362-028X.JFP-11-442

339. Abdel-Galil FY, Abdel-Latif HK, Ammar AM, Serry FME. Studies on prevalence, antimicrobial resistance and survival of *Cronobacter sakazakii*. *Zagazig J. Pharm. Sci.* **2014**, *23*, 1–15. DOI:10.21608/zjps.2014.38173
340. Carvalho GG, Calarga AP, Brocchi M, Kabuki DY. Antibiotic Resistance Of *Cronobacter sakazakii* Isolated From Powdered Infant Cereals. *Anais do 13º Simpósio Latino Americano de Ciência de Alimentos. O Futuro dos Alimentos. 2019, Campinas. Anais eletrônicos...* Campinas, Galoá, 2019. Available online: <https://proceedings.science/slaca/slaca-2019/trabalhos/antibiotic-resistance-of-cronobacter-sakazakii-isolated-from-powdered-infant-cer?lang=en>. (accessed on 19 February 2024).
341. Tayeb BA, Mohamed Sharif YH, Ameren AM. Incidence rate and antibiotic resistance profile of *Cronobacter sakazakii* isolated from various food products. *Food Res.* **2020**, *4*, 2217–2223. DOI:10.26656/fr.2017.4(6).304
342. Li H, Fu S, Song D, Qin X, Zhang W, Man C, et al. Identification, Typing and Drug Resistance of *Cronobacter* spp. in Powdered Infant Formula and Processing Environment. *Foods* **2023**, *12*, 1084. DOI:10.3390/foods12051084
343. Taji Z, Bonyadian M, Moshtaghi H. Contamination and antibiotic resistance profile of *Cronobacter sakazakii* isolated from raw milk and infant formula. *Iran J. Microbiol.* **2026**, *18*, 279–284. DOI:10.18502/ijm.v18i2.21305
344. Carruthers LV. Trends in Antimicrobial Resistant Non-Typhoidal *Salmonella* Before and After the Food and Drug Administration’s Guidance for Industry #213, United States 2013–2020. *Public Health Theses.* 2022. Available online: <https://elischolar.library.yale.edu/ysphtdl/2138> (accessed on 10 August 2026).
345. Zhao C, Wang Y, Mulchandani R, Van Boeckel TP. Global surveillance of antimicrobial resistance in food animals using priority drugs maps. *Nat. Commun.* **2024**, *15*, 76. DOI:10.1038/s41467-024-45111-7