Check for updates

RESEARCH ARTICLE OPEN ACCESS

Time-of-Day Defines the Risk of Thermally Abused Frying Oil to Renal Injury by Modulating the Diurnal Dynamics of Oxylipins

Yanjun Liu[1,2,7,8] | Meishan Lu[3] | Yanpeng Du[3] | Jie Xu[2,7] | Mingjie Fan[4,5] | Changhu Xue[2,6,7,8] | Yuanfa Liu[1,3]

[1]School of Food and Health, Key Laboratory of Geriatric Nutrition and Health, Beijing Technology and Business University, Beijing, China | [2]Laboratory of Food Science and Human Health, State Key Laboratory of Marine Food Processing & Safety Control, College of Food Science and Engineering, Ocean University of China, Qingdao, China | [3]State Key Laboratory of Food Science and Technology, School of Food Science and Technology, Jiangnan University, Wuxi, China | [4]Department of Diabetes Complications and Metabolism, Arthur Riggs Diabetes & Metabolism Research Institute, Beckman Research Institute, City of Hope National Medical Center, Duarte, California, USA | [5]Shandong First Medical University & Shandong Academy of Medical Sciences, Jinan, China | [6]Qingdao Marine Science and Technology Center, Qingdao, China | [7]Sanya Ocean Institute, Ocean University of China, Sanya, China | [8]Qingdao Institute of Marine Bioresources For Nutrition & Health Innovation, Qingdao, China

Correspondence: Yanjun Liu (fs-lauyang@hotmail.com) | Yuanfa Liu (yfliu@jiangnan.edu.cn)

Received: 3 April 2024 | **Revised:** 26 March 2025 | **Accepted:** 26 April 2025

Keywords: circadian clock | epoxide hydrolases | renal injury

ABSTRACT

Modulation of circadian rhythms impairs homeostasis, resulting in altered susceptibility to disease development. New perspectives on nutrition emphasize the circadian timing of food intake. Here, we found that the ad libitum feeding of oxidized frying oil disrupted the conversion of epoxides to diols in the kidneys, leading to aggravated renal injury. Interestingly, the circadian rhythms of epoxides and their diols in the kidneys and plasma showed differences between the light and dark phases. We demonstrated that time-imposed feeding of fried oil during the active period resulted in limited damage to renal function, conversely, renal function was impaired during the inactive period. Remarkably, feeding with epoxy stearic acid (EpSA) from fried oil during the inactive period disrupted the rhythmicity of epoxide hydrolases and related metabolites, and fueled the progression of renal fibrosis and injury. The effect of EpSA on SMA and E-cadherin, two specific biomarkers for renal inflammation and fibrosis, are lost in knockouts of Ephx1 and Ephx2, which encodes for epoxide hydrolases. In addition, dietary supplementation with linoleic acid, which inhibits epoxide hydrolases, protected mice from damage caused by time-imposed feeding of EpSA. Our findings revealed that the time-of-day dependence of epoxide hydrolases, and circadian disruption induced by time-imposed feeding of EpSA from thermo-induced oxidized oil have adverse effects on kidney function.

1 | Introduction

Staying up late and eating late at night are more common among young people [1]. Modern lifestyles contribute to staying up late and late-night eating because of several physiological and non-physiological factors, such as increased energy intake during the day, low food quality, shift work, the presence of night eating syndrome, religious practices, disrupted circadian rhythms, or special physiological state, including the third trimester of pregnancy [2–5]. Notably, a growing number of young Chinese

Yanjun Liu and Meishan Lu contributed equally to this study.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2026 The Author(s). *Exploration* published by Henan University and John Wiley & Sons Australia, Ltd.

individuals are abandoning the established norm of three meals a day and making late-night meals a staple of their lives. Many obsess over skewers, hotpots, and cooked seafood, eating late almost every weekend. Late-night eating has been associated with several metabolic dysfunction-related diseases, such as anxiety or depressive disorder, obesity, diabetes, and hypertension [1, 5]. Circadian disruption associated with late-night eating is believed to be partly responsible for the higher incidence of chronic diseases among late-night eaters.

It is noteworthy that the types of foods and beverages consumed late at night are critical [2, 6]. People who eat late at night often tend to follow less healthy diets. Huang et al., [7] and Noel et al. [8] found that dietary choices among late-night eaters include high-sodium foods, fried food, high saturated fat, low fruit and vegetable etc., Since food choices consistently focused on taste and convenience. Similarly, the dietary patterns of rotating shift workers include increased snacking or eating at night and the consumption of fewer nutrient-dense foods [9]. The proportion of energy derived from saturated fat and fried foods tends to be high among shift workers [10]. Thus, the adoption of unhealthy dietary patterns may exacerbates various unhealthy adverse health outcomes. Demand for convenient and quick meals has increased across China. Grilled or fried foods are more frequently consumed as late-night meals, driven by the "lazy economy" and the rise of convenient delivery services. These foods' nutritional characteristics may negatively affect the metabolic and circadian rhythms that are required for optimal health [2]. Recently, the harmful effects of dietary oxidized frying oil have recently gradually been realized. Although cooking oils are frequently used for frying, repeated heating of cooking oils is both genotoxic and mutagenic, with potential tumorigenic effects. Remarkably, during repeated heating, deep-frying oil undergoes various chemical reactions, including hydrolysis, polymerization, lipid oxidation, and the Maillard reaction, producing oxidized molecules that may increase the risk of cancer, anxiety, depressive disorders, and cardiovascular disease [11]. Poor diet quality (such as fried oil) among late eaters may contribute to these associations. Currently, there are no studies examining the specific dietary patterns of individuals consuming heated cooking oils and their disease risk; therefore, we investigated the risk and synergistic effects of heated cooking oils and unhealthy dietary patterns in this research.

Studies on late-night eating have primarily focused on obesity, inflammation, and other metabolic health issues, with limited research on its potential impacts on renal function [12]. In the National Health and Nutrition Examination Survey (2013–2020), a strong correlation was found between meal timing and the incidence of chronic kidney disease, suggesting that eating late may worsen renal outcomes for patients with diabetes [12]. Sleep disorders are prevalent in patients with chronic kidney disease, raising the possibility that these patients may have a disorganized circadian timing system [13]. Thus, alterations in peripheral clock rhythms resulting from late-night eating may contribute to an increased risk of chronic kidney disease. Chronic kidney disease incidence is increasing globally and has become a major health problem [14]. Several renal functions, including renal plasma flow, glomerular filtration rate, tubular transport activities and diuresis, exhibit circadian rhythms [14]. Numerous physiological functions exhibit substantial circadian oscillations. In the kidneys, renal plasma flow, the glomerular filtration rate and tubular reabsorption and/or secretion processes have been shown to peak during the active phase and decline during the inactive phase [14]. In this study, we investigated the timing of fried oil intake was investigated as a novel factor in the etiology and progression of kidney injury.

2 | Results

2.1 | Dietary Oxidized Oil Elicits Time-Dependent Renal Injury

We investigated the effects of thermally induced oxidized oil intake and feeding time on renal injury in mice. Although fried oil had no effect on kidney injury in chow diet-treated mice (Figure S1A,B), histological examination of mice treated with oxidized oil revealed severe renal injury after ad libitum feeding of a high-fat diet containing fried oil (FO, thermo-induced oxidized oil), as indicated by remarkable tubular injury and swelling (Figure S2A,B). The oxidized oil also increased the levels of blood urea nitrogen (BUN, Figure S2C) and creatinine (Figure S2D), without affecting body weight or food intake. In addition, Masson's trichrome staining showed a slight increase in fibrotic areas in the kidneys of oxidized oil-treated mice (Figure S2E), indicating that renal dysfunction and fibrosis induced by dietary thermally induced oxidized oil. To assess renal function in mice with a timed intake of oxidized oil (Figure S2A and Figure 1A), we tested two different timed feeding pattern protocols (light-restricted feeding and 12-h active/12-h inactive feeding). Time-restricted feeding of oxidized oil during the inactive period promoted kidney injury in mice, as indicated by vacuolar degeneration, glomerular hypertrophy, and tubular lumen expansion in renal tissue (Figure S2B) and elevated serum BUN and creatinine levels (Figures S2C,D). Likewise, as for the mice with access to HFD containing oxidized oil (Figure 1A) during light-phase (LFO) and HFD during dark-phase (DFO), these mice show exacerbated pathological renal damage and glomerular fibrosis in mice (Figures 1B–D), with significantly increased BUN, creatinine, and urine albumin-creatinine ratio (UACR) (Figures 1E,F and Figure S3A), confirming that timed feeding of oxidized oil during the daytime leads to aggravated renal dysfunction. By contrast, micefed an HFD containing oxidized oil during the dark phase (DFO) did not show significant changes in renal dysfunction. Consistent with these pathological changes, timed feeding of oxidized oil during the light/inactive period increased renal α-smooth muscle actin (α-SMA) and E-cadherin (Figures 1G–I and Figure S3A,B) and serum KIM-1 (kidney injury molecule 1) and NGAL (neutrophil gelatinase-associated lipocalin) levels (Figure S4B,C), indicating marked activation of the fibrogenic pathway during timed feeding of oxidized oil in the inactive period.

2.2 | Altered Oxylipin Metabolism and Dysregulation of Epoxide Hydrolase in Mice With Timed Feeding of Oxidized Oil

We compared the ultra-high-performance liquid chromatography-mass spectrometry (UPLC-MS/MS) oxylipin

27662098, 2026, 1, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/EXP.20240073 by CochraneChina, Wiley Online Library on [08/06/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License




FIGURE 1 | Time feeding of fried oil in the inactive period worsened renal injury in HFD-treated mice. (A) Schematic of the experimental protocols indicating time window of food access during a 24 h day under ad libitum feeding of Normal diet or HFD ($n = 8$), LFO (light-feeding fried oil, mice were access to HFD containing fried oil for 12 h from ZT0 to ZT12, $n = 8$), and DFO (dark-feeding fried oil, mice were access to HFD containing fried oil for 12 h from ZT12 to ZT24, $n = 8$). (B–D) Representative images of HE, PAS, Masson staining (B) in mice. Its renal damage score (C) and fibrotic area (D) were measured. Arrows represent renal glomerular damage, slight swelling of renal tubular epithelial cells, or glomerular fibrosis. (E) Serum BUN in mice ($n = 8$). (F) Serum creatinine in mice ($n = 8$). (G–I) Immunofluorescence of α-SMA and E-cadherin in kidney. The fluorescent intensity of the α-SMA (H) and E-cadherin (I) was examined. Two-way ANOVA were used for statistical tests and Holm–Sidak method was used for p value adjustment for multiple comparisons. Data are mean ± SEM. ns, no significant difference, * $P < 0.05$, ** $P < 0.01$.

profile signatures in the kidneys of mice with access to oxidized oil during the light phase (inactive) versus the dark phase (active). Oxygenated polyunsaturated fatty acids (oxylipins) are essential bioactive lipid mediators derived from arachidonic acid (AA), linoleic acid (LA), and other polyunsaturated fatty acids (PUFAs). Twenty-seven oxylipin metabolites were detected and their changes in these mice are shown in a normalized heatmap (Figure 2A). The renal oxylipidome was affected in mice treated with LFO (access to oxidized oil in the light phase) but not in DFO-treated mice (access to oxidized oil in the dark phase). Cytochrome P450 (CYP450) epoxygenases can utilize arachidonic acid (AA) and linoleic acid (LA) as substrates to produce endogenous epoxides or other oxylipins, which are subsequently hydrolyzed by epoxide hydrolases (EHs, including microsomal epoxide hydrolase and soluble epoxide hydrolase) to dihydroxyeicosatrienoic acids (DHETs)

27662098, 2026, 1, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/EXP.20240073 by CochraneChina, Wiley Online Library on [08/06/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License




FIGURE 2 | Timed-imposed oxidized oil during the inactive period contributes to alteration of oxylipins metabolism. (A) Heat map of dietary fried oil effects on oxylipins profile in kidney ($n = 8$). (B) Cytochrome P450 (CYP) epoxygenases and soluble epoxide hydrolase catalyze the epoxidation of PUFA and converting epoxides to their corresponding diols, respectively. (C) Volcano plot of oxylipins profile (HFD versus DFO; fold change, ≥2; q value < 0.05). (D) Volcano plot of oxylipins profile (HFD versus LFO; fold change, ≥2; q value < 0.05). (E) mRNA expression of genes encoding epoxide hydrolase (Ephx1 and Ephx2, $n = 8$). (F) Heat map of gene expression related to cytochrome P450 monooxygenase eicosanoid pathway. (G) Protein levels of Ephx1 and Ephx2 ($n = 4$) in kidney were measured. Data are mean ± SEM. ns, no significant difference, * P <0.05, ** P <0.01.

or DiHOME, the diol form of epoxides (Figure 2B). Notably, we plotted fold changes in oxylipins levels and identified reduced levels of 9,10-epoxyoctadecenoic (EpOME) and several epoxyeicosatrienoic acids (EETs, including 11,12-EET and 14,15-EET) along with increased levels of 9,10-DiHOME and several dihydroxyeicosatrienoic acids (14,15-DHET and 8,9-DHET) in the kidneys of LFO-fed mice (Figure 2C). These changes were only observed between control and LFO-fed mice. The imbalance between the epoxides and their corresponding diols was associated with increased mRNA and protein expression of genes encoding EHs (Ephx1 and Ephx2) in the kidneys of LFO-fed mice, as reflected by Figure 2D,E,G, and Figure S5, which are involved in the conversion of bioactive epoxides to less active diols. In addition, feeding oxidized oil during the light phase failed to modify the cytochrome P450 monooxygenase eicosanoid pathway (*Cyp2c29*, *Cyp2c44*, *Cyp2j2*, *Cyp2j5*, *Cyp2j9*, *Cyp2c8*, *Cyp2c9*, *Cyp4a12a*, *Cyp4a12b*, *Cyp4f16*, and *Cyp4f13*), the predominant epoxygenase isoforms involved in EETs formation, except for *Cyp2c37* (Figure 2F). These results suggested that timed feeding of oxidized oil during the inactive period worsens oxylipins metabolism in the kidney via the activation of epoxide hydrolase.

27662098, 2026, 1, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/EXP.20240073 by CochraneChina, Wiley Online Library on [08/06/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

2.3 | Renal Transcriptional Oscillations of Epoxide Hydrolase Genes Were Disrupted by Epoxy Stearic Acid in Oxidized Oil

Currently, palm oil and vegetable oil with high oleic acid contents are the preferred choices for deep frying and are widely used in commercial restaurants [15, 16]. The safety of fried food is controlled by the chemical structural changes and oxidation of the frying oil [17, 18]. Our research group and others have reported that epoxy-fatty acids (EpFAs) in fried oils, ranging from 0.3 to 5.9%, are the main group of oxidized and polar lipids [19]. During the deep frying of palm oil and high oleic acid oil, we found that epoxy stearic acid (EpSA, 9,10-epoxy stearic acid) derived from oleic acid during thermal oxidation is the main form of EpFAs. The presence of EpFAs in oxidized edible oils is thought to be a hazardous lipid oxidation products [20], and several studies have suggested the potential health risks associated with consuming epoxy fatty acids in the diet [21, 22]. In our study, through in vivo EHs activity assessment, we found that 9,10-EpSA in oxidized edible oils activated EH activity in TCMK-1 cells treated with high glucose (Figure S6A). In addition, 9,10-EpSA increased the protein levels of Ephx1 (gene encodes for microsomal epoxide hydrolase, mEH) and Ephx2 (gene encodes for soluble epoxide hydrolase, sEHs) in TCMK-1 tubular epithelial cells, the tubular epithelial cells (Figures S6B,C), suggesting that 9,10-EpSA in oxidized edible oils is involved in the regulation of oxylipins metabolism. We hypothesized that renal injury induced by timed feeding of oxidized edible oils may be related to the loss of the circadian rhythmicity in oxylipin metabolism.

To understand the molecular basis underlying the loss of circadian rhythmicity with timed feeding of EpSA from oxidized oil, we initiated transcriptomics analysis and assessed the diurnal rhythms of kidneys from mice collected at 4-h intervals over 24 h. Mice were fed with HFD during the dark phase or accessed with HFD containing EpSAs during the light phase (Figure 3A). As expected, renal transcriptome analysis by RNA sequencing and rhythmic analysis using MetaCycle identified a total of 282 dark/inactive and 982 light/active in-phase circadian transcripts. When mice were fed EpSA during the light phase, these 982 light in-phase circadian transcripts were reduced to 533 transcripts, with 55 common to both conditions (Figure 3B). However, the number of dark in-phase circadian transcripts increased slightly to 328 transcripts when fed with EpSA during the light-phase. In addition, 29 transcripts were not classified as day- or night-in-phase circadian transcripts in either group. Disruption of circadian rhythmicity during timed feeding of EpSA was observed for both day- and night-peaking transcripts, with the majority of transcripts showing a loss of rhythmicity (Figure 3B–D) and a significant reduction in circadian amplitude of expression (Figure 3D), as indicated by the JTK_CYCLE results of diurnally oscillating transcripts. Furthermore, we identified the rhythmicity of four genes (including Ephx1, Ephx2, Ephx3, and Ephx4) encoding epoxide hydrolases. Consistent with the findings of previous studies [23, 24], the expression profiles of epoxide hydrolase transcripts oscillated in a circadian manner and appeared to be clearly circadian in control mice (e.g., Ephx1 and Ephx2; Figure 3E). The rhythmic oscillations of Ephx1 and Ephx2 at the protein level in the control group were similar to the circadian rhythmicity of transcripts (Figure 3F). Notably, the phase distribution of Ephx1 and Ephx2 peaked around ZT17 and ZT20, respectively, whereas Ephx1 and Ephx2 were inhibited by the circadian clock during the light/inactive phase. However, we observed that timed feeding of EpSA completely disrupted the phase of epoxide hydrolases-coding genes in the kidney (Figure 3E). In particular, the inhibition of Ephx1 and Ephx2 during the light/inactive phase was disrupted by timed feeding of EpSA in this phase.

2.4 | Loss of Circadian Rhythmicity in Oxylipins by Timed Feeding of Epoxy Stearic Acid

To match our findings on diurnal transcriptomes, we assessed diurnal lipidomes and diurnal oxylipins using targeted lipidomics of oxylipins and untargeted lipidomics (Figure 4A,B) of the kidney and plasma. The percentage of rhythmic lipids in all lipid classes was approximately 24.3% in the renal lipidome. Timed feeding of EpSA specifically reprogrammed the lipidome of the renal samples, as indicated by principal component analysis (PCA) and heatmap visualization of diurnally oscillating transcripts (Figures S7 and S8). Regarding specific lipid classes, the oxylipins class (37.0%) was one of the most rhythmic lipid classes in the kidneys of mice (Figures 4B and 5A). Subsequently, we identified rhythmic oxylipin metabolites in the plasma and kidney computed by CircaCompare. As expected, several epoxides and their corresponding diols exhibited pronounced circadian variations in the mouse kidneys and plasma (Figures 4C–Q and 5D–R; Figure S9). In the kidney, 8,9-EET, 11,12-EET, and 14,15-EET formation are rhythmic (Figure 4C–E) with the zenith reached around the phase of ZT5 (light phase), whereas the corresponding 8,9-DHET and 11,12-DHET formation also exhibited rhythmicity with a 24-h period (Figure 4H–J). However, 14,15-DHET showed less rhythmicity in the kidneys. Both 9,10-EpOME and 12,13-EpOME exhibited rhythmic variations with zenith levels around ZT6.1 and ZT3.7 (light phase) respectively (Figure 4F,G) and exhibited nadir levels around ZT13 (early of dark phase), whereas 9,10-DiHOME showed only slight circadian variations (Figure 4K,L). Certain oxylipins, such as multiple types of EETs (including 8,9-EET, 14,15-EET), prostaglandins, thromboxane (TX)-A2, and leukotrienes, exhibit a rhythmic patterns, with their production or concentration significantly changing at different times of the day in the plasma, brain, urine, and vasculature [24–27]. Consistent with the findings of previous studies, the total EETs and EpOMEs formation in the kidney and plasma showed a significant day-night rhythm in amplitude across 24 h (Figures S9A and S9B; Figure 5B,C), peaking during the light/inactive phase.

Based on the previous data, we hypothesized that the observed alteration in oxylipin metabolism in the kidneys of timed-fed oxidized oil might be due to transcriptional oscillations of epoxide hydrolase genes induced by EpSA. Indeed, the timed feeding of EpSA during the light phase, but not the dark phase, selectively dampened the oscillation of rhythmic EETs and EpOMEs abundance in the kidney while increasing the oscillation of respective diols (DHETs and DiHOME) abundance in the kidney (Figure 4C–Q). We speculated that the balance between EETs/EpOME and DHET/DiHOME generation in the kidney may be due to the rhythmic expression of epoxide hydrolase genes. EETs and EpOMEs could be hydrolyzed to DHETs and DiHOME, respectively, by epoxide hydrolase. Interestingly,

27662098, 2026, 1, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/EXP.20240073 by CochraneChina, Wiley Online Library on [08/06/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License




FIGURE 3 | Timed-imposed oxidized oil during the inactive period disrupts the circadian transcriptome related to oxylipins metabolism. (A) Schematic of the experimental protocols indicating time window of food access during a 24 h day under timed feeding of HFD and HFD containing EpSA (n = 28 mice per group, n = 4 mice per time point for 7 time points). Transcriptomics were then performed. (B) A total of 282 dark/night and 982 light/day in-phase circadian transcripts, and these 982 light in-phase circadian transcripts reduced to 533 transcripts when fed with EpSA under light-phase, with 55 common to both. (C) Median expression profiles of circadian transcripts in HFD and EpSA treated mice. White and black blocks indicate the light (ZT1, 5, and 9) and dark (ZT13, 17, and 21) time points, respectively. (D) Scatter plots of JTK_CYCLE results of diurnally oscillating transcripts in kidney of HFD and EpSA treated mice. (E) Representative expression of transcripts related to epoxide hydrolases (Ephx1, Ephx2, Ephx3, Ephx4) oscillating in the kidney of Ctrl (green) mice, mice timed-fed with EpSA during the light phase (yellow). Phase parameter were calculated by MetaCycle (meta2d_phase). AMP, amplitude. Mesor, rhythm-adjusted mean expression level. (F) Protein levels of Ephx1 and Ephx2 (n = 4) in kidney were measured in kidney of HFD and EpSA treated mice.

27662098, 2026, 1, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/EXP.20240073 by CochraneChina, Wiley Online Library on [08/06/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License




FIGURE 4 | Loss of circadian rhythmicity in renal oxylipins by timed feeding of epoxy stearic acid during the inactive period. (A) Schematic of the experimental protocols indicating time window of food access during a 24 h day under timed feeding of Normal diet, HFD, HFD containing EpSA. Lipidomics and lipidomics targeted Oxylipins were then performed ($n = 21$ mice per group, $n = 3$ mice per time point for 7 time points). (B) Heat map of dietary EpSA effects on circadian oxylipins profile in kidney. White and black blocks indicate the light (ZT1, 5, and 9) and dark (ZT13, 17, and 21) time points, respectively. (C–L) Representative circadian profiles of oxylipins oscillating in the kidney of Ctrl (green) mice, mice timed-fed with EpSA during the dark phase (red), mice timed-fed with EpSA during the light phase (yellow). (C) 8,9-EET, (D) 11,12-EET, (E) 14,15-EET, (F) 9,10-EpOME, (G) 12,13-EpOME, (H) 8,9-DHET, (I) 11,12-DHET, (J) 14,15-DHET, (K) 9,10-DiHOME, (L) 12,13-DiHOME). (M–Q) Representative circadian ratio of diols/epoxides oscillating in Ctrl (green) mice, mice timed-fed with EpSA during the dark phase (red), mice timed-fed with EpSA during the light phase (yellow). (M) 8,9-DHET/8,9-EET, (N) 11,12-DHET/11,12-EET, (O) 14,15-DHET/14,15-EET, (P) 9,10-DiHOME/9,10-EpOME, (Q) 12,13-DiHOME/12,13-EpOME. Statistics and comparison of rhythmicity parameters were computed by CircaCompare. Arrhythmic when $p > 0.05$.

27662098, 2026, 1, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/EXP.20240073 by CochraneChina, Wiley Online Library on [08/06/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License




FIGURE 5 | Loss of circadian rhythmicity in plasma oxylipins by timed feeding of epoxy stearic acid during the inactive period. (A) Heat map of dietary EpSA effects on circadian oxylipins profile in plasma ($n = 21$ mice per group, $n = 3$ mice per time point for 7 time points). White and black blocks indicate the light (ZT1, 5, and 9) and dark (ZT13, 17, and 21) time points, respectively. (B, C) Representative circadian profiles of total EETs (B) and total EpOMEs (C) oscillating in the plasma of Ctrl (green) mice, mice timed-fed with EpSA during the dark phase (red), mice timed-fed with EpSA during the light phase (yellow). (D–M) Representative circadian profiles of oxylipins oscillating in the plasma of Ctrl (green) mice, mice timed-fed with EpSA during the dark phase (red), mice timed-fed with EpSA during the light phase (yellow). (D) 8,9-EET, (E) 11,12-EET, (F) 14,15-EET, (G) 9,10-EpOME, (H) 12,13-EpOME, (I) 8,9-DHET, (J) 11,12-DHET, (K) 14,15-DHET, (L) 9,10-DiHOME, (M) 12,13-DiHOME). (N–R) Representative circadian ratio of diols/epoxides oscillating in Ctrl (green) mice, mice timed-fed with EpSA during the dark phase (red), mice timed-fed with EpSA during the light phase (yellow). (N) 8,9-DHET/8,9-EET, (O) 11,12-DHET/11,12-EET, (P) 14,15-DHET/14,15-EET, (Q) 9,10-DiHOME/9,10-EpOME, (R) 12,13-DiHOME/12,13-EpOME. Statistics and comparison of rhythmicity parameters were computed by CircaCompare. Arrhythmic when $p > 0.05$.

27662098, 2026, 1, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/EXP.20240073 by CochraneChina, Wiley Online Library on [08/06/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

previous studies have shown that reduced EET levels contribute to kidney inflammation, fibrosis, and loss of renal function [28]. By contrast, DHETs play opposing roles in the regulation of renal function [28]. In the control group, except for the renal 9,10-DiHOME/EpOME ratio, the ratios of 8,9-DHET/EET, 11,12-DHET/EET, 14,15-DHET/EET, and 12,13-DiHOME/EpOME were found to be rhythmically up- and downregulated over a 24-h period in a diurnal fashion (Figure 4N–Q). However, the rhythmic ratio of DHET/EET in the kidney peaked at different times in the control, DFO, and LFO mice. The rhythmic DHET/EET ratio peaked almost exclusively during the light phase in the kidneys of mice with timed feeding of EpSA during the light phase, with the exception of the 12,13-DiHOME/EpOME ratio peaking towards the middle of the dark phase. In addition, timed feeding of EpSA reduced the total EETs and total EpOMEs levels (Figure S6), and elevated the peak levels of diols and the diols/epoxides ratio in the kidney and plasma (Figures 4N–Q, and 5N–R). These data support the novel hypothesis that rhythmic disruption of epoxide hydrolase genes results in rhythmic variations in the metabolism of EETs, EpOMEs, and the corresponding diols in the kidney, potentially contributing to diurnal changes in the regulation of renal function.

2.5 | Timed Feeding of Epoxy Stearic Acid Elicits Time-of-the-Day-Dependent Effects on Renal Injury by Disturbing EHs

Diet quality, timing, and frequency of food intake play a key role in metabolic health, including renal function, and previous observational studies have suggested that late-night dinner (≥3 times per week) and bedtime snacking (≥3 times per week) are associated with an elevated risk of insulin resistance and chronic kidney disease [1, 6, 29]. Late-night snacks are typically high in energy density, often including fried foods [30]. Thus, to explore how timed feeding of EpSA regulates renal function, we compared the phenotypes of mice with time-restricted access to HFD containing EpSA in the light (LFO) or dark (DFO) phase and mice with ad libitum access to HFD, as illustrated in Figure 6A. Compared with observations in the control group, histological analysis of kidneys using HE, PAS, and Masson staining (Figure 6B) demonstrated a significant increase in glomerular size and fibrosis area increased significantly in the LFO group, accompanied by renal tubular vacuolar changes, mesangial cell proliferation, and basement membrane thickening. DFO treatment showed no obvious tubular injury or fibrosis in the kidney, although EpSA intake was low in mice fed with time-imposed EpSA during the inactive period (Figure S10). These structural changes are direct evidence of time-of-day-dependent differences in the harmful effects of dietary EpSA intake. In addition, serum BUN and creatinine levels in the LFO group were significantly higher than those in the control and DFO groups (Figure 6C).

By performing RNA-seq-based profiling of renal samples from the control and LFO group, KEGG enrichment analysis showed that timed feeding (light phase) of EpSA primarily affected pathways related to lipid metabolism (including oxylipin metabolism), oxidative phosphorylation, PI3K-AKT, AMPK, MAPK signaling, insulin signaling, VEGF signaling, and insulin resistance (Figure 6D and Figure S11). The LFO group also showed increased expression and protein levels of Ephx1 and Ephx2 expression and protein levels (Figure 6E and Figure S12). Remarkably, a variety of studies have found that the physiological role of EETs is linked to the activation of the AMPK, PI3K-AKT, and MAPK signaling pathways in insulin resistance, obesity, and kidney injury [31, 32]. Given that increased expression of EH-encoding genes and the altered diol/epoxide ratio during the light phase with EpSA feeding, our data allow us to suppose that timed feeding of fried oil and its oxidation production (EpSA) disrupts the renal epoxide hydrolase clock to exacerbate the impairment of renal function. As expected, administration of EpSA elevated α-SMA and E-cadherin expression in TCMK-1 cells treated with high glucose (Figure 6F,G and Figure S13). However, in Ephx1-KO (Figures 6H,I) and Ephx2-KO TCMK-1 cells (Figure S14) treated with high glucose, the α-SMA and E-cadherin expression levels remained unchanged across groups. We also treated mice with AUDA, a potent pharmacological inhibitor of soluble EH (sEHs), to mimic the effects of sEH blockade. Timed feeding of EpSA during the light phase aggravated renal injury, whereas it failed to cause renal injury and fibrosis in AUDA-treated mice (Figures 7D–F), as shown by insignificant changes in serum BUN and creatinine levels (Figure 7A–C). These data support the hypothesis that EpSA elicits time-dependent effects on renal injury by disturbing EHs.

2.6 | Linoleic Acid Inhibits Epoxide Hydrolysis and Rescues Renal Injury Induced by EpSA

Among food lipids and associated minor content of components, linoleic acid (LA) has been identified as a natural competitive-type inhibitors of epoxide hydrolase, with a Ki value of 3.52 μM [33]. In this study, we assessed the effect of LA on EH activity in mouse renal tubular epithelial (TCMK-1) cells. Through in vivo sEHs activity assessment, linoleic acid inhibited sEH activity (Figure S15), and markedly reduced Ephx2 protein levels in TCMK-1 cells (Figure S15). In addition, dietary LA attenuated kidney injury caused by time-imposed feeding of EpSA during the inactive period. Histological assessment of kidney morphology using HE and Masson staining showed improvements in renal proximal tubular dilatation and tubular basement membrane thickness, along with reduced tubulointerstitial fibrosis in mice treated with an LA-rich diet compared to those treated with EpSA alone (Figure 7G–J).

3 | Discussion

A major goal of chrononutrition is the application of circadian clock biology for disease prevention [34, 35]. More specifically, the emerging field of chronononutrition implements and translates findings from temporal eating patterns and biological rhythms to improve overall health by studying the interactions between the circadian system, dietary patterns, and nutrition [36]. A growing body of evidence has demonstrated the adverse metabolic consequences of circadian misalignment due to the timing of food intake [37]. In modern society, the observation that many young Chinese are abandoning the established norm of three meals per day and making late-night snacks (usually full of fried foods, such as barbecue skewers, hotpot, and grilled seafood) a staple has raised interest in potential health outcomes, which may be attributed to mistimed food

27662098, 2026, 1, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/EXP.20240073 by CochraneChina, Wiley Online Library on [08/06/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License




FIGURE 6 | Epoxy stearic acid elicits time-of-the-day-dependent effects on renal injury via disturbing EHs. (A) Schematic of the experimental protocols indicating time window of food access during a 24 h day under timed feeding of Normal diet, HFD, HFD containing EpSA (n = 7–8 mice per group). (B) Representative images of HE, PAS, Masson staining in kidney of mice (n = 3 mice per group). Arrows represent renal glomerular damage, swelling of renal tubular epithelial cells, or glomerular fibrosis. (C) Volcano plot of gene expression from RNA-seq analysis (Ctrl versus Light-fed EpSA; fold change, ≥2; q value < 0.05) (n = 4 mice per group). (D) KEGG analysis of differentially expressed genes (n = 4 mice per group). (F, G). Immunofluorescence of α-SMA and E-cadherin (F) in EpSA-treated TCMK-1 cells under high glucose. The fluorescent intensity of the α-SMA and E-cadherin (G) was examined (n = 4). (H, I). Immunofluorescence of α-SMA and E-cadherin (H) in WT and Ephx1-KO TCMK-1 cells treated with EpSA. The fluorescent intensity of the α-SMA and E-cadherin (I) was examined (n = 3). Two-way ANOVA were used for statistical test. Data are mean ± SEM. ns, no significant difference, * P <0.05, ** P <0.01.

intake and low food quality. We hypothesized that mistimed fried food intake may adversely affects metabolic health. Although eating during specific periods of time is fundamentally linked to circadian rhythms, the mechanisms by which alterations in dietary nutrients regulate the circadian rhythms in specific metabolic pathways have yet to be elucidated. Our work shows that by investigating the time-dependent effects of fried oil intake on renal function, we found that time-restricted feeding of fried oil during the active period resulted in limited renal damage, whereas fried food intake during the inactive period posed a substantial risk to renal health. In addition, circadian disruption of oxylipin metabolism and epoxide hydrolase was found to exacerbate renal injury under a time-restricted diet of fried oil during the active period. These data may help late-night diners establish guidelines for personalizing food timing to avoid health risks associated with late-night fried food intake.

Metabolic processes and circadian rhythms are tightly linked, with many transcripts and metabolites oscillating in the appropriate molecular environment [38]. This is supported by our findings that 282 dark/night and 982 light/day in-phase circadian transcripts were identified across all renal transcriptomes and 24.3% of rhythmic lipids were identified in all renal lipidomes, under regular eating patterns. Interestingly, renal oxylipins—one of the most rhythmic lipid classes in the kidneys of mice—exhibit circadian oscillations, including epoxide substrates and their

27662098, 2026, 1, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/EXP.20240073 by CochraneChina, Wiley Online Library on [08/06/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License




FIGURE 7 | Timed feeding of epoxy stearic acid during the inactive period failed to cause Renal injury in AUDA-treated mice. (A) Representative images of HE, PAS, Masson staining in kidney of EpSA-treated mice. Arrows represent renal glomerular damage, swelling of renal tubular epithelial cells, or glomerular fibrosis. (B) Serum BUN levels in EpSA-treated mice ($n = 6$). (C) Serum creatinine levels in EpSA-treated mice ($n = 6$). (D). Representative images of HE, PAS, Masson staining in kidney of AUDA-treated mice. (E) Serum BUN levels in AUDA-treated mice ($n = 6$). (F) Serum creatinine levels in AUDA-treated mice ($n = 6$). (G). Representative images of HE, PAS, Masson staining in mice treated with LA-rich diet. Arrows represent renal glomerular damage, slight swelling of renal tubular epithelial cells, or glomerular fibrosis. (H) Renal damage scores in mice treated with LA-rich diet ($n = 6$). (I) Serum BUN levels in mice treated with LA-rich diet ($n = 6$). (J) Serum creatinine levels in mice treated with LA-rich diet ($n = 6$). Two-way ANOVA were used for statistical test. Data are mean ± SEM. ns, no significant difference, * $P < 0.05$, ** $P < 0.01$.

associated EH hydrolysates. Remarkably, our renal lipidomic analysis showed that numerous EETs and EpOME species accumulated in a circadian manner, all reaching their peak levels around ZT5-8 (light phase). In this study, expression analysis of renal enzymes involved in oxylipin metabolism revealed a circadian accumulation pattern of epoxide hydrolase transcripts (e.g., Ephx1 and Ephx2; Figure 3E) in control mice. Similarly, the associated diols also accumulate in a circadian manner, peaking during the night phase. Certain oxylipins, such as multiple types of EETs, prostaglandins, thromboxane A2, and leukotrienes, exhibit rhythmic patterns in the plasma, brain, urine, and vasculature [24–27]. A plausible explanation for this is that post-transcriptional control of epoxide hydrolases in a temporal manner may determine oxylipin metabolism. Notably, the circadian rhythm of epoxide substrates and the formation of associated hydrolysates in the kidney and plasma exhibited a significant light–dark pattern, suggesting that irregular feeding times and poor food quality may alter oxylipin metabolism.

Many studies have highlighted the risk of deep-frying oil for metabolic syndromes [39, 40]. Our previous studies identified epoxy fats, such as epoxy triglyceride monomers and epoxy fatty acids, as early components of deep oxidative polymerization and as the principal cytotoxic products associated with deep-frying oil. Notably, we propose that the risks of epoxy triglycerides and epoxy fatty acids generated from deep frying [41, 42] may have been underestimated, given our evidence that the intake of epoxy fats causes hepatic injury, nerve damage, and intestinal injury [43, 44]. However, the effects of epoxy fat on kidney injury are not thoroughly understood. Since the kidneys are a major site of detoxification in the human body, changes in renal injury, as determined by histological examination, have been observed in response to the dietary intake of oxidized oil in murine models under both ad libitum feeding and extreme time-restricted feeding patterns that consolidate all-calorie intake to the light/inactive phase. Similarly, following a period of time-feeding with oxidized oil during the inactive phase of the day, the mice showed aggravated renal dysfunction and renal fibrosis (Figure 1). By contrast, the mice that received oxidized oil during the active phase did not show any significant pathological renal damage. These findings demonstrate that renal function is influenced not only by the time of day but also by diet quality.

Furthermore, timeimposed feeding with fried oil activated the expression of EHs encoding genes (Ephx1 and EPhx2). This activation was corroborated by changes in oxylipin levels. Timed feeding of oxidized oil in the inactive period worsens renal oxylipin metabolism via EH activation. Specifically, 9,10-EpOME and several epoxyeicosatrienoic acids were reduced by the inactive timed-feeding of oxidized oil, whereas 9,10-DiHOME and DHETs were increased. An imbalance between epoxides and the corresponding diols was observed in the kidneys of mice fed with timed feeding of oxidized oil during the inactive period. Epoxide hydrolases (EHs), which are key enzymes involved in the detox-

27662098, 2026, 1, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/EXP.20240073 by CochraneChina, Wiley Online Library on [08/06/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

ification of xenobiotics and the biotransformation of epoxides, catalyze the hydrolysis of exogenous and endogenous epoxides to their corresponding diols, which are usually inactive or less active than epoxide substrates [45]. The inhibition of epoxide hydrolase, resulting in the upregulation of epoxide substrates (e.g., EETs and EpOME) was renoprotective in many renal diseases [46–49]. As previously mentioned, we unveil the time-of-day dependence of oxidized oil on the regulation of renal injury and oxylipin metabolism. Given that circadian alignment and dietary timing are associated with longevity and lipids metabolism [50–53], we speculated that renal injury induced by time-imposed feeding of oxidized edible oils during the daytime (inactive phase) may be related to the loss of circadian rhythmicity in oxylipins metabolism. Although the effects of the epoxide hydrolase on renal function have now been well documented, much less is known about how the circadian clock alters EH-derived oxylipin metabolic processes, thereby regulating renal function [14, 54].

The presence of epoxy fatty acids in oxidized edible oils is considered a hazardous lipid oxidation products [20, 55]. Intriguingly, EpSA in oxidized edible oils also activates EH activity and the expression of EH-encoding genes in tubular epithelial cells, suggesting that EpSA in oxidized edible oils was involved in the regulation of oxylipin metabolism. Circadian rhythms in renal physiology are becoming increasingly important in kidney health [14]. Consistent with previous study showing that the acrophase of plasma EET levels occurs during the light phase of a 24-h light–dark cycle [25], we determined that the circadian rhythm of epoxide substrates and the associated hydrolysate formation in kidney and plasma showed a significant light–dark manner, by assessment of the diurnal rhythms in the kidneys of mice collected at 4-h intervals over 24 h. In addition, certain oxylipins show rhythmic variations and the rhythmic production of multiple types of EETs has been observed in both endothelial cells and astrocytes [24–27]. We investigated the circadian rhythmicity phenotype after time-imposed feeding of epoxy stearic acid (EpSAs) from oxidized oil. Disruption of circadian rhythmicity during timed feeding of EpSA was observed for both day- and night-peaking transcripts, with many transcripts losing rhythmicity and showing a significant reduction in circadian amplitude of expression, as indicated by the JTK_CYCLE results of diurnally oscillating transcripts. Notably, we observed that the timed feeding of EpSA completely disrupted the rhythmicity of epoxide hydrolases-coding genes in the kidney (Figure 3). Similarly, timed feeding of EpSA during the day (inactive) periods, but not during the dark period, selectively dampened the oscillation of rhythmic EETs and EpOMEs abundance in the kidney while increasing the oscillation of respective diols (DHETs and DiHOME) abundance in the kidney (Figure 4). Nevertheless, evidence suggests that EH inhibitors and other means of manipulating EETs have beneficial effects in chronic kidney disease and inflammation [46–49]. Indeed, our results suggest that the dietary intake of EpSA produces time-of-day-dependent changes in harmful effects, with time-imposed feeding of EpSA only during the daytime (inactive phase) worsening renal injury.

Clinical trials on EH inhibition have highlighted the therapeutic potential of epoxide hydrolases regulation [56]. Our data allow us to suppose that the alteration in oxylipin profiles may be associated with the development of kidney injury following exposure to oxidized oil and EpSA during the light phase. Mechanistically, we demonstrated that time-imposed feeding of EpSA during the daytime (inactive phase) of EpSA induced the circadian degradation of EETs and disrupted their transcriptional oscillations, which can impede the renoprotective actions of EETs. The effect of EpSA on SMA and E-cadherin, two specific biomarkers for renal inflammation and fibrosis, was also lost in *Ephx1* and *Ephx2* knockout cells, which lack EHs. This is further supported by the fact that renal injury did not worsen with daytime-restricted feeding of EpSA in mice treated with AUDA (an sEH inhibitor). In our study, we also found that dietary LA, a natural competitive-type EH inhibitor [33], revealed markedly reduced kidney injury induced by dietary EpSA through EHs inhibition.

Although our results reveal that the time-of-day dependence of epoxide hydrolases and circadian disruption during the inactive phase by EpSA from thermally induced oxidized oil have adverse effects on kidney function, it is unclear how frying oil regulates the circadian rhythm of epoxide hydrolase. Additionally, other hydrolases (such as triglycerol hydrolase Ces1d) related to lipid metabolism and activated by dietary changes also exhibit circadian features. Further in-depth research is required to understand the interaction between the circadian rhythms of EHs and renal injury. In conclusion, our data demonstrate that the time of day defines circadian variation in the production of epoxy-oxylipin metabolites and EH expression, with this rhythmic signaling contributing to diurnal differences in renal injury when fried oil and EpSA from fried oil are consumed. By understanding these relations, we will be better informed about the role of meal timing and food choice in renal health and the importance of intentionally modifying dietary patterns and nutrition in late-night eating for better health.

4 | Methods

4.1 | Preparation of Fried Oil (Thermo-Induced Oxidized Oil From Palm Oil)

Fried oil was obtained by heating palm oil at 180°C to a TPC (Total Polar Compounds) value of 27 in an oil bath in the laboratory. Total polar compound levels in frying oils were analyzed according to our method previously reported [57]. The composition of TPC in fried oil samples was 20.02% polymerized triglycerides (PG), 30.01% oxidized triglycerides (ox-TG) and 49.47% triglyceride degradation products (including free fatty acids and 2-monoacylglycerol).

4.2 | Mice and Diet

All of the experimental procedures followed the Guide for the Care and Use of Laboratory Animals: Eighth Edition (ISBN-10: 0-309-15396-4), and the animal experiments was approved by the animal ethics committee of Jiangnan University (JN.No20210630c1921001) in July 2021. 6-week-old male C57BL/6 mice were maintained in a 12:12-h light–dark (LD) cycle. Long-term high-fat diet (HFD) feeding was used to induce renal injury in mice [58]. 6-week-old mice (SiPeiFu Biotechnology Co., Ltd, Beijing, China) fed a standard chow diet (Control, XTCON50J,

27662098, 2026, 1, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/EXP.20240073 by CochraneChina, Wiley Online Library on [08/06/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

13% calories from fat), a high-fat diet (HFD, XTHF60, 60% energy from fat), or a high-fat diet containing 3% fried oil (thermo-induced oxidized oil from palm oil) for 14 weeks, in which palm oil or oxidized oil as a fat source was incorporated (XTHF60, Xietong Shengwu, Jiangsu, China). Another cohort of 6-week-old mice fed a standard chow diet, or a high-fat diet containing 3% palm oil, or a high-fat diet containing 3% fried oil (thermo-induced oxidized oil from palm oil) for 14 weeks. These mice were fed under AL feeding (Ad libitum feeding) or light-restricted feeding (ZT0-ZT12). The third cohort of 6-week-old mice fed a standard chow diet, or a high-fat diet containing 3% palm oil, or a high-fat diet containing 3% fried oil (thermo-induced oxidized oil from palm oil) for 14 weeks. A high-fat diet containing 3% fried oil was time-imposed on mice during the light phase (LFO, ZT0-ZT12) or dark phase (DFO, ZT12-ZT24). The other cohort of 6-week-old mice fed a standard chow diet, a high-fat diet, or a high-fat diet containing 3% EpSA for 14 weeks. A high-fat diet containing 3% EpSA was time-imposed to mice during the light phase (ZT0-ZT12). ZT0 denotes lights-on and ZT12 denotes lights-off. The last cohort of 6-week-old mice fed a high-fat diet containing 3% EpSA or a high-fat diet containing 3% EpSA and 3% linoleic acid for 14 weeks. These mice were time-restricted to mice during the light phase (ZT0-ZT12). The renal injury score was assessed in a blinded manner and scored by the percentage of injured tubules, as follows: 0, no damage; 1, <25%; 2, 25%–50%; 3, 50%–75%; 4, >75%. Semi quantitative assessment of fibrosis area was carried out using Image Pro Plus 6.0 (Media Cybernetics).

4.3 | Plasmid Construction

The PX459 plasmid (#48139, Addgene, Cambridge, MA, USA) was used to construct Ephx1 and Ephx2 KO plasmids [59]. PX459 plasmid was a gift from Feng Zhang, Broad Institute, Cambridge, MA, obtained via Addgene (62988). Ephx1 and Ephx2 KO by CRISPR/Cas9 was performed as described previously. Insert oligonucleotides that include a guiding RNA sequence were designed as follows: for Ephx1 KO, GGAGCTCTTGTACCCATTCA and GCCCCTCGGGTGATTCCGTTC; for Ephx2 KO, GCTTGATCATCCCAACCTGACG and GTCCCTCTGGGAATTCCGTC. After annealing, these oligonucleotides were inserted to the BbsI cloning site of PX459. These plasmids and unmodified PX459 plasmid were transfected to TCMK-1 cells by Lipofectamin 2000. The validation of knockout clones were shown in Figures S16 and S17.

4.4 | Cell Culture

TCMK-1 cell (ATCC CCL-139TM, Pricella Biotechnology Co., Ltd., China), the tubular epithelial cells in mice were maintained in MEM medium with 10% FBS. TCMK-1 cell was cultured at 37°C, 95% air, and 5% CO_2.

4.5 | Oxylipin Profiling

As for serum samples, precisely aspirate 90 μL of plasma under dark conditions and place it in a 1.5 mL centrifuge tube. Add 5 μL of methanol, vortex for 30 s, and then add 500 μL of ethyl acetate, vortex for 3 min and let it stand for 1 min. Centrifuge at 20,000 rpm at 4°C for 5 min. Transfer the supernatant to a 10 mL covered glass tube, dry it with nitrogen gas at room temperature, dissolve the residue with 200 μL mobile phase, vortex for 30 s, and transfer it to a 1.5 mL centrifuge tube. Centrifuge at 20,000 rpm at 4°C for 5 min, and pass the supernatant through a 0.22 μm filter membrane, Perform HPLC MS/MS analysis. As for kidney samples, grind the solid sample in a liquid nitrogen environment until crushed, take approximately 100 mg of each tissue sample, rinse with physiological saline, dry with filter paper, and weigh. Add 500 μL of methanol (containing 0.01 mol/L BHT) and 10 μL of formic acid, mix well, homogenize in a homogenizer, and sonicate to break the cells (working time 3 s, gap time 2 s, working 60 times, power 400 W). The above process is carried out in an ice bath. Centrifuge at 4°C for 20 min (15000 r/min), remove the supernatant, add 700 μL of water and 1 mL of ethyl acetate, vortex for 30 s, centrifuge at 12000 r/min for 10 min, remove the organic phase, extract the aqueous phase again, merge the ethyl acetate layer, freeze dry the solvent, dissolve the residue in 200 μL of methanol, centrifuge at 20000 rpm for 5 min at 4°C, take the supernatant and pass through a 0.22 μm filter membrane for HPLC MS/MS analysis.

4.6 | Lipidomic Analysis

Collected kidney were frozen in liquid nitrogen and homogenized. To the homogenized sample, methanol was added and homogenized in the identical condition. 200 μL of suspension in methanol was mixed with 100 μL chloroform and incubated for 1 h. Subsequently, 20 μL of water was added. After extraction, samples were centrifuged at 3500×g for 25 min, and the supernatants were collected. Untargeted lipidomic was performed by using an UPLC (AB SCIEX ExionLC) coupled with a TOF-MS (AB SCIEX, Triple TOF 5600+). Chromatographic separations were performed on a Kinetex C18 columns (100 mm x 2.1 mm, 2.6 μm, Phenomenex). The mobile phase A was an acetonitrile-water mixture containing 10 mM ammonium acetate (acetonitrile: water = 2:3, v/v), and mobile phase B was an acetonitrile-isopropanol mixture containing 10 mM ammonium acetate (acetonitrile: isopropanol = 1:9, v/v). The elution gradient was started with 35% B keeping for 1 min, increasing to 60% B in 1 min, then rising to 100% B in 11 min and holding for 3.5 min. Next, the elution fell to 35% B in 0.5 min and maintaining for 1.5 min. The ESI source was set at 500°C, decluttering potential 90 V, collision energy 40 ± 20 V, spray voltage of 5500 V in positive mode or −4500 V in negative mode.

4.7 | RNA-seq

Frozen tissue samples were homogenized and RNA was isolated with Spin Column Animal Total RNA Purification Kit (Sangon Biotech (Shanghai) Co., Ltd., B518651) according to the manufacturer's instructions. 1 μg total RNA was used for following library preparation. Sequencing was performed on an Illumina Novaseq platform using a 2 × 150 paired-end (PE) configuration according to manufacturer's instructions. Renal transcriptome analysis, rhythmic analysis, and Phase parameter analysis were calculated by MetaCycle (meta2d_phase). MetaCy-

27662098, 2026, 1, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/EXP.20240073 by CochraneChina, Wiley Online Library on [08/06/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

cle package is available on the CRAN repository (https://cran.r-project.org/web/packages/MetaCycle/index.html) and GitHub (https://github.com/gangwug/MetaCycle).

4.8 | Immunofluorescence Staining

Immunofluorescence staining were performed in OCT-embedded kidneys section. The following antibodies were used for the immunofluorescence analyses. Nuclei were counterstained with DAPI. The slides were inoculated in primary antibody anti-E-cadherin (ab231303, Abcam, Cambridge, USA, 1:1000) and anti-α-SMA (ab5694, Abcam, Cambridge, USA, 1:200). Alexa Fluor 488-labeled goat anti-rabbit IgG (111-585-003; Jackson) and Goat anti-mouse IgG-CY3 (GB21301; Servicebio) were used as secondary antibodies. Fluorescence images were gathered at 200× and 400× magnification using an upright fluorescence microscope (NIKON ECLIPSE C1, Nikon, Japan). The fluorescence intensity was computed using Image Pro Plus 6.0 (Media Cybernetics).

4.9 | Quantitative Real-Time PCR Analysis (QT-PCR)

TRIzol reagent (Yeason, China) was used to extract the total RNA from kidney, and cDNA Synthesis Mix was used to reverse-transcribe the recovered RNA (Yeason, China). The QT-PCR was carried out on a QuantStudio 5 System. The internal reference utilized to calculate the relative expression level of mRNA was 36B4. Primer sequences were detailed in Table S1.

4.10 | Western Blot Analysis

Western blot analysis protocols have been previously described [60]. All kidney tissue were homogenized in RIPA lysis buffer (Beyotime, Shanghai, China, #P0013B) supplemented with PMSF (Beyotime, Shanghai, China, #ST505), phosphatase inhibitor cocktail (Beyotime, Shanghai, China, #P1082) and protease inhibitor cocktail (Beyotime, Shanghai, China, #P1006). The total protein were extracted and were separated by SDS-PAGE, transferred to PVDF membranes (Millipore, Billerica). Membranes were blocked in a blocking buffer (Beyotime, Shanghai, China, #P0023B) and incubated with primary antibodies. Primary antibody anti-E-cadherin (ab231303, Abcam, Cambridge, USA, 1:2000), anti-α-SMA (ab5694, Abcam, Cambridge, USA, 1:2000), anti-Ephx1 (66982-1-Ig, Proteintech, Wuhan, 1:2000), anti-Ephx2 (10833-1-AP, Proteintech, Wuhan, 1:1000), and Tubulin antibody (10094-1-AP, Proteintech, 1:5000) were used. BeyoECL Plus regent was used to detect the signal.

4.11 | sEHs Activity

Soluble epoxide hydrolases (sEHs) activity was performed using a Soluble Epoxide Hydrolase Cell-Based Assay Kit (Cayman, No.600090, Michigan, USA) according to the manufacturer's instructions.

4.12 | Serum KIM-1 and Nga Levels

Serum KIM-1 and Nga (mouse) were detected by using ELISA Kit (Sangon Biotech (Shanghai) Co., Ltd., Shanghai, China).

4.13 | Statistical Analysis

All results were expressed as means ± SEM. GraphPad Prism (version 8.0, Boston, USA) was used for statistical analyses. Statistical significance was determined by Student's t test or two-way ANOVA models between-group comparisons. When comparing more than 2 groups, we used two-way ANOVA were used for group comparisons and Holm–Sidak method was used for p-value adjustment for multiple comparisons. A p value of 0.05 or less was indicated a significant difference.

Author Contributions

J. Y. L. and M. L. performed the experiments. J. Y. L. designed experiments. J. Y. L. analyzed data. F.Y.L., C. X., Y. Du., M. F. and J.X. helped with data interpretation and discussion. J. Y.L. wrote the paper.

Acknowledgments

This work was supported by National Natural Science Foundation of China (No. 32202056), Youth Cross-innovation Research Program of CFSE-OUC (No. 2024104), Youth Talent Project Launch Fund of Ocean University of China (No. 862401013170), Postdoctoral Fellowship Program of CPSF (No. GZB20240695), and National Science Foundation for Postdoctoral Scientists of China (No. 2021M701465 and No. 2025M772986).

Conflicts of Interest

The authors declare that they have no competing interests.

Data Availability Statement

The data are available from the author (Yanjun Liu) upon reasonable request.

References

1. R. Michishita, T. Matsuda, S. Kawakami, et al., "The Association Between Unhealthy Lifestyle Behaviors and the Prevalence of Chronic Kidney Disease (CKD) in Middle-Aged and Older Men," *Journal of Epidemiology* 26 (2016): 378–385, https://doi.org/10.2188/jea.JE20150202.

2. A. Gallant, J. Lundgren, and V. Drapeau, "Nutritional Aspects of Late Eating and Night Eating," *Current Obesity Reports* 3 (2014): 101–107, https://doi.org/10.1007/s13679-013-0081-8.

3. F. Osman, S. Haldar, and C. J. Henry, "Effects of Time-Restricted Feeding During Ramadan on Dietary Intake, Body Composition and Metabolic Outcomes," *Nutrients* 12 (2020): 2478, https://doi.org/10.3390/nu12082478.

4. J. Kovac, J. Husse, and H. Oster, "A Time to Fast, a Time to Feast: The Crosstalk Between Metabolism and the Circadian Clock," *Molecules and Cells* 28 (2009): 75–80, https://doi.org/10.1007/s10059-009-0113-0.

5. N. A. Costa, A. G. Pereira, M. S. Dorna, et al., "Meal Timing and Frequency Implications in the Development and Prognosis of Chronic Kidney Disease," *Nutrition (Burbank, Los Angeles County, California)* 91-92 (2021): 111427, https://doi.org/10.1016/j.nut.2021.111427.

6. X. Y. Liu, C. L. Zheng, C. Xu, et al., "Nighttime Snacking is Associated With Risk of Obesity and Hyperglycemia in Adults: A Cross-Sectional

27662098, 2026, 1, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/EXP.20240073 by CochraneChina, Wiley Online Library on [08/06/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Survey from Chinese Adult Teachers," *Journal of Biomedical Research* 31, no. 6 (2017): 541–547.

7. T. T. Huang, S. B. Roberts, N. C. Howarth, and M. A. McCrory, "Effect of Screening out Implausible Energy Intake Reports on Relationships Between Diet and BMI," *Obesity Research* 13 (2005): 1205–1217, https://doi.org/10.1038/oby.2005.143.

8. S. E. Noel, C. Mattocks, P. Emmett, C. J. Riddoch, A. R. Ness, and P. K. Newby, "Use of Accelerometer Data in Prediction Equations for Capturing Implausible Dietary Intakes in Adolescents," *American Journal of Clinical Nutrition* 92 (2010): 1436–1445, https://doi.org/10.3945/ajcn.2010.29386.

9. A. B. Clark, A. M. Coates, Z. E. Davidson, and M. P. Bonham, "Dietary Patterns Under the Influence of Rotational Shift Work Schedules: A Systematic Review and Meta-Analysis," *Advances in Nutrition* 14 (2023): 295–316, https://doi.org/10.1016/j.advnut.2023.01.006.

10. L. C. Balieiro, L. T. Rossato, J. Waterhouse, S. L. Paim, M. C. Mota, and C. A. Crispim, "Nutritional Status and Eating Habits of Bus Drivers During the Day and Night," *Chronobiology International* 31 (2014): 1123–1129, https://doi.org/10.3109/07420528.2014.957299.

11. F. M. Suk, F. Y. Hsu, Y. C. Lee, T. L. Chen, W. C. Chiu, and Y. J. Liao, "Dietary Oxidized Frying Oil Activates Hepatic Stellate Cells and Accelerates the Severity of Carbon Tetrachloride- and Thioacetamide-Induced Liver Fibrosis in Mice," *Journal of Nutritional Biochemistry* 115 (2023): 109267, https://doi.org/10.1016/j.jnutbio.2023.109267.

12. H. Motohashi, Y. Tahara, D. Whittaker, et al., "The Circadian Clock Is Disrupted in Mice With Adenine-Induced Tubulointerstitial Nephropathy," *Kidney International* 97, no. 4 (2020): 728–740, https://doi.org/10.1016/j.kint.2019.09.032.

13. C. Lu, X. Cang, W. Liu, et al., "A Late Eating Midpoint is Associated With Increased Risk of Diabetic Kidney Disease: A Cross-Sectional Study Based on NHANES 2013–2020," *Nutrition Journal* 23, no. 1 (2024): 39, https://doi.org/10.1186/s12937-024-00939-z.

14. D. Firsov and O. Bonny, "Circadian Rhythms and the Kidney," *Nature Reviews Nephrology* 14 (2018): 626–635, https://doi.org/10.1038/s41581-018-0048-9.

15. A. Zambelli, "Current Status of High Oleic Seed Oils in Food Processing," *Journal of the American Oil Chemists' Society* 98 (2021): 129–137, https://doi.org/10.1002/aocs.12450.

16. A. Mancini, E. Imperlini, E. Nigro, et al., "Biological and Nutritional Properties of Palm Oil and Palmitic Acid: Effects on Health," *Molecules* 20 (2015): 17339–17361, https://doi.org/10.3390/molecules200917339.

17. Y. Shen, K. Krätschmer, T. Bovee, J. Louisse, and S. P. J. van Leeuwen, "Chlorinated Paraffins (CPs) in Vegetable Oils From the Dutch Market and the Effects of the Refining Process on Their Levels," *Food Control* 153 (2023): 109889, https://doi.org/10.1016/j.foodcont.2023.109889.

18. K. M. Schaich, "Thinking Outside the Classical Chain Reaction Box of Lipid Oxidation," *Lipid Technology* 24 (2012): 55–58, https://doi.org/10.1002/lite.201200170.

19. S. Marmesat, J. Velasco, and M. C. Dobarganes, "Quantitative Determination of Epoxy Acids, Keto Acids and Hydroxy Acids Formed in Fats and Oils at Frying Temperatures," *Journal of Chromatography A* 1211 (2008): 129–134, https://doi.org/10.1016/j.chroma.2008.09.077.

20. L. Brühl, "Fatty Acid Alterations in Oils and Fats During Heating and Frying," *European Journal of Lipid Science and Technology* 116 (2014): 707–715.

21. A. Fankhauser-Noti, K. Fiselier, S. Biedermann-Brem, and K. Grob, "Assessment of Epoxidized Soy Bean Oil (ESBO) Migrating Into Foods: Comparison With ESBO-Like Epoxy Fatty Acids in Our Normal Diet," *Food and Chemical Toxicology* 44 (2006): 1279–1286, https://doi.org/10.1016/j.fct.2006.02.005.

22. M. Grootveld, B. C. Percival, J. Leenders, and P. B. Wilson, "Potential Adverse Public Health Effects Afforded by the Ingestion of Dietary Lipid Oxidation Product Toxins: Significance of Fried Food Sources," *Nutrients* 12 (2020): 974, https://doi.org/10.3390/nu12040974.

23. H. Xin, F. Deng, M. Zhou, et al., "A Multi-Tissue Multi-Omics Analysis Reveals Distinct Kineztics in Entrainment of Diurnal Transcriptomes by Inverted Feeding," *Iscience* 24 (2021): 102335, https://doi.org/10.1016/j.isci.2021.102335.

24. K. A. Carver, D. Lourim, A. K. Tryba, and D. R. Harder, "Rhythmic Expression of Cytochrome P450 Epoxygenases CYP4x1 and CYP2c11 in the Rat Brain and Vasculature," *American Journal of Physiology. Cell Physiology* 307 (2014): C989–C998, https://doi.org/10.1152/ajpcell.00401.2013.

25. T. Kezeli, N. Gongadze, G. Sukoyan, et al., "Circadian Variation in Vasoconstriction and Vasodilation Mediators and Baroreflex Sensitivity in Hypertensive Rats," *J Circadian Rhythms* 17 (2019): 10, https://doi.org/10.5334/jcr.185.

26. K. Kamperis, M. N. Hansen, S. Hagstroem, G. Hvistendahl, J. C. Djurhuus, and S. Rittig, "The Circadian Rhythm of Urine Production, and Urinary Vasopressin and Prostaglandin E2 Excretion in Healthy Children," *Journal of Urology* 171 (2004): 2571–2575, https://doi.org/10.1097/01.ju.0000110421.71910.c0.

27. K. Kurokawa, H. Tanaka, S. Tanaka, and S. Abe, "Circadian Characteristics of Urinary Leukotriene E4 in Healthy Subjects and Nocturnal Asthmatic Patients," *Chest* 120 (2001): 1822–1828, https://doi.org/10.1378/chest.120.6.1822.

28. M. R. Noh, H.-S. Jang, F. E. Salem, F. A. Ferrer, J. Kim, and B. J. Padanilam, "Epoxyeicosatrienoic Acid Administration or Soluble Epoxide Hydrolase Inhibition Attenuates Renal Fibrogenesis in Obstructive Nephropathy," *American Journal of Physiology. Renal Physiology* 324 (2022): F138–F151, https://doi.org/10.1152/ajprenal.00052.2022.

29. R. Mohandas, L. G. Douma, Y. Scindia, and M. L. Gumz, "Circadian Rhythms and Renal Pathophysiology," *Journal of Clinical Investigation* 132 (2022): e148277, https://doi.org/10.1172/JCI148277.

30. W. Si Hassen, K. Castetbon, C. Tichit, et al., "Energy, Nutrient and Food Content of Snacks in French Adults," *Nutrition Journal* 17 (2018): 33, https://doi.org/10.1186/s12937-018-0336-z.

31. M. Peyman, E. Barroso, A. L. Turcu, et al., "Soluble Epoxide Hydrolase-Targeting PROTAC Activates AMPK and Inhibits Endoplasmic Reticulum Stress," *Biomedicine & Pharmacotherapy* 168 (2023): 115667, https://doi.org/10.1016/j.biopha.2023.115667.

32. Z. Shi, Z. He, and D. W. Wang, "CYP450 Epoxygenase Metabolites, Epoxyeicosatrienoic Acids, as Novel Anti-Inflammatory Mediators," *Molecules (Basel, Switzerland)* 27 (2022): 3873.

33. X. He, W. Y. Zhao, B. Shao, et al., "Natural Soluble Epoxide Hydrolase Inhibitors From Inula Helenium and Their Interactions With Soluble Epoxide Hydrolase," *International Journal of Biological Macromolecules* 158 (2020): 1362–1368, https://doi.org/10.1016/j.ijbiomac.2020.04.227.

34. R. Adafer, W. Messaadi, M. Meddahi, et al., "Food Timing, Circadian Rhythm and Chrononutrition: A Systematic Review of Time-Restricted Eating's Effects on Human Health," *Nutrients* 12 (2020): 3770, https://doi.org/10.3390/nu12123770.

35. V. A. Acosta-Rodríguez, F. Rijo-Ferreira, C. B. Green, and J. S. Takahashi, "Importance of Circadian Timing for Aging and Longevity," *Nature Communications* 12 (2021): 2862, https://doi.org/10.1038/s41467-021-22922-6.

36. A. Flanagan, D. A. Bechtold, G. K. Pot, and J. D. Johnston, "Chrono-Nutrition: From Molecular and Neuronal Mechanisms to Human Epidemiology and Timed Feeding Patterns," *Journal of Neurochemistry* 157 (2021): 53–72, https://doi.org/10.1111/jnc.15246.

37. K. G. Baron and K. J. Reid, "Circadian Misalignment and Health," *International Review of Psychiatry* 26 (2014): 139–154, https://doi.org/10.3109/09540261.2014.911149.

27662098, 2026, 1, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/EXP.20240073 by CochraneChina, Wiley Online Library on [08/06/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

38. K. L. Eckel-Mahan, V. R. Patel, S. de Mateo, et al., "Reprogramming of the Circadian Clock by Nutritional Challenge," *Cell* 155 (2013): 1464–1478, https://doi.org/10.1016/j.cell.2013.11.034.

39. Y. F. Chiang, H. M. Shaw, M. F. Yang, C. Y. Huang, C. H. Hsieh, and P. M. Chao, "Dietary Oxidised Frying Oil Causes Oxidative Damage of Pancreatic Islets and Impairment of Insulin Secretion, Effects Associated With Vitamin E Deficiency," *British Journal of Nutrition* 105 (2011): 1311–1319, https://doi.org/10.1017/S0007114510005039.

40. A. Koch, B. König, S. Luci, G. I. Stangl, and K. Eder, "Dietary Oxidised Fat up Regulates the Expression of Organic Cation Transporters in Liver and Small Intestine and Alters Carnitine Concentrations in Liver, Muscle and Plasma of Rats," *British Journal of Nutrition* 98 (2007): 882–889, https://doi.org/10.1017/S000711450775691X.

41. C. Wei, X. Wang, Y. Ai, et al., "Effect of Oleic/Linoleic Acid Ratio on Simultaneous Formation of Epoxy Fatty Acids and Polar Compounds in Deep Frying Oils," *LWT* 187 (2023): 115389, https://doi.org/10.1016/j.lwt.2023.115389.

42. Y. P. Khor, K. S. Hew, F. Abas, et al., "Oxidation and Polymerization of Triacylglycerols: In-Depth Investigations Towards the Impact of Heating Profiles," *Foods* 8 (2019): 475, https://doi.org/10.3390/foods8100475.

43. X. Li, Y. J. Liu, Y. Wang, Y. F. Liu, and Y. J. Xu, "Epoxy Triglyceride Enhances Intestinal Permeability via Caspase-1/NLRP3/GSDMD and cGAS-STING Pathways in Dextran Sulfate Sodium-Induced Colitis Mice," *Journal of Agricultural and Food Chemistry* 71 (2023): 4371–4381, https://doi.org/10.1021/acs.jafc.2c08134.

44. R. Wilson, C. E. Fernie, C. M. Scrimgeour, K. Lyall, L. Smyth, and R. A. Riemersma, "Dietary Epoxy Fatty Acids are Absorbed in Healthy Women," *European Journal of Clinical Investigation* 32 (2002): 79–83, https://doi.org/10.1046/j.1365-2362.2002.00951.x.

45. J. W. Newman, C. Morisseau, and B. D. Hammock, "Epoxide Hydrolases: Their Roles and Interactions With Lipid Metabolism," *Progress in Lipid Research* 44 (2005): 1–51, https://doi.org/10.1016/j.plipres.2004.10.001.

46. J. Gautheron and I. Jéru, "The Multifaceted Role of Epoxide Hydrolases in Human Health and Disease," *International Journal of Molecular Sciences* 22 (2020): 13, https://doi.org/10.3390/ijms22010013.

47. D. Y. Hu, Y. Luo, C. B. Li, et al., "Oxylipin Profiling of human Plasma Reflects the Renal Dysfunction in Uremic Patients," *Metabolomics* 14 (2018): 104, https://doi.org/10.1007/s11306-018-1402-4.

48. M. M. Shuey, F. T. T. Billings, S. Wei, et al., "Association of Gain-of-Function EPHX2 Polymorphism Lys55Arg With Acute Kidney Injury Following Cardiac Surgery," *PLoS ONE* 12 (2017): e0175292, https://doi.org/10.1371/journal.pone.0175292.

49. X. S. Jiang, X. Y. Xiang, X. M. Chen, et al., "Inhibition of Soluble Epoxide Hydrolase Attenuates Renal Tubular Mitochondrial Dysfunction and ER Stress by Restoring Autophagic Flux in Diabetic Nephropathy," *Cell Death & Disease* 11 (2020): 385, https://doi.org/10.1038/s41419-020-2594-x.

50. V. Acosta-Rodríguez, F. Rijo-Ferreira, M. Izumo, et al., "Circadian Alignment of Early Onset Caloric Restriction Promotes Longevity in Male C57BL/6J Mice," *Science* 376 (2022): 1192–1202, https://doi.org/10.1126/science.abk0297.

51. K. G. Baron, K. J. Reid, T. Kim, et al., "Circadian Timing and Alignment in Healthy Adults: Associations With BMI, Body Fat, Caloric Intake and Physical Activity," *Int J Obes (Lond)* 41 (2017): 203–209, https://doi.org/10.1038/ijo.2016.194.

52. J. P. Chaput, A. W. McHill, R. C. Cox, et al., "The Role of Insufficient Sleep and Circadian Misalignment in Obesity," *Nature Reviews Endocrinology* 19 (2023): 82–97, https://doi.org/10.1038/s41574-022-00747-7.

53. L. C. Ruddick-Collins, P. J. Morgan, C. L. Fyfe, et al., "Timing of Daily Calorie Loading Affects Appetite and Hunger Responses Without Changes in Energy Metabolism in Healthy Subjects With Obesity," *Cell Metabolism* 34 (2022): 1472–1485.e6, https://doi.org/10.1016/j.cmet.2022.08.001.

54. J. G. Johnston and D. M. Pollock, "Circadian Regulation of Renal Function," *Free Radical Biology & Medicine* 119 (2018): 93–107, https://doi.org/10.1016/j.freeradbiomed.2018.01.018.

55. N. Kalogeropoulos, F. N. Salta, A. Chiou, and N. K. Andrikopoulos, "Formation and Distribution of Oxidized Fatty Acids During Deep- and Pan-Frying of Potatoes," *European Journal of Lipid Science and Technology* 109 (2007): 1111–1123, https://doi.org/10.1002/ejlt.200700007.

56. J. D. Imig and B. D. Hammock, "Soluble Epoxide Hydrolase as a Therapeutic Target for Cardiovascular Diseases," *Nature Reviews Drug Discovery* 8 (2009): 794–805, https://doi.org/10.1038/nrd2875.

57. M. Lu, J. Shi, X. Li, Y. Liu, and Y. Liu, "Long-term Intake of Thermo-induced Oxidized Oil Results in Anxiety-Like and Depression-Like Behaviors: Involvement of Microglia and Astrocytes," *Food Funct* 15, no. 8 (2024): 4037–4050, https://doi.org/10.1039/D3FO05302D.

58. Y. Sun, X. Ge, X. Li, et al., "High-fat Diet Promotes Renal Injury by Inducing Oxidative Stress and Mitochondrial Dysfunction," *Cell Death & Disease* 11 (2020): 914, https://doi.org/10.1038/s41419-020-03122-4.

59. F. Ran, P. D. Hsu, J. Wright, V. Agarwala, D. A. Scott, and F. Zhang, "Genome Engineering Using the CRISPR-Cas9 System," *Nature Protocols* 8, no. 11 (2013): 2281–2308, https://doi.org/10.1038/nprot.2013.143.

60. Y. Liu, C. Jiao, T. Zhang, et al., "Early-Life Gut Microbiota Governs Susceptibility to Colitis via Microbial-Derived Ether Lipids," *Research (Washington D C)* 6 (2023): 0037.

Supporting Information

Additional supporting information can be found online in the Supporting Information section.

Supporting File 1: exp270124-sup-0001-SuppMat.pdf.

27662098, 2026, 1, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/EXP.20240073 by CochraneChina, Wiley Online Library on [08/06/2026]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License