



Original Article

Asian Pacific Journal of Tropical Biomedicine



apjtb.org

doi: 10.4103/apjtb.apjtb_339_26

Ecological and biological determinants of fecal bacterial diversity in bats across four continents and phylogenetic screening for pathogen–affiliated bacterial lineages

Lucas Palheta Gomes^{1✉}, Raphael de Andrade Morraye², Lucas da Silva e Silva³, João Paulo Bianchi Ximenez⁴, Maíra Pompeu Martins³, Wilson Araújo Silva Jr², Maria Paula Cruz Schneider³

¹Immunoparasitology Laboratory, Institute of Biomedical Sciences, University of São Paulo, São Paulo, Brazil

²Department of Genetics, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, SP, Brazil

³Laboratory of Genomics and Biotechnology, Biological Sciences Institute, Federal University of Pará, Belém, PA, Brazil

⁴Department of Clinical Analyses, Toxicology and Food Science, School of Pharmaceutical Sciences of Ribeirão Preto, University of São Paulo, Ribeirão Preto, SP, Brazil

ABSTRACT

Objective: To characterize the diversity and composition of bacterial communities in bat fecal samples under distinct ecological and biological conditions and to investigate the phylogenetic relationships of selected bacterial lineages with clinically relevant taxa.

Methods: Publicly available 16S rRNA gene sequencing data from 194 fecal samples covering 20 bat species from South America, Europe, Asia, and Africa were reanalyzed using DADA2 and the SILVA database. Alpha and beta diversity, taxonomic composition, and community differences according to host diet, habitat, sex, and geographic region were evaluated. Phylogenetic analyses were performed on amplicon sequence variants assigned to *Pseudomonas*, *Staphylococcus*, *Clostridium*, and *Streptococcus*.

Results: A total of 3778110 reads passed quality filtering, representing 123 bacterial families and 203 genera. Proteobacteria (57%) and Firmicutes (40%) predominated, with *Pseudomonas*, *Staphylococcus*, and *Carnobacterium* among the most abundant genera. Microbial composition and diversity varied significantly according to diet, habitat, sex, and geographic region. Frugivorous bats showed the highest alpha diversity, whereas piscivorous bats showed the lowest, and *Rousettus aegyptiacus* exhibited the greatest bacterial richness. Bats inhabiting forest and cave habitats possessed higher bacterial richness, while communities from freshwater and coastal aquatic habitats were less diverse. Males showed higher relative abundance of *Staphylococcus*, whereas females were enriched in *Pseudomonas*. Distinct taxonomic signatures were linked to dietary guilds. Insectivorous bats were enriched in Gammaproteobacteria, Pseudomonadales, and

Moraxellaceae; frugivorous bats were enriched in Lactobacillales and *Streptococcus*; and piscivorous bats were enriched in Firmicutes, including Staphylococcaceae and *Clostridium*. Sixty-two genera containing species with recognized pathogenic potential were identified. Phylogenetic analyses revealed amplicon sequence variants affiliated with *Pseudomonas*, *Clostridium*, and *Streptococcus* clustering with or related to clinically relevant lineages, such as *Clostridioides difficile* and *Streptococcus pneumoniae*.

Conclusions: Bat fecal bacterial communities are highly diverse and vary significantly under different ecological and biological conditions. Detection of bacterial genera containing clinically relevant species and the phylogenetic proximity of selected amplicon sequence variants to known pathogens support further genomic, functional, and epidemiological studies to clarify their pathogenic and zoonotic relevance.

KEYWORDS: Bat gut microbiome; Bacterial diversity; Ecological drivers; Zoonotic potential; Chiroptera; Gut microbiota; 16S rRNA sequencing; Bacterial pathogens; One Health

✉To whom correspondence may be addressed. E-mail: lucas_palheta@usp.br

This is an open access article distributed under the terms of the Creative Commons Attribution-Non-Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

For reprints contact: reprints@medknow.com

©2026 Asian Pacific Journal of Tropical Biomedicine Produced by Wolters Kluwer-Medknow.

How to cite this article: Gomes LP, Morraye RA, Silva LS, Ximenez JPB, Martins MP, Silva Jr WA, et al. Ecological and biological determinants of fecal bacterial diversity in bats across four continents and phylogenetic screening for pathogen-affiliated bacterial lineages. Asian Pac J Trop Biomed 2026; 16(12): 495–507.

Article history: Received 10 June 2026; Revision 18 July 2026; Accepted 13 October 2026; Available online 22 October 2026

Summary

Question: How do ecological and biological factors influence the diversity of the bat fecal bacteriome and what is its association with bacterial lineages of zoonotic potential?

Findings: The reanalysis of 194 fecal samples from 20 bat species using 16S rRNA gene sequencing showed that diet, habitat, sex, and geographic region significantly influenced composition and diversity of bacterial community. Sixty-two potentially pathogenic genera were identified, and phylogenetic analyses revealed amplicon sequence variants related to clinically relevant species, including lineages associated with *Clostridioides* and *Streptococcus*.

Meaning: These findings indicate that bats harbor ecologically structured bacteriomes with potential zoonotic and epidemiological relevance.

1. Introduction

Bats (order Chiroptera) are widely recognized as natural reservoir hosts for a broad range of emerging pathogens that pose significant threats to both veterinary and human health. Several high-impact viral diseases, including Ebola virus, Marburg virus, Nipah virus, Hendra virus, severe acute respiratory syndrome coronavirus (SARS-CoV), and SARS-CoV-2, have been linked to bats as their primary reservoirs and probable evolutionary origins[1]. Despite harboring an exceptional diversity of viruses, bats generally remain asymptomatic upon infection with many pathogens that cause severe disease in other mammals. In contrast, their role in bacterial pathogen transmission remains comparatively underexplored[1,2]. The lack of overt inflammatory responses in bats is hypothesized to reflect their unique evolutionary trajectory, which has facilitated long-term co-evolution with a wide array of pathogens[3].

Although research has predominantly focused on bat-borne viruses, bats also host rich and diverse bacterial communities with potential relevance to public health[4]. Metagenomic studies of bat feces and urine samples have identified several zoonotic and potentially pathogenic bacterial genera, such as *Bartonella*, *Clostridium*, *Corynebacterium*, and *Rickettsia*[5,6]. These findings indicate that the impacts of bacterial microbiota on bat health and disease dynamics remain poorly understood. Previous studies have reported complex and diverse bacterial communities across different bat species, reflecting variation in host dietary habits and habitat preferences[6]. Moreover, unique biological and ecological traits of bats, such as powered flight, migratory behavior, long lifespan, high sociality, and adaptation to diverse ecological niches, may drive bacterial genetic variation and shape microbial community structure[6,7]. Collectively, these features may facilitate cross-species microbial transmission and enhance pathogen dispersal[8].

Given that host ecology and diet are major determinants of microbiota variation, the present study evaluated 16S rRNA

sequencing data derived from fecal samples of bat populations covering different geographic regions, sexes, and ecological habits. We assessed bacterial richness and diversity using 16S rRNA gene sequences and examined the phylogenetic proximity of detected strains to clinically relevant bacteria. Our analyses expand upon previous studies conducted using the same sample set, providing additional insights into bat-associated microbiomes[6,9,10]. Furthermore, our findings reveal a complex bacterial assemblage with intrinsic pathogenic potential, which merits epidemiological concerns. Aerial mobility, a defining characteristic of bats, may play a critical role in the spread of zoonotic diseases and the emergence of infectious-disease outbreaks.

2. Methods

2.1. Samples, species and public data collection

We conducted a comprehensive survey of publicly available datasets of bat fecal microbiome, focusing on 16S rRNA sequencing data. Following data quality assessment and strict screening based on our research objectives, a total of 194 bat fecal samples were selected, including 87 samples from the Dataverse Academic Portal[6], 96 samples from the NCBI SRA database (project accession number PRJEB47836)[9], and 11 samples from the Dryad Digital Repository[10]. The final dataset comprised specimens from 20 bat species: *Barbastella barbastellus*, *Miniopterus schreibersii*, *Myotis bechsteinii*, *Myotis emarginatus*, *Myotis myotis*, *Myotis nattererii*, *Myotis capaccinii*, *Myotis daubentonii*, *Myotis vivesi*, *Myotis blythii*, *Myotis ricketti* (= *Myotis pilosus*), *Rhinolophus hipposideros*, *Rhinolophus blasii*, *Rhinolophus ferrumequinum*, *Pipistrellus pipistrellus*, *Eptesicus bottae*, *Hypsugo ariel*, *Noctilio leporinus*, *Pipistrellus kuhlii*, and *Rousettus aegyptiacus*. Samples were obtained from bats captured across diverse geographic regions, including South America (two species), Europe (14 species), Asia (three species), and Africa (one species), and were stratified according to sex- and diet-based classification. Foraging behavior was inferred from capture habitats (Supplementary Table 1)[6,9,10]. Four piscivorous bat species originating from well-characterized fishing colonies distributed across four geographically distinct regions were included; notably, fishing behavior is thought to have evolved independently within these lineages[9]. In addition, facultative fishing species exhibiting both insectivorous and piscivorous feeding strategies were classified as piscivores to ensure consistency across analytical groupings.

2.2. 16S raw data processing

The sequences were organized into R1 (forward) and R2 (reverse) files and subjected to quality validation. Low-quality bases,

particularly those located at the read terminals, were trimmed, and sequence error rates were assessed through data filtering and sorting. All sequences were processed independently according to dietary grouping. R (v4.1.2) and the DADA2 package (v1.16.1)[11] were used to process and analyze the 16S rRNA sequences *via* a custom analytical pipeline. Forward and reverse reads were merged to obtain unique amplicon sequence variants (ASVs). Chimera detection and removal were performed during ASV table construction. After completion of all quality-control procedures, a total of 3 778 110 sequences were retained, with an average of 19 677 reads per sample.

Taxonomic annotation was performed against the SILVA database V138.1[12], with classification assigned down to the genus level. The ASV count table with 194 bat feces samples, sample metadata, and the taxonomic assignment table were imported into the Microbiome Analyst[13] for data processing and tabular output, including identification of microorganisms and analysis of diversity profiles. Additionally, diversity analyses were performed based on the relative abundance of each ASV (calculated as the read count of a given ASV divided by the total read depth per sample library). For statistical analyses, heat tree analysis was used to visualize the hierarchical taxonomic structure using median abundance, and the non-parametric Wilcoxon rank-sum test ($\alpha=0.05$)[14] was used to detect taxonomic differences between microbial communities. Separate workflows[15] were applied to compute and visualize alpha- and beta-diversity indices, respectively. One-way analysis of variance (ANOVA) ($\alpha=0.05$) and permutational multivariate analysis of variance (PERMANOVA) ($\alpha=0.05$) were also used for single-factor statistical comparisons and covariate-adjusted linear regression analyses.

2.3. Phylogenetic analysis of bacterial genera

To investigate the pathogen-related association of the bacterial taxa detected, phylogenetic analyses were conducted. Four genera containing well-documented pathogenic species and exhibiting high relative abundance in bat fecal samples were selected: *Pseudomonas*, *Staphylococcus*, *Clostridium*, and *Streptococcus*. For each genus, all ASVs targeting the V4 hypervariable region of the 16S rRNA gene were retrieved. A reference database was constructed for each genus, comprising confirmed and potential pathogenic bacterial species capable of infecting humans, animals, or plants. Full-length 16S rRNA sequences from reference taxa were aligned against the ASV dataset using MAFFT version 7[16], and resulting multiple sequence alignments were manually curated using BioEdit version 7.2.5[17]. Maximum likelihood phylogenetic trees were subsequently reconstructed in MEGA7 version 7.0.26[18] based on combined datasets containing ASVs (V4 region) and reference sequences, with branch robustness assessed by bootstrap analysis based on 1 000

replicates. Final phylogenetic trees were visualized and annotated using iTOL[19].

2.4. Statistical analysis

Statistical analyses were performed to assess differences in microbial composition and diversity among the evaluated groups. Relative abundance was calculated for each ASV as the number of reads assigned to a given ASV divided by the total number of reads in the corresponding sample. Differences in the relative abundance of bacterial taxa between groups were assessed using the non-parametric Wilcoxon rank-sum test. Alpha-diversity metrics were compared using ANOVA, whereas differences in overall microbial community composition based on beta-diversity were evaluated using PERMANOVA. Multivariable linear regression analyses were additionally performed to assess associations while accounting for potential confounding factors. All available covariates included in the sample metadata, including sex, diet, geographic region, and foraging behavior, were considered in the multivariable analyses. Heat tree analyses were used to visualize differences in the hierarchical taxonomic structure of microbial communities based on median taxon abundance. Statistical analyses and diversity assessments were conducted using MicrobiomeAnalyst and the corresponding R-based workflows described above. Statistical significance was defined as $P<0.05$.

3. Results

3.1. Bacterial profile of the fecal microbiota of bats

Analysis of the fecal microbiome of 20 bat species based on 16S rRNA sequencing data revealed a bacterial community dominated by two major phyla, Proteobacteria (57%) and Firmicutes (40%), which together accounted for 97% of the total identified taxa (Figure 1A). Across all samples, 123 bacterial families and 203 genera were detected (Supplementary Tables 1 and 2). In addition to these dominant phyla, a minor fraction of microbial diversity was observed at low relative abundances ($<1\%$), including Actinobacteriota, Bacteroidota, and several low-abundance phyla, such as Deinococcota, Spirochaetota, Campylobacterota, Patescibacteria, Verrucomicrobiota, Fusobacteriota, Desulfobacterota, Halobacterota, Chloroflexi, Cyanobacteria, Thermoplasmatota, and Synergistota. A subset of sequencing reads remained taxonomically unassigned, indicating the presence of poorly characterized or potentially novel microbial taxa. At the family level, Peptostreptococcaceae predominated in samples from the piscivorous species *Myotis vivesi* and *Myotis ricketti*. The highest family-level bacterial diversity was

observed in the frugivorous bat *Rousettus aegyptiacus* sampled in South Africa. In contrast, species such as *Rhinolophus hipposideros* (arthropod-feeding; forested habitats in Europe), *Rhinolophus blasii* (insectivorous; freshwater environments), *Hypsugo ariel* (insectivorous; desert habitats), and *Noctilio leporinus* (facultative piscivore–insectivore; predominantly whitewater habitats) exhibited high bacterial abundance but relatively low diversity at the family level. Rhodanobacteraceae and Thermoactinomycetaceae were the rarest families detected, each accounting for less than 1% of the processed dataset. At the genus level, the most prevalent taxa were *Pseudomonas*, *Staphylococcus*, *Carnobacterium*, and *Lonsdalea*. Consistent with the family-level patterns, *Rousettus aegyptiacus* showed the greatest genus-level diversity, while high relative abundances were also observed in the insectivorous species *Myotis myotis*, *Myotis bechsteini*, and *Barbastella barbastellus*. Analysis of the core microbiome revealed that no taxon was consistently present across all samples at a relative abundance threshold of $\geq 1\%$.

Relative abundance patterns were further examined by stratifying samples according to biological and ecological factors, including diet, habitat, and sex. At the genus level, insectivorous bats exhibited the highest microbial diversity (165 genera), with *Pseudomonas* representing the most abundant taxon (781 085 sequences) (Figure

1B). Frugivorous bats displayed a distinct microbial composition, also comprising 165 genera, with *Streptococcus* as the dominant genus (112 139 sequences) (Figure 1C). In contrast, piscivorous bats harbored the lowest bacterial diversity (16 genera), with *Clostridium* predominating (20 859 sequences) (Figure 1D). Habitat exerted a strong influence on microbial community structure. Bats foraging in forests and open fields exhibited the highest relative diversity (204 genera each), with *Pseudomonas* (502 480 sequences) and *Acinetobacter* (14 141 sequences) as the most abundant genera, respectively (Figure 2A and 2B). Conversely, bats captured in caves and desert habitats showed distinct microbial profiles, characterized by a predominance of *Staphylococcus* (298 808 sequences) in cave-dwelling bats (Figure 2C) and *Romboutsia* (1 517 sequences) in desert-dwelling bats (Figure 2D). Among piscivorous bats, individuals foraging in freshwater and coastal water environments exhibited greater *Bartonella* enrichment (19 220 sequences and 6 434 sequences, respectively) (Figure 2E and 2F). Sex-stratified comparisons further revealed distinct microbial signatures, with male bats showing enrichment of *Staphylococcus* (338 648 sequences) (Figure 3A) and female bats dominated by *Pseudomonas* (168 689 sequences) (Figure 3B).




Figure 1. Relative abundance of the predominant bacterial genera from 20 bat species with distinct dietary habits. (A) All samples; (B) Insectivorous bats; (C) Frugivorous bats; (D) Piscivorous bats.




Figure 2. Relative abundance of the predominant bacterial genera from 20 bat species grouped according to habitat. (A) Forest-associated bats; (B) Open-area bats; (C) Cave-dwelling bats; (D) Desert-dwelling bats; (E) Freshwater piscivorous bats; (F) Coastal piscivorous bats.




Figure 3. Taxonomic composition of the bacterial microbiota identified by 16S rRNA gene sequencing from 20 bat species grouped according to host sex. (A) Male bats; (B) Female bats.




Figure 4. Heat tree analysis of differentially abundant bacterial taxa identified by 16S rRNA gene sequencing in the fecal microbiota of 20 bat species. (A) Differential bacterial taxa between insectivorous and piscivorous bats; (B) Differential bacterial taxa between piscivorous and frugivorous bats; (C) Differential bacterial taxa between insectivorous and frugivorous bats; (D) Habitat-associated differences between forest-dwelling and desert-dwelling bats; (E) Habitat-specific bacterial taxa in freshwater and coastal piscivorous bats.

3.2. Phylogenetic analysis of bat fecal microbiota communities

Heat-gradient phylogenetic tree analyses were conducted to compare divergent fecal microbiota abundance patterns among the 20 bat species (Supplementary Table 3). Comparison between insectivorous and piscivorous bats revealed significant microbial divergence (Figure 4; $P=1.28 \times 10^{-4}$). Insectivorous bats were dominated by Proteobacteria, particularly Gammaproteobacteria, with marked enrichment of Pseudomonadales and Moraxellaceae. In contrast, piscivorous bats were enriched in Firmicutes, mainly

Bacilli and Staphylococcaceae, showing reduced overall diversity but dominance of specific bacterial taxa.

Comparison between piscivorous and frugivorous bats showed marginal microbial community differentiation (Figure 4; $P=0.05$). Frugivorous bats displayed enrichment in Proteobacteria, including *Pseudomonas* and other members of Pseudomonadales, corresponding to higher microbial diversity and community complexity. Piscivorous bats again showed enrichment in Firmicutes, with predominance of *Staphylococcus* and other Bacillales, reflecting a more restricted gut microbiota.

Significant taxonomic differences were also detected between




Figure 5. Alpha diversity of the fecal microbiota identified by 16S rRNA gene sequencing in 20 bat species, estimated using the Chao1 richness index. (A) Alpha diversity across individual bat species (ANOVA, $P < 0.001$); (B) Comparison of alpha diversity according to dietary guild; (C) Comparison of alpha diversity according to habitat type.




Figure 6. Beta diversity of the fecal microbiota identified by 16S rRNA gene sequencing from 20 bat species, visualized by principal coordinates analysis (PCoA). (A) Ordination of samples according to bat species; (B) Beta diversity grouped by habitat type (caves, desert, forests, freshwater, meadow, and ocean coasts); (C) Beta diversity grouped by dietary guild; (D) Beta diversity according to climatic region (semiarid, temperate, and tropical).

insectivorous and frugivorous bats (Figure 4; $P=0.007$). Both groups exhibited relatively high microbial diversity but distinct taxonomic signatures: insectivorous bats were enriched in Proteobacteria, particularly Gammaproteobacteria and Moraxellaceae, whereas frugivorous bats were dominated by Firmicutes, especially Lactobacillales, Staphylococcaceae, and *Streptococcus*. These patterns indicate that host gut microbial community composition is shaped by diet-driven selective pressures, with insectivorous bats adapted to protein- and chitin-rich diets and frugivorous bats harboring taxa specialized in the metabolism of fermentable carbohydrates.

Habitat-driven differences were also evident. Comparison between forest-dwelling and desert bats yielded significant microbial differences (Figure 4; $P=0.002$). Forest-dwelling bats exhibited greater bacterial diversity, with enrichment of Proteobacteria (particularly Gammaproteobacteria, Pseudomonadales, Pseudomonadaceae, and *Acinetobacter*) and Bacilli (including Lactobacillales and Staphylococcaceae). In contrast, desert bats showed markedly reduced diversity and were dominated by Romboutsia, reflecting adaptive microbial shifts in response to nutritionally limited diets in extreme environments.

Finally, significant microbial divergence was observed between freshwater and coastal piscivorous bats (Figure 4; $P=0.006$). Freshwater bats were enriched in *Bartonella* (identified in the dataset although not highlighted in the figure), whereas coastal bats were dominated by Romboutsia, suggesting that aquatic foraging ecology directly influences gut microbiota composition by selecting for taxa adapted to distinct dietary resources.

3.3. Diversity profile of bacterial communities in the fecal microbiota of bats

To investigate differences in bacterial diversity of bat fecal microbiota, analyses were conducted focusing on patterns of alpha and beta diversity. Alpha diversity analysis revealed substantial variations in microbial richness across bat species, with distinct taxa exhibiting divergent diversity levels. Alpha diversity indices among bat species differed significantly ($P<0.001$), with *Rousettus aegyptiacus* exhibiting the highest microbial richness (Figure 5A). When samples were grouped by diet (frugivorous, insectivorous, and piscivorous), significant differences were also observed, with frugivorous bats showing the highest alpha diversity values ($P<0.001$) and piscivorous bats the lowest (Figure 5B). Habitat-based analyses further demonstrated that bats inhabiting caves and forests possessed significantly greater fecal microbial diversity ($P<0.001$), whereas bats from freshwater and coastal environments hosted less diverse and structurally simpler bacterial communities (Figure 5C).

Beta diversity analyses were performed to assess dissimilarity in microbial composition among samples. Microbial communities revealed partial clustering according to host species (Figure 6A) and habitat (Figure 6B). However, overlapping clusters indicated concurrent patterns of host or habitat specificity and microbiota sharing across conditions. PERMANOVA results confirmed significant overall compositional divergence among bat species ($F=4.667$, $R^2=0.340$, $P=0.001$). It demonstrated significant




Figure 7. Distribution and relative abundance of potentially pathogenic bacterial genera identified by 16S rRNA gene sequencing in the fecal microbiota of 20 bat species. Boxplots show the log-transformed bacterial counts of (A) *Streptococcus*, (B) *Mycobacterium*, (C) *Escherichia-Shigella*, and (D) *Yersinia* across bat species.




Figure 8. Single-factor analysis of bacterial genera across dietary guilds (A-F) and habitat types (G-H).

differences between species with distinct dietary habits, such as insectivorous *Myotis daubentonii* and frugivorous *Rousettus aegyptiacus* ($P=0.001$; $FDR=0.007$). Although several bacterial communities overlapped across forest, grassland, desert, and cave habitats, a distinct microbial assemblage was identified as exclusive to piscivorous bats ($P=0.001$; $FDR=0.008$). Diet-driven microbial differentiation was further validated by PERMANOVA analysis ($F=20.34$, $R^2=0.177$, $P=0.001$). Core bacterial taxa were common to all three dietary groups, alongside a subset of genera exclusively associated with insectivorous bat hosts (Figure 6C). Finally, PERMANOVA analysis incorporating climatic variables detected weak but significant climatic effects on microbial community structure ($F=1.119$, $R^2=0.012$, $P=0.034$). A shared core microbiota was conserved across climatic regions; nevertheless, temperate environments harbored a distinct bacterial assemblage ($P=0.004$; $FDR=0.012$) (Figure 6D).

3.4. Potentially pathogenic genera in bat fecal microbiota

Genera containing pathogenic bacterial members with relatively high abundance were evaluated in relation to environmental factors, bat species, diet, and habitat. Interspecific comparison revealed a total of 62 bacterial genera. Samples obtained from the frugivorous bat *Rousettus aegyptiacus* harbored a higher diversity of potentially pathogenic taxa. This species exhibited significant enrichment of the genera *Streptococcus* ($P<0.001$), *Mycobacterium* ($P<0.001$), and *Shinella* ($P<0.01$) (Figure 7A-C). In contrast, other potentially pathogenic genera, such as *Yersinia*, were predominantly detected in insectivorous bats, including *Barbastella barbastellus*, *Myotis bechsteinii*, *Myotis nattereri*, and *Pipistrellus pipistrellus* ($P<0.001$)

(Figure 7D), indicating occurrence of host-specific patterns of these pathogenic bacteria.

When examining differences across dietary guilds, univariable statistical analyses revealed 118 bacterial genera with statistically significant differences in relative abundance among feeding groups. Enrichment was primarily observed in frugivorous bats, particularly for *Streptococcus* ($P<0.001$), *Mycobacterium* ($P<0.001$), *Shinella* ($P<0.001$), and *Helicobacter* ($P<0.05$) (Figure 8A-D). Conversely, insectivorous bats exhibited marked enrichment of *Pseudomonas* ($P<0.05$) and *Staphylococcus* ($P<0.001$) (Figure 8E and F), indicating the influence of dietary specialization on the structure of potentially pathogenic bacterial communities.

Habitat-related patterns further supported the role of environmental context in shaping the distribution of these genera. The genus *Pseudomonas* ($P<0.001$) was significantly enriched in samples from cave and forest environments (Figure 8G), suggesting an association with more complex or sheltered habitats. In contrast, *Yersinia* ($P<0.05$) predominated in samples obtained from open grassland habitats (Figure 8H), highlighting the contribution of habitat-specific ecological pressures to the composition and abundance of potentially pathogenic bacteria in bat fecal microbiota.

3.5. Phylogenetic relationship of the main potentially pathogenic bacteria in the bat microbiota

Phylogenetic analyses based on 16S rRNA sequencing of bat fecal samples revealed a high number of ASVs, reflecting substantial bacterial diversity, particularly within the genera *Pseudomonas* (Supplementary Figure 1) and *Staphylococcus* (Supplementary Figure 2). Although *Staphylococcus* exhibited a high ASV richness, most

sequences did not cluster closely with the pathogenic reference strains included in the analysis, suggesting a predominance of non-pathogenic or distantly related lineages. In contrast, *Pseudomonas*-derived ASVs showed closer phylogenetic relationships to several disease-associated species. In the *Clostridium* phylogeny (Supplementary Figure 3 and Supplementary Table 4), three distinct ASVs clustered basally, appearing to represent ancestral lineages shared by known pathogenic *Clostridium* species. The majority of *Clostridium* ASVs formed a separate, well-supported clade, displaying a more distant relationship to established pathogens while sharing a common ancestor. Similarly, the phylogenetic reconstruction for *Streptococcus* (Supplementary Figure 4) revealed a closer association with multiple pathogenic species. Collectively, these findings for *Clostridium*, *Pseudomonas*, and *Streptococcus* indicate the presence of ASVs that may represent ancestral lineages of clinically relevant pathogenic bacteria within the bat fecal microbiota.

4. Discussion

Bats are widely recognized as natural reservoirs of viruses, and in recent years, increasingly comprehensive studies of bat-associated microbiomes have suggested that these mammals also host a wide diversity of bacteria, including potentially pathogenic strains[20]. Benefiting from flight capacity and migratory behavior, bats have successfully colonized a broad range of terrestrial and aquatic-associated habitats, ranging from forests, deserts, and urban environments to freshwater systems, coastal zones, and oceanic regions. Although bats harbor a remarkable diversity of pathogenic microorganisms, their long-distance mobility may facilitate the dispersal of infectious agents, including bacteria associated with human disease[6]. It has been proposed that the elevated metabolic rates and increased body temperatures required for flight have shaped bat immune systems, creating selective pressures that reduce microbial virulence while allowing bats to tolerate and carry a wide range of pathogens[21]. Moreover, adaptation to diverse dietary niches and variable ecosystem distributions has driven the evolution of complex and heterogeneous bacterial communities in bats[22]. In this study, we evaluated the complex interactions among sex, dietary preference, and geographic origin, assessing patterns of bacterial diversity in bats and exploring their reservoir potential for pathogenic bacteria.

Across all bat samples analyzed, members of the phyla Firmicutes (notably the families Staphylococcaceae and Carnobacteriaceae) and Proteobacteria (particularly Pseudomonadaceae and Pectobacteriaceae) dominated the fecal microbiota, consistent with previous studies of bat-associated bacterial communities[23]. Geographic variation was evident at the genus level, with

Pseudomonas predominating at terrestrial sampling locations, whereas *Bartonella* and *Romboutsia* were more abundant in samples from freshwater and marine environments. Insectivorous bats exhibited a higher relative abundance of *Pseudomonas*, a pattern similar to that observed in bats foraging in wet habitats. This pattern differed from that of bats consuming a broader range of arthropods other than insects. Notably, *Pseudomonas* was more prevalent in female bats than in males, although it was abundant in both sexes. The genus *Paraclostridium*, significantly enriched in piscivorous bats and previously described as characteristic of piscivorous bats when compared with arthropodophagous bats[9], was also detected in insectivorous bats in our dataset. This finding indicates that no strictly diet-specific gut microbiota exists in these bat hosts.

From an ecological perspective, fruit bats interact extensively with other organisms, including plants and animals, due to their feeding behavior and habitat use. This complex interaction network likely contributes to the higher and more distinct diversity of bacterial genera observed in the intestinal microbiomes of frugivorous bats compared with other dietary guilds. While zoonotic and potentially zoonotic pathogens of bats have been widely investigated, the broader ecological context of bat-associated pathogens remains poorly understood, particularly the diverse community of taxa that bats interact with[24,25]. Fruits themselves constitute complex microecosystems supporting diverse arthropod assemblages that often co-evolve with host plants[26]. In this context, fruit bats may ingest insects incidentally, which can constitute a meaningful nutritional resource[27–29]. These trophic interactions may facilitate the acquisition of viruses and bacteria associated with fruit-dwelling arthropods[30].

Alpha diversity analyses at both family and genus levels revealed that the frugivorous species *Rousettus aegyptiacus* from South Africa, a bat typically inhabiting terrestrial environments, harbored the most diverse bacterial communities. These findings were supported by both capture location and dietary preference. In contrast, bats with arthropod-based diets exhibited higher beta diversity indices and are recognized as major nocturnal consumers, playing key roles in regulating arthropod populations[31]. Their broad consumption of diverse arthropod taxa mirrors spatial variation in arthropod availability and abundance across habitats[32,33]. Such dietary breadth likely contributes to the elevated and differentiated abundance of bacterial genera observed in their microbiomes. For example, *Diplorickettsia*, a potential human pathogen previously isolated from Ixodes ticks, important vectors of human disease[34], was detected at high abundance in arthropod-feeding bats.

Arthropodophagous bats constituted the majority of dietary categories in our dataset. Terrestrial bat samples from Slovenia showed a distinct relative abundance of bacterial genera, including *Streptococcus*, *Mycobacterium*, *Pseudomonas*, *Mycoplasma*, and *Ureaplasma*, which were largely restricted to continental bat

populations. In addition, many bats frequently inhabit urban environments and often come into close contact with humans and domestic animals. The accumulation of bat guano in buildings potentially facilitates the maintenance and indirect transmission of these bacteria to humans[35,36]. Similar patterns were observed in bats inhabiting coastal and freshwater environments. Piscivorous bats foraging near oceans, lakes, and rivers exhibited high abundances of *Aeromonas*, a genus commonly found in freshwater and marine environments and associated with gastrointestinal disease, wound infections, and septicemia in humans[37].

The fecal bacteriome of bats revealed high diversity within the genera *Pseudomonas* and *Staphylococcus*, with numerous ASVs diverging from known pathogenic strains. In the case of *Pseudomonas*, a limited number of ASVs clustered near pathogenic species, whereas *Staphylococcus* ASVs were largely phylogenetically distant from established pathogens, despite their high overall diversity. Bats are known to harbor non-viral pathogens while remaining clinically asymptomatic, which supports their role as natural reservoirs of diverse bacterial taxa. Nevertheless, the magnitude of their contribution to disease spillover and zoonotic bacterial transmission remains insufficiently explored[5]. Most *Pseudomonas* ASVs occupied basal phylogenetic positions relative to reference pathogenic strains, with sequences such as *Pseudomonas*_467, *Pseudomonas*_485, and *Pseudomonas*_473 clustering closer to the outgroup than to pathogenic ingroups. These findings suggest that diversification of these clades may have co-occurred alongside their mammalian hosts. Several ASVs also clustered with pathogenic species infecting fungi (e.g., *Pseudomonas tolaasi*) and warm-blooded animals (e.g., *Pseudomonas oryziatrans*), as well as with putative ancestral lineages related to *Pseudomonas aeruginosa*[38].

Similarly, phylogenetic reconstructions of *Clostridium* and *Streptococcus* revealed basal lineages related to pathogenic taxa, including putative ancestral strains associated with *Clostridium difficile* (*Clostridium* 014) and *Streptococcus pneumoniae* (*Streptococcus* 012 and *Streptococcus* 034), both of which carry well-characterized virulence factors. These phylogenetic patterns support prediction from the “bat seeding hypothesis”; however, the present 16S rRNA-based analysis does not establish bats as ancestral sources of these pathogenic lineages. Further genomic analyses are required to disentangle their evolutionary histories and assess their potential epidemiological significance[39]. This long-term cohabitation and coevolution between bats and microbial symbionts may explain the capacity of bats to host infectious agents with little or no overt clinical signs[5,40]. Combined with the long lifespan of many bat species and their adaptability to diverse ecological niches, these traits underscore bats’ potential contribution to the emergence and transmission of zoonotic pathogens[41].

The high immune tolerance of bats is thought to be a consequence

of long-term coevolution with microbial pathogens, which has shaped distinctive immunological response profiles[42]. The bat immune system appears to limit immunopathology while maintaining effective pathogen control. Microbial community composition may therefore serve as an important predictor of immune response magnitude and infection severity[43]. Robust antiviral defenses in bats include restrained inflammatory cytokine expression, activation of interferon regulatory factors, and engagement of Toll-like receptor signaling pathways[41,44]. Additionally, bats exhibit reduced sensing of self and exogenous DNA, partly due to dampened expression of stimulator of interferon genes, a key adaptor protein in DNA-sensing pathways. This reduced functionality of stimulator of interferon genes may contribute to balanced immune responses, highlighting a fundamental distinction in immune tolerance between bats and other mammals[45].

In summary, our study shows that bats harbor complex fecal bacterial communities, whose composition and diversity vary significantly according to diet, habitat, sex, and geographic origin. Proteobacteria and Firmicutes predominated across all analyzed samples, with recurrent detection of genera such as *Pseudomonas*, *Staphylococcus*, *Clostridium*, and *Streptococcus*, some of which include species of recognized clinical relevance. Frugivorous and insectivorous bats exhibited richer bacterial communities than piscivorous bats, supporting a significant association between dietary habits and microbial diversity. Similarly, cave- and forest-associated bats showed higher bacterial richness than bats from freshwater, coastal, and desert environments, indicating an association between habitat characteristics and fecal microbiota composition.

The absence of a universally shared core microbiome further highlights the variability of bat-associated bacterial communities across ecological and biological contexts. Phylogenetic analyses identified ASVs within *Pseudomonas*, *Clostridium*, and *Streptococcus* that clustered with or were phylogenetically related to bacterial taxa of recognized pathogenic potential, including lineages related to *Clostridioides difficile* and *Streptococcus pneumoniae*. However, these phylogenetic relationships do not confirm pathogenicity or zoonotic spillover risk. Rather, they indicate the presence of bacterial lineages that warrant further genomic, functional, and epidemiological investigation. Collectively, these findings expand current knowledge of bat-associated bacterial diversity and reinforce the need for continued investigation of how ecological factors contribute to the distribution of bacterial taxa with potential clinical relevance.

Conflict of interest statement

The authors declare that there is no conflict of interest.

Funding

This work was supported by the Fundação Amazônia de Amparo a Estudos e Pesquisas (FAPESPA, grant number 2014/15440-7); the São Paulo Research Foundation (FAPESP, grant number 13/08135-2); the National Council for Scientific and Technological Development (CNPq, grant numbers 421516/2025-9, PS); and the Coordination for the Improvement of Higher Education Personnel (CAPES, Finance Code 001).

Data availability statement

The data supporting the findings of this study are available from the corresponding author upon request.

Authors' contributions

LPG, WAS Jr., and MPCS contributed to the conception and design of the study, definition of the intellectual content, literature review, data collection and analysis, statistical analysis, preparation and editing of the manuscript, as well as to the critical review of the final version of the paper. RAM, LSS, JPXB, and MPM contributed to the literature review, data analysis, statistical analysis, preparation and editing of the manuscript, as well as the critical review of the manuscript.

References

- [1] Letko M, Seifert SN, Olival KJ, Plowright RK, Munster VJ. Bat-borne virus diversity, spillover and emergence. *Nat Rev Microbiol* 2020; **18**(8): 461-471.
- [2] Szentivanyi T, McKee C, Jones G, Foster JT. Trends in bacterial pathogens of bats: Global distribution and knowledge gaps. *Transbound Emerg Dis* 2023; **2023**. doi: 10.1155/2023/9285855.
- [3] Irving AT, Ahn M, Goh G, Anderson DE, Wang LF. Lessons from the host defences of bats, a unique viral reservoir. *Nature* 2021; **589**(7842): 363-370.
- [4] Dhivahar J, Parthasarathy A, Krishnan K, Kovi BS, Pandian GN. Bat-associated microbes: Opportunities and perils, an overview. *Heliyon* 2023; **9**(12). doi: 10.1016/j.heliyon.2023.e22351.
- [5] Veikkolainen V, Vesterinen EJ, Lilley TM, Pulliainen AT. Bats as reservoir hosts of human bacterial pathogen, *Bartonella mayotimonensis*. *Emerg Infect Dis* 2014; **20**(6): 960-967. doi: 10.3201/eid2006.130956.
- [6] Vengust M, Knapic T, Weese JS. The fecal bacterial microbiota of bats; Slovenia. *PLoS One* 2018; **13**(5). doi: 10.1371/journal.pone.0196728.
- [7] McKee CD, Bai Y, Webb CT, Kosoy MY. Bats are key hosts in the radiation of mammal-associated *Bartonella* bacteria. *Infect Genet Evol* 2021; **89**. doi: 10.1016/j.meegid.2021.104719.
- [8] Corduneanu A, Wu-Chuang A, Maitre A, Obregon D, Sándor AD, Cabezas-Cruz A. Structural differences in the gut microbiome of bats using terrestrial *vs.* aquatic feeding resources. *BMC Microbiol* 2023; **23**: 93. doi: 10.1186/s12866-023-02836-7.
- [9] Aizpurua O, Nyholm L, Morris E, Chaverri G, Herrera Montalvo LG, Flores-Martinez JJ, et al. The role of the gut microbiota in the dietary niche expansion of fishing bats. *Anim Microbiome* 2021; **3**(1): 76. doi: 10.1186/s42523-021-00137-w.
- [10] Dietrich M, Markotter W. Studying the microbiota of bats: Accuracy of direct and indirect samplings. *Ecol Evol* 2019; **9**(4): 1730-1735. doi: 10.1002/ece3.4842.
- [11] Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat Methods* 2016; **13**(7): 581-583. doi: 10.1038/nmeth.3869.
- [12] Quast C, Priesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Res* 2013; **41**: D590-D596. doi: 10.1093/nar/gks1219.
- [13] Dhariwal A, Chong J, Habib S, King IL, Agellon LB, Xia J. MicrobiomeAnalyst: A web-based tool for comprehensive statistical, visual and meta-analysis of microbiome data. *Nucleic Acids Res* 2017; **45**(W1): W180-W188. doi: 10.1093/nar/gkx295.
- [14] Foster ZSL, Sharpton TJ, Grünwald NJ. Metacoder: An R package for visualization and manipulation of community taxonomic diversity data. *PLoS Comput Biol* 2017; **13**(2). doi: 10.1371/journal.pcbi.1005404.
- [15] Menéndez ML, Pardo JA, Pardo L, Pardo MC. The Jensen-Shannon divergence. *J Franklin Inst* 1997; **334**(2): 307-318. doi: 10.1016/S0016-0032(96)00063-4.
- [16] Katoh K, Rozewicki J, Yamada KD. MAFFT online service: Multiple sequence alignment, interactive sequence choice and visualization. *Brief Bioinform* 2019; **20**(4): 1160-1166. doi: 10.1093/bib/bbx108.
- [17] Hall TA. BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 1999; **41**: 95-98.
- [18] Kumar S, Stecher G, Tamura K. MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol* 2016; **33**(7): 1870-1874. doi: 10.1093/molbev/msw054.
- [19] Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: An online tool for phylogenetic tree display and annotation. *Nucleic Acids Res* 2021; **49**(W1): W293-W296. doi: 10.1093/nar/gkab301.
- [20] Federici L, Masulli M, De Laurenzi V, Allocati N. An overview of bats microbiota and its implication in transmissible diseases. *Front Microbiol* 2022; **13**. doi: 10.3389/fmicb.2022.1012189.
- [21] Morales AE, Dong Y, Brown T, Baid K, Kontopoulou DG, Gonzalez V, et al. Bat genomes illuminate adaptations to viral tolerance and disease resistance. *Nature* 2025; **638**(8050): 449-458. doi: 10.1038/s41586-024-08471-0.

- [22] Dai W, Leng H, Li J, Li A, Li Z, Zhu Y, et al. The role of host traits and geography in shaping the gut microbiome of insectivorous bats. *mSphere* 2024; **9**(4). doi: 10.1128/msphere.00087-24.
- [23] Huang Y, Sun Y, Huang Q, Lv X, Pu J, Zhu W, et al. The threat of potentially pathogenic bacteria in the feces of bats. *Microbiol Spectr* 2022; **10**(6). doi: 10.1128/spectrum.01802-22.
- [24] Aziz SA, McConkey KR, Tanalgo K, Sritongchuay T, Low MR, Yong JY, et al. The critical importance of Old World fruit bats for healthy ecosystems and economies. *Front Ecol Evol* 2021; **9**. doi: 10.3389/fevo.2021.641411.
- [25] Santos RT, Flores RJD, Montecillo AD, De Leon MP. Mapping the bacterial diversity and functional landscape of the Philippine bat gut microbiome using a 16S-based metataxonomics approach. *SciEnggJ* 2025; **18**(2): 275-294. doi: 10.54645/2025182HWH-76.
- [26] Borges RM. Interactions between figs and gall-inducing fig wasps: Adaptations, constraints, and unanswered questions. *Front Ecol Evol* 2021; **9**. doi: 10.3389/fevo.2021.685542.
- [27] Pellón JJ, Medina-Espinoza EF, Lim BK, Cornejo F, Medellín RA. Eat what you can, when you can: Relatively high arthropod consumption by frugivorous bats in Amazonian Peru. *Mamm Biol* 2023; **103**(1): 137-144. doi: 10.1007/s42991-022-00327-2.
- [28] Ingala MR, Simmons NB, Wultsch C, Krampis K, Provost KL, Perkins SL. Molecular diet analysis of neotropical bats based on fecal DNA metabarcoding. *Ecol Evol* 2021; **11**(12): 7474-7491. doi: 10.1002/ece3.7579.
- [29] Adhikari BN, Paskey AC, Frey KG, Bennett AJ, Long KA, Kuhn JH, et al. Virome profiling of fig wasps (*Ceratosolen* spp.) reveals virus diversity spanning four realms. *Virology* 2024; **591**. doi: 10.1016/j.virol.2024.109992.
- [30] Dong Y, Zhang ZR, Mishra S, Wong ACN, Huang JF, Wang B, et al. Diversity and metabolic potentials of microbial communities associated with pollinator and cheater fig wasps in fig-fig wasp mutualism system. *Front Microbiol* 2022; **13**. doi: 10.3389/fmicb.2022.1009919.
- [31] Tuneu-Corral C, Puig-Montserrat X, Riba-Bertolin D, Russo D, Rebelo H, Cabeza M, et al. Pest suppression by bats and management strategies to favour it: A global review. *Biol Rev Camb Philos Soc* 2023; **98**(5): 1564-1582. doi: 10.1111/brev.12967.
- [32] Zhu D, Liu Y, Gong L, Si M, Wang Q, Feng J, et al. The consumption and diversity variation responses of agricultural pests and their dietary niche differentiation in insectivorous bats. *Animals (Basel)* 2024; **14**(5). doi: 10.3390/ani14050815.
- [33] Aihartzia J, Vallejo N, Aldasoro M, García-Mudarra JL, Goiti U, Noguera J, et al. Aerospace-foraging bats eat seasonably across varying habitats. *Sci Rep* 2023; **13**: 19576. doi: 10.1038/s41598-023-46939-7.
- [34] Merenstein C, Ward J, Allen D. *Diplorickettsia* bacteria in an *Ixodes scapularis* tick, Vermont, USA. *Emerg Infect Dis* 2020; **26**(5): 1036-1038. doi: 10.3201/eid2605.191135.
- [35] Gerbáčová K, Maliničová L, Kisková J, Maslišová V, Uhrin M, Pristaš P. The faecal microbiome of building-dwelling insectivorous bats (*Myotis myotis* and *Rhinolophus hipposideros*) also contains antibiotic-resistant bacterial representatives. *Curr Microbiol* 2020; **77**(9): 2333-2344. doi: 10.1007/s00284-020-02095-z.
- [36] Jackson RT, Lunn TJ, DeAnglis IK, Ogola JG, Webala PW, Forbes KM. Frequent and intense human-bat interactions occur in buildings of rural Kenya. *PLoS Negl Trop Dis* 2024; **18**(2). doi: 10.1371/journal.pntd.0011988.
- [37] Aizpurua O, Nyholm L, Morris E, Chaverri G, Herrera Montalvo LG, Flores-Martinez JJ, et al. The role of the gut microbiota in the dietary niche expansion of fishing bats. *Anim Microbiome* 2021; **3**(1): 76. doi: 10.1186/s42523-021-00137-w.
- [38] Liu Z, Zhao Y, Sossah FL, Okorley BA, Amoako DG, Liu P, et al. Characterization, pathogenicity, phylogeny, and comparative genomic analysis of *Pseudomonas tolaasii* strains isolated from various mushrooms in China. *Phytopathology* 2022; **112**(3): 521-534. doi: 10.1094/PHYTO-12-20-0550-R.
- [39] Savage PM, Savage TJ, Kruse DL, Stone PA. *Leclercia adecarboxylata* and *Pseudomonas oryzihabitans* infection after Achilles tendon repair: A case report and literature review. *J Foot Ankle Surg* 2023; **62**(4): 742-745. doi: 10.1053/j.jfas.2022.11.004.
- [40] Quiroz-Morales SE, García-Reyes S, Ponce-Soto GY, Servín-González L, Soberón-Chávez G. Tracking the origins of *Pseudomonas aeruginosa* phylogroups by diversity and evolutionary analysis of important pathogenic marker genes. *Diversity* 2022; **14**(5). doi: 10.3390/d14050345.
- [41] Nabi G, Wang Y, Lü L, Jiang C, Ahmad S, Wu Y, et al. Bats and birds as viral reservoirs: A physiological and ecological perspective. *Sci Total Environ* 2021; **754**. doi: 10.1016/j.scitotenv.2020.142372.
- [42] Das S, Jain D, Chaudhary P, Quintela-Tizon RM, Banerjee A, Kesavardhana S. Bat adaptations in inflammation and cell death regulation contribute to viral tolerance. *mBio* 2025; **16**(3). doi: 10.1128/mbio.03204-23.
- [43] Berman TS, Weinberg M, Moreno KR, Czirják GÁ, Yovel Y. In sickness and in health: The dynamics of the fruit bat gut microbiota under a bacterial antigen challenge and its association with the immune response. *Front Immunol* 2023; **14**. doi: 10.3389/fimmu.2023.1152107.
- [44] Clayton E, Munir M. Fundamental characteristics of bat interferon systems. *Front Cell Infect Microbiol* 2020; **10**. doi: 10.3389/fcimb.2020.527921.
- [45] Irving AT, Ahn M, Goh G, Anderson DE, Wang LF. Lessons from the host defences of bats, a unique viral reservoir. *Nature* 2021; **589**(7842): 363-370. doi: 10.1038/s41586-020-03128-0.

Publisher's note

The Publisher of the *Journal* remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Supplementary Table 1. List of bacterial families found in fecal microbiota analyses of bats.

	Barbastella_barb	Eptesicus_bottae	Hypsugo_ariel	Miniopterus_schr	Myotis_bechsteiri	Myotis_blythii	Myotis_capaccinii	Myotis_daubentonii	Myotis_emarginatus	Myotis_myotis	Myotis_nattererii	Myotis_ricketti	Myotis_viviparus	Noctilio_leporinus	Pipistrellus_kuhlmanni	Pipistrellus_pipistrellus	Rhinolophus_blaesi	Rhinolophus_ferrugineus	Rhinolophus_hipposideros	Rousettus_aegyptiacus
Acetobacteraceae	0	0	0	192.537.255.959	0	0	254.544.083.264	751.410.982.845	248.714.296.587	491.674.449.797	0	0	0	0	0	0	0	0	1	0
Actinomycetaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Aerococcaceae	0	0	0	0	0	142.254.990.687	0	188.186.706.148	131.696.258.684	0	85.013.839.462.1	0	0	0	0	0	0	0	0	79.358.658.219.4
Aeromonadaceae	0	0	0	0	0	0	0	0	380.546.464.723	0	0	0	0	0	0	0	0	0	0	0
Akkermansiaceae	0	0	0	0	0	0	0	0	693.138.203.603	0	85.155.058.464.6	0	0	0	0	0	0	0	0	0
Alcaligenaceae	0	0	0	0	0	0	0	0	159.014.058.473	423.786.345.850	0	0	0	0	0	0	0	0	0	538.203.666.116
Alcanivoraceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	140.706.840.814
Aquaspirillaceae	0	0	0	0	0	0	0	0	36.967.370.858.6	310.639.505.939	0	0	0	0	0	0	511.591.118.219	0	0	0
Arcobacteraceae	0	0	0	0	0	0	0	0	250.980.263.076	0	0	0	0	0	0	0	0	0	0	0
Atopobiaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	102.012.459.590
Azospirillaceae	0	0	0	0	0	0	143.298.891.319	150.282.196.569	0	0	0	0	0	0	0	0	0	0	0	0
Bacillaceae	74.480.804.819.1	0	0	155.570.102.814	574.114.039.546	437.558.638.669	0	972.381.471.874	888.847.814.032	610.787.213.995	847.314.014.573	0	0	0	411.608.848.357	0	950.284.867.204	0	851.276.386.929	0
Balneolaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	261.011.189.711
Beijerinckiaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	260.659.422.809
Bifidobacteriaceae	0	0	0	0	49.995.417.088.7	506.472.048.554	0	0	6.39E+09	429.957.991.664	0	0	0	0	0	0	480.162.312.322	0	175.883.551.018	0
Bogoriellaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	306.037.378.772
Brevibacteriaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	33.523.404.824.1
Budviciaceae	0	0	0	0	0	0	0	0	7.88E+09	0	0	0	0	0	0	0	0	0	0	0
Burkholderiaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	524.132.982.035
Campylobacteraceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	263.825.326.527
Carnobacteriaceae	54.832.494.345.1	0	0	940.660.017.713	479.206.072.776	78.359.251.971.1	0	360.065.010.965	147.959.183.673	959.217.764.465	105.617.691.916	0	0	0	764.187.565.469	0	89.739.717.105.6	0	378.501.401.791	0
Caulobacteraceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	10.975.133.583.6
Cellulomonadaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	113.620.773.957
Chitinophagaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	291.966.694.690
Chlamydiaceae	137.081.235.863	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	334.367.573.853	0	0	0
Christensenellaceae	124.896.237.120	0	0	0	0	0	0	0	189.263.804.943	0	0	0	0	0	0	0	9.87E+09	0	0	0
Clostridiaceae	0	839.484.765.915	0	0	0	0	827.701.938.306	350.658.458.661	0	339.440.519.734	0	379.075.425.790	746.740.918.871	0	0	0	641.671.453.739	0	993.038.529.050	0
Comamonadaceae	0	0	0	0	433.293.614.751	0	0	768.109.004.686	0	0	0	0	0	0	0	0	0	0	0	478.403.258.770
Corynebacteriaceae	178.205.606.622	0	0	0	0	0	0	0	0	0	197.706.603.400	0	0	0	0	0	0	0	0	164.521.473.622
Corynebacteriales	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	23.638.749.256.6
Deinococcaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	478.403.258.770
Dermabacteraceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	407.698.071.260
Dermacoccaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	439.708.877.546
Desulfobulbaceae	0	0	0	0	0	0	0	0	130.473.073.619	224.236.464.552	0	0	0	0	0	0	986.515.296.225	0	0	0
Devosiaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	358.802.444.077
Dietziaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	239.201.629.385
Diploricetisidae	0	0	0	0	0	102.401.452.440	0	0	278.614.375.958	0	0	0	0	0	0	0	0	274.129.029.216	0	0
Enterobacteriaceae	122.611.549.855	0	0	36.212.407.100	1984.993.042.304	32.519.380.166.1	3.77E+08	463.648.406.452	231.480.978.113	130.386.303.883	120.036.152.064	0	0	0	210.733.871.464	0	267.528.979.978	0	113.040.358.239	0
Enterococcaceae	332.498.153.211	0	0	180.599.946.089	64.594.078.876	1282.960.121.553	0	446.338.123.810	268.027.029.671	650.635.473.797	286.815.793.933	0	0	0	501.940.969.868	0	672.174.855.907	0	221.613.274.283	0
Erwiniaceae	0	0	0	0	0	285.063.502.738	0	66.903.407.509.6	174.371.826.514	117.466.991.980	0	0	0	0	0	0	0	0	0	946.605.271.581
Erysipelotrichaceae	0	0	0	0	0	0	0	0	269.100.714.340	162.520.006.418	0	0	0	0	0	0	6.98E+08	0	0	0
Euzebyaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	545.239.008.157
Family_XI	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	837.205.702.848
Flavobacteriaceae	0	0	0	0	0	0	495.374.647.950	0	0	0	0	0	0	0	0	0	0	0	0	310.258.583.996
Fusibacteraceae	0	0	0	0	0	0	0	202.505.083.013	0	0	0	0	0	0	0	0	275.002.051.602	0	0	0
Fusobacteriaceae	0	0	0	0	0	0	0	0	724.139.775.434	0	525.547.445.255	0	0	0	0	0	37.016.149.168.6	0	12.663.615.673.3	0
Gemellaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	12.663.967.440.4
Hafniaceae	470.721.732.706	0	0	51.076.283.260.6	636.191.682.429	405.232.990.980	0	345.871.692.400	12.558.169.245.1	513.028.344.312	28.300.288.086.1	0	0	0	753.958.962.351	0	572.213.792.706	0	0	0
Halomonadaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	160.757.565.630
Helicobacteraceae	0	0	0	328.083.484.154	0	0	299.042.160.042	0	0	145.445.119.668	0	150.851.581.508	277.180.103.784	0	0	0	0	0	0	914.594.465.296
Hungateiclostridi	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5.63E+09
Hyphomicrobiaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	112.565.472.651
Iamiaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	6.68E+09
Intrasporangiaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	115.449.962.888
JG30_KF_CM45	0	0	0	448.996.880.896	0	0	0	0	35.472.366.890.1	0	0	0	0	0	0	0	0	0	0	0
JH49D030	0	0	0	0	0	0	200.618.447.846	0	0	4.11E+07	0	126.520.681.265	0	0	0	0	0	0	0	0
Lactobacillaceae	0	0	0	223.343.216.912	0	130.907.802.713	0	400.752.524.184	486.555.837.039	584.249.136.998	144.043.382.477	0	0	0	0	0	189.795.066.725	0	0	0
Leptotrichiaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	199.451.946.855
Listeriaceae	25.740.809.845.4	0	0	0	0	0	0	0	181.034.943.858	392.588.826.752	0	0	0	0	0	0	619.845.894.088	0	0	0
LWQ8	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	6.33E+08
Marinococcaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	299.002.036.731
Methanomethylo	0	0	0	0	0	0	0	0	0	0	0	0	0	0	121.541.501.976	0	0	0	0	0
Methanosarcinac	0	0	0	0	0	0	0	0	0	0	0	0	0	0	188.735.177.865	0	0	0	0	0
Methanospirillac	0	0	0	0	0	0	0	0	0	0	0	0	0	0	205.533.596.837	0	0	0	0	0
Methylophilaceae	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	8.09E+09
Microbacteriaceae	0	0	0	0	3.433.018.639.95	0	0	0	0	812.600.032.092	0	0	0	0	0	0	8.03E+09	0	238.146.328.079	0
Micrococcaceae	642.758.683.715	0	0	877.969.887.173	930.748.014.765	521.417.125.397	0	0	137.404.455.655	138.244.866.219	0	0	0	0	0	0	23.362.079.050.4	0	559.802.166.181	0
Microcystaceae	0	0	0	0	0	0	0	0	0	0	0	293.917.274.939	0	0	0	0	0	0	0	0

[illegible]

Supplementary Table 2. List of bacterial genera found in the fecal microbiota analysis of bats.

	Barbastella_barb	Eptesicus_bottae	Hypsugo_ariel	Miniopterus_schr	Myotis_bechstein	Myotis_blythii	Myotis_capaccini	Myotis_daubentc	Myotis_emargina	Myotis_myotis	Myotis_nattereri	Myotis_ricketti	Myotis_vivesi	Noctilio_leporinu	Pipistrellus_kuhl	Pipistrellus_pipis	Rhinolophus_bla	Rhinolophus_fer	Rhinolophus_hip	Rousettus_aegyptiacus
1174_901_12	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0 125.229.088.325
Achromobacter	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0 538.203.666.116
Acinetobacter	872.598.222.513	0	0	0	331.934.229.273	214.355.350.759	133.705.853.211	0	167.447.763.021	461.426.179.422	148.901.241.323	199.697.791.334	0	0	0	0	184.854.273.214	0	289.843.432.165	0 161.812.866.937
Actinobacillus	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0 338.399.952.159
Actinobaculum	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0 211.412.028.324
Actinomyces	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0 42.458.289.215.8
Actinotignum	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0 4.92E+09
Aerococcus	0	0	0	0	0	0	142.254.990.687	0	188.186.706.148	131.696.258.684	0	85.013.839.462.2	0	0	0	0	0	0	0	0
Aeromonas	0	0	0	0	0	0	0	0	380.546.464.723	0	0	0	0	0	0	0	0	0	0	0
Akkermansia	0	0	0	0	0	0	0	0	693.138.203.603	0	85.155.058.464.8	0	0	0	0	0	0	0	0	0
Alcanivorax	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Alkalibacterium	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Allorhizobium_Nit	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Alloscardovia	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Anaerococcus	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Anoxybacillus	761.562.421.463	0	0	0	0	0	0	0	36.791.308.123.6	115.523.033.933	623.336.227.149	114.387.391.967	0	0	0	0	0	0	0	0
Aquaspirillum	0	0	0	0	0	0	0	0	36.967.370.858.8	310.639.505.939	0	0	0	0	0	0	0	0	0	0
Arcobacter	0	0	0	0	0	0	0	0	0	250.980.263.076	0	0	0	0	0	0	0	0	0	0
Arsenophonus	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Asaia	0	0	0	192.537.255.959	0	0	196.093.219.699	0	0	0	0	0	0	0	0	0	0	0	0	0
Atopobium	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Bartonella	0	0	0	17.091.147.136.8	0	151.664.853.884	0	158.074.606.761	241.918.824.002	265.997.934.555	0	0	0	0	0	0	0	1	0	0
Bifidobacterium	0	0	0	0	49.995.417.086.1	506.472.048.554	0	0	6.39E+09	429.957.991.664	0	0	0	0	0	0	0	0	480.162.312.322	0
Bosea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Brachybacterium	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Brevibacterium	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Brevundimonas	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Brochothrix	25.740.809.845.4	0	0	0	0	0	0	0	0	181.034.943.858	392.588.826.752	0	0	0	0	0	0	0	619.845.894.088	0
Brucella	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Budvicia	0	0	0	0	0	0	0	0	7.88E+09	0	0	0	0	0	0	0	0	0	0	0
Buttiauxella	0	0	0	0	0	249.084.614.043	0	0	0	333.268.873.921	0	0	0	0	0	0	0	0	0	0
Campylobacter	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Candidatus_Amp	137.081.235.863	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Candidatus_Arth	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	224.366.753.212	0	0
Candidatus_Metf	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	121.541.501.976	0	0	0	0
Candidatus_Saci	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	576.416.337.285	0	0	0
Candidatus_Sole	0	0	0	0	0	0	0	0	0	92.368.965.673.3	0	0	0	0	0	0	0	0	0	0
Carnobacterium	531.951.351.392	0	0	940.660.017.713	479.206.072.776	767.900.189.027	0	339.749.084.391	146.620.475.574	946.257.308.257	105.617.691.916	0	0	0	0	0	764.187.565.469	0	886.239.944.964	0
Caulobacter	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Cetobacterium	0	0	0	0	0	0	0	0	0	0	0	525.547.445.255	0	0	0	0	0	0	0	0
Chishuiella	495.015.573.951	0	0	0	0	1.94E+09	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Christensenellae	124.896.237.120	0	0	0	0	0	0	0	0	189.263.804.943	0	0	0	0	0	0	0	9.87E+09	0	0
Chryseobacterium	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Chungangia	0	0	0	549.886.403.018	659.106.248.593	448.352.305.278	0	0	582.507.909.930	102.037.877.447	0	0	0	0	0	0	617.413.272.536	0	276.224.282.943	0
Clostridium_sens	0	660.886.797.126	0	0	0	0	0	629.572.365.940	100.188.131.046	0	27.834.122.618.2	0	350.656.934.306	710.036.704.214	0	0	0	0	417.304.700.527	0
Commensalibacter	0	0	0	0	0	0	0	0	551.034.720.753	248.714.296.587	491.674.449.797	0	0	0	0	0	0	0	0	0
Corynebacterium	178.205.606.622	0	0	0	0	0	0	0	0	0	0	197.706.603.400	0	0	0	0	0	0	0	0
Crossiella	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Deinococcus	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Desemzia	163.735.920.614	0	0	0	0	0	852.422.901.392	0	0	0	0	0	0	0	0	0	0	324.764.327.606	0	0
Desulfobulbus	0	0	0	0	0	0	0	0	130.473.073.619	224.236.464.552	0	0	0	0	0	0	0	986.515.296.225	0	0
Devosia	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Dietzia	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Diploricickettsia	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	146.056.645.184	0	0
Egibacter	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Ellin6055	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Enteractinococcus	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Enterobacter	0	0	0	221.032.769.840	518.285.823.799	166.886.691.409	0	595.562.778.996	163.227.251.475	738.540.282.332	0	0	0	0	0	0	0	15.967.579.440.8	0	138.103.764.259
Enterococcus	252.838.723.926	0	0	139.628.018.021	546.699.885.843	217.312.487.718	0	370.083.624.069	238.847.270.394	518.623.969.849	221.290.176.806	0	0	0	0	0	388.686.918.479	0	556.726.375.577	0
Erysipelothrix	0	0	0	0	0	0	0	0	0	216.096.028.182	162.520.006.418	0	0	0	0	0	0	0	0	0
Escherichia_Shig	0	0	0	0	0	0	0	0	0	491.992.215.106	818.771.677.905	0	0	0	0	0	0</			

Fusobacterium		0	0	0	0	0	0	0	0	724.139.775.434	0	0	0	0	0	0	0	37.016.149.168.1	0	12.863.615.673.1
Garicola		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	587.802.827.503
Gemella		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	12.663.967.440.4
Gemmobacter		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Geobacillus	668.651.806.045	0	0	155.570.102.814.574.114.039.546	437.558.638.669	0	604.468.390.644	705.913.692.061	425.843.561.121	732.926.622.606	0	0	0	0	411.608.848.357	0	670.131.983.524	0	0	0
Georgenia		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	306.037.378.772
Geothermophilum		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	4.22E+07
Glutamicibacter	642.758.683.715	0	0	877.969.887.173.802.426.444.242	454.994.561.652	0	0	921.466.082.437	138.244.866.219	0	0	0	0	0	0	213.017.462.193	0	0	0	0
Gordonia		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	182.918.893.059
Gracilibacillus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	228.648.616.324
Haemophilus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	138.024.839.232
Haemophilus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	399.959.195.016
Hafnia_Obesum	470.721.732.706	0	0	51.076.283.260.1636.191.682.429	405.232.990.980	0	345.871.692.400	12.558.169.245.1	513.028.344.312	28.300.288.086.1	0	0	0	0	753.958.962.351	0	572.213.792.706	0	0	0
Halomonas		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	110.806.637.141
Helicobacter		0	0	328.083.484.154	0	0	299.042.160.042	0	0	145.445.119.668	0	150.851.581.508	277.180.103.784	0	0	0	0	0	0	914.594.465.296
Hoyosella		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	275.081.873.792
Hyphomicrobium		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	112.565.472.651
Iamia		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	6.68E+09
Ignatzschineria	9.90E+09	0	0	373.229.619.931	599.945.005.041	580.920.672.085	0	191.081.029.933	146.496.797.973	374.413.179.343	0	0	0	0	338.246.349.128	0	363.552.712.218	0	0	0
Isobaculum		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	8.82E+09	0	0	0
Jeotgaliococcus		0	0	0	0	716.533.406.398	0	0	22.153.240.624.1	0	314.212.280.404	0	0	0	0	0	74.556.111.767.8	0	0	0
Klebsiella	73.109.992.460.1	0	0	130.155.185.028	387.464.482.422	808.141.192.229	0	0	392.778.315.375	20.572.152.711.2	0	0	0	0	0	0	476.670.222.777	0	0	0
Koukoulia		0	0	308.059.609.534	289.140.162.151	869.028.542.329	0	161.636.851.421	36.624.879.040.1	397.659.711.907	0	0	0	0	0	0	115.448.480.329	0	0	0
Kurtzia		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	207.779.327.877
Lactococcus	146.372.297.405	0	0	246.293.657.822	808.259.242.902	275.902.724.155	0	973.494.673.331	276.929.098.757	245.652.075.524	0	0	0	0	0	0	24.387.007.331.1	0	0	0
Lautropia		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	524.132.982.035
Leucobacter		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	8.03E+09	0	0	0
Leuconostoc		0	0	223.343.216.912	0	130.907.802.713	0	0	1.90E+09	121.375.700.996	508.388.408.744	0	0	0	0	0	1.75E+09	0	0	0
Lonsdalea	227.097.914.080	0	0	377.835.111.093	299.222.571.264	102.622.860.985	0	174.666.874.464	16.783.594.098.1	360.465.259.805	262.949.782.522	0	0	0	485.550.557.643	0	203.239.611.470	0	0	0
Luteipulveratus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	270.860.668.568
Lysinibacillus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	6.88E+09	0	0	0
Marinilactibacillus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2.18E+09	0	0	0
Mariniluteococcus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	137.189.169.794
Mariniluteococcus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	158.295.195.916
Marixanthomona		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	183.622.427.263
Methanimicrococcus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Methanospirillum		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Methylobacillus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	8.09E+09
Methylobacterium		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Methylocella		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	830.170.360.807
Microbacterium		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	204.024.919.181
Microcystis_PCC		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	111.861.938.447
Mobiluncus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Moellerella	170.894.607.376	0	0	215.102.622.357	0	709.060.867.976	0	117.999.354.343	421.727.028.584	578.427.217.780	831.779.924.306	0	0	0	214.283.073.510	0	116.242.930.701	0	0	0
Moheibacter		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	204.024.919.181
Monoglobus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Moraxella		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	29.548.436.571.1
Morgarella		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Mycobacterium		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	355.284.773.057
Mycoplasma		0	0	0	0	2.49E+09	0	0	0	9.26E+08	0	0	0	0	0	0	0	0	0	0
Myroides		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Natribacillus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Nesterenkonia		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Nissabacter		0	0	508.298.355.731	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Nitratireductor		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Nitrococcus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	200.507.248.161
Nitrosococcus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	225.130.945.303
Nitrosococcus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	513.579.968.974
Nocardiodides		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	102.012.459.590
Nocardiopeis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	12.065.611.599.1
Not_Assigned	643.672.558.621	11.716.621.253.4	0	657.861.296.160	219.329.894.759	963.902.104.211	153.103.552.304	788.814.551.769	583.078.729.627	741.461.528.017	189.346.438.456	29.294.403.892.1	592.330.084.799	883.620.689.655	138.998.682.476	480.004.929.447	0	892.359.831.890	1	163.642.055.867
Oceanobacillus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
OLB8		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	734.841.476.155
Olivibacter		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Orbus		0	0	146.328.314.528	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Others		0	0	0	0	0	3.77E+08	0	0	1.44E+09	0	0	0	0	0	0	0	0	0	0
Paenacaligenes		0	0	0	0	0	0	0	159.014.058.473	423.786.345.850	0	0	0	0	0	0	0	0	0	0
Paenibacillus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Paeniciostidium		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Paenisporosarcin		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Paludicola		0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Pantoea		0	0	0	0	0	213.936.007.395	0	376.262.092.150	899.720.570.167	726.196.990.705	0	0	0	0	0	0	0	0	0

Paraclostridium	0	470.646.519.692	0	0	0	0	424.994.343.464	0	0	41.144.305.422.1	0	522.433.090.024	221.490.950.512	0	0	0	0	0	0
Paracoccus	0	0	0	0	0	0	0	0	0	0	0	0	0	0	691.699.604.743	0	0	0	265.232.394.935
Parvimonas	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	831.225.662.113
Pasteurella	0	0	0	0	0	0	0	0	911.952.420.819	169.308.816.813	0	0	0	0	0	0	334.454.876.091	0	0
Peptoanaerobac	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	260.160.671.039	0	0
Peptococcus	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5.63E+09
Peptostreptococ	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	12.663.615.673.1
Porphyromonas	0	0	0	0	0	0	0	0	169.886.814.608	281.838.492.143	0	0	0	0	0	0	910.562.348.639	0	0
Providencia	868.181.160.468	0	0	196.388.001.078	648.273.908.225	50.370.444.173.1	0	0	800.506.670.435	604.204.125.128	0	0	0	0	100.807.196.993	0	945.483.244.081	0	232.166.287.344
Pseudocitrobacte	0	0	0	0	478.289.490.130	0	0	0	815.456.710.121	0	0	0	0	0	0	0	0	0	0
Pseudogracillibac	0	0	0	0	0	0	0	0	456.701.790.188	0	0	0	0	0	0	0	264.525.782.970	0	0
Pseudomonas	321.067.101.264	0	0	437.059.571.026	172.242.544.433	244.257.907.744	0	111.937.972.414	208.504.126.210	424.923.985.895	13.377.676.100.0	0	0	0	165.247.396.635	0	202.563.019.120	0	274.730.106.690
Pseudonocardia	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	267.342.997.548
Psychrobacter	0	0	0	0	0	251.852.220.866	0	0	110.304.110.989	191.526.741.741	0	0	0	0	0	0	290.245.022.462	0	0
Rhodobacter	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	140.706.840.814
Rhodococcus	700.637.427.746	0	0	0	0	440.603.006.174	0	712.448.931.883	114.163.939.417	968.948.392.697	0	0	0	0	0	0	132.699.402.678	0	119.249.047.590
Rhodohalobacter	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	348.249.431.016
Rhodopseudomon	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	151.259.853.875
Rickettsia	0	0	0	94.497.285.224.0	0	0	678.784.222.037	0	181.697.345.960	0	0	0	0	0	205.862.977.602	0	0	0	0
Rickettsiella	0	0	0	0	0	102.401.452.440	0	0	278.614.375.958	0	0	0	0	0	0	0	128.072.384.032	0	0
Romboutsia	0	149.120.634.134	0	0	0	0	0	0	0	148.279.952.550	194.647.201.946	814.706.999.114	116.379.310.344	0	0	0	103.016.641.552	0	0
Roseomonas	0	0	0	0	0	0	795.685.949.166	0	0	0	0	0	0	0	0	0	0	0	0
Rothia	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	221.507.744.152
Rubellimicrobium	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	7.74E+09
Rubribacterium	0	0	0	0	0	0	701.410.362.772	0	0	0	0	0	0	0	0	0	0	0	0
Rubritepida	0	0	0	0	0	0	603.363.752.922	0	0	0	0	0	0	0	0	0	0	0	0
Saccharopolyspc	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	236.739.259.670
Salinisphaera	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	659.915.083.421
Sarcina	0	681.198.910.081	0	0	0	0	878.648.465.193	0	0	335.326.089.192	0	284.184.914.841	367.042.146.563	0	0	0	0	0	0
Savagea	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	563.099.438.995	0	0
Selenomonas	0	0	0	0	0	0	0	0	146.782.207.821	0	0	0	0	0	0	0	100.397.574.394	0	0
Serratia	122.514.612.402	0	0	31.499.095.074	1.685.770.471.040	154.432.460.706	0	127.628.546.938	196.389.157.687	45.073.586.590.1	0	0	0	0	258.795.982.500	0	100.493.606.857	0	353.174.170.445
Shinella	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	685.945.848.972
Skermanella	0	0	0	0	0	0	143.298.891.319	150.282.196.569	0	0	0	0	0	0	0	0	0	0	0
Sneathia	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	199.451.946.855
Solibacillus	0	0	0	0	0	0	0	0	360.160.046.970	0	0	0	0	0	0	0	0	0	0
Sphingobacteriur	134.415.767.388	0	0	0	173.650.748.681	807.034.149.500	0	6.12E+09	0	252.893.473.278	0	0	0	0	0	0	163.429.790.666	0	0
Sphingomonas	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	545.239.008.157
Sporosarcina	935.960.215.979	0	0	0	0	444.173.218.975	0	58.443.076.443.1	190.123.731.964	269.207.190.378	162.401.852.793	0	0	0	0	0	221.494.509.562	0	0
Stackebrandtia	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	144.224.511.835
Staphylococcus	248.931.908.703	0	0	310.678.116.215	104.723.733.657	134.104.388.594	0	25.741.670.470.1	464.851.097.604	276.242.866.606	221.290.176.806	0	0	0	103.888.101.546	0	257.934.463.955	0	278.916.135.205
Stenotrophomon	784.409.294.107	0	0	0	677.437.901.525	112.364.837.001	0	0	0	368.241.533.530	155.340.902.671	0	0	0	0	0	355.320.111.118	0	1.688.482.089.71
Streptococcus	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	394.468.110.553
Streptomyces	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	207.542.590.201
Tabrizicola	0	0	0	0	0	641.073.987.480	0	0	0	0	0	0	0	0	0	0	0	0	0
Taibaiella	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5.28E+09
Terribacillus	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5.98E+09
Tetragenococcus	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	545.239.008.157
Thermus	506.439.010.273	0	0	901.074.357.888	367.466.315.587	45.914.597.188.0	0	342.309.447.740	264.479.792.982	126.313.017.646	152.516.522.623	0	0	0	123.236.182.143	0	209.088.861.456	0	0
TM7a	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	256.086.450.282
TM7x	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	823.135.018.766
Tomitella	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	23.638.749.256.0
Treponema	132.664.173.819	0	0	0	0	0	0	0	777.402.063.649	120.347.093.360	0	0	0	0	0	0	262.168.622.527	0	0
Truepera	883.412.408.898	0	0	106.280.565.289	0	0	0	0	0	0	0	0	0	0	0	0	0	0	471.367.916.729
Ureaplasma	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	12.909.852.644.1
Vagococcus	251.315.599.083	0	0	437.444.645.538	0	142.531.751.369	0	0	201.689.626.303	432.838.093.043	231.457.944.981	0	0	0	0	0	389.891.797.605	0	0
Varibaculum	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	921.629.807.337
Veillonella	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	274.378.339.588
Virgibacillus	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	14.308.836.907.1	0	140.706.840.814
Viridibacillus	0	0	0	0	0	0	207.055.470.828	241.918.824.002	0	0	0	0	0	0	0	0	481.035.334.708	0	0
Weissella	0	0	0	0	0	0	0	400.752.524.184	467.528.513.802	46.287.343.600.1	932.045.416.031	0	0	0	0	0	188.049.021.953	0	0
Wenzhouxiangeli	3.81E+09	0	0	0	0	1.38E+09	0	0	0	0	0	0	0	0	0	0	0	0	0
Wohlfahrtiimonas	0	0	0	0	0	0	0	0	646.928.990.029	159.022.740.457	141.219.002.428	0	0	0	0	0	0	0	0
Yersinia	319.094.654.593	0	0	122.453.694.789	455.791.552.441	857.958.115.038	0	209.838.474.468	512.378.632.859	285.952.922.685	114.528.610.969	0	0	0	147.883.418.571	0	583.178.953.874	0	0
Yokenella	258.931.223.297	0	0	469.790.904.540	123.322.028.814	251.852.220.866	0	0	172.605.003.642	9.46E+09	33.892.560.582.0	0	0	0	0	0	269.763.917.286	0	0
ZOR006	0	0	0	0	0	0	0	0	5.30E+09	0	0	0	0	0	0	0	6.98E+08	0	0

Supplementary Table 3. Biological information on the bat species studied.

<i>Rousettus_aeg</i>	Pteropodidae	Rousettus	F	Caves	Temperate_clim	Frugivores
<i>Eptesicus_botta</i>	Vespertilionidae	Eptesicus	Not_Available	Desert	Semiarid	Insectivores
<i>Hypsugo_ariel</i>	Vespertilionidae	Hypsugo	Not_Available	Desert	Semiarid	Insectivores
<i>Myotis_ricketti</i>	Vespertilionidae	Myotis	Not_Available	Fresh_water	Temperate_clim	Piscivores
SampleType	Family	Genus	Sex	Habitat	Climate	Diet
<i>Barbastella_ba</i>	Vespertilionidae	Barbastella	M	Forests	Temperate_clim	Insectivores
<i>Miniopterus_sc</i>	Miniopteridae	Miniopterus	M	Caves	Temperate_clim	Insectivores
<i>Miniopterus_sc</i>	Miniopteridae	Miniopterus	Not_Available	Forests	Temperate_clim	Insectivores
<i>Myotis_bechste</i>	Vespertilionidae	Myotis	M	Forests	Temperate_clim	Insectivores
<i>Myotis_blythii</i>	Vespertilionidae	Myotis	M	Forests	Temperate_clim	Insectivores
<i>Myotis_capacci</i>	Vespertilionidae	Myotis	Not_Available	Fresh_water	Temperate_clim	Piscivores
<i>Myotis_dauben</i>	Vespertilionidae	Myotis	M	Lakes	Temperate_clim	Insectivores
<i>Myotis_dauben</i>	Vespertilionidae	Myotis	Not_Available	Lakes	Temperate_clim	Insectivores
<i>Myotis_emargin</i>	Vespertilionidae	Myotis	M	Forests	Temperate_clim	Insectivores
<i>Myotis_emargin</i>	Vespertilionidae	Myotis	Not_Available	Forests	Temperate_clim	Insectivores
<i>Myotis_myotis</i>	Vespertilionidae	Myotis	M	Forests	Temperate_clim	Insectivores
<i>Myotis_myotis</i>	Vespertilionidae	Myotis	Not_Available	Forests	Temperate_clim	Insectivores
<i>Myotis_natterer</i>	Vespertilionidae	Myotis	F	Meadow	Temperate_clim	Insectivores
<i>Pipistrellus_kuh</i>	Vespertilionidae	Pipistrellus	Not_Available	Forests	Temperate_clim	Insectivores
<i>Pipistrellus_pip</i>	Vespertilionidae	Pipistrellus	F	Forests	Temperate_clim	Insectivores
<i>Rhinolophus_b</i>	Rhinolophidae	Rhinolophus	Not_Available	Forests	Temperate_clim	Insectivores
<i>Rhinolophus_fe</i>	Rhinolophidae	Rhinolophus	M	Caves	Temperate_clim	Insectivores
<i>Rhinolophus_fe</i>	Rhinolophidae	Rhinolophus	Not_Available	Caves	Temperate_clim	Insectivores
<i>Rhinolophus_hi</i>	Rhinolophidae	Rhinolophus	Not_Available	Caves	Temperate_clim	Insectivores
<i>Myotis_vivesi</i>	Vespertilionidae	Myotis	Not_Available	Ocean_coasts	Tropical	Piscivores
<i>Noctilio_leporin</i>	Noctilionidae	Noctilio	Not_Available	Fresh_water	Tropical	Piscivores


Africa	South_Africa	Dietrich and Markotter, 2019
Asia	Israel	Aizpurua et al., 2021
Asia	Israel	Aizpurua et al., 2021
Asia	China	Aizpurua et al., 2021
Continent	Capture_locatio	Source
Europe	Slovenia	Vengust et al., 2018
Europe	Slovenia	Vengust et al., 2018
Europe	Not_specified	Aizpurua et al., 2021
Europe	Slovenia	Vengust et al., 2018
Europe	Slovenia	Vengust et al., 2018
Europe	Not_specified	Aizpurua et al., 2021
Europe	Slovenia	Vengust et al., 2018
Europe	Not_specified	Aizpurua et al., 2021
Europe	Slovenia	Vengust et al., 2018
Europe	Not_specified	Aizpurua et al., 2021
Europe	Slovenia	Vengust et al., 2018
Europe	Not_specified	Aizpurua et al., 2021
Europe	Slovenia	Vengust et al., 2018
Europe	Not_specified	Aizpurua et al., 2021
Europe	Slovenia	Vengust et al., 2018
Europe	Not_specified	Aizpurua et al., 2021
Europe	Not_specified	Aizpurua et al., 2021
South_America	Mexico	Aizpurua et al., 2021
South_America	Costa_Rica	Aizpurua et al., 2021

Supplementary Table 4. GenBank accession numbers of selected pathogenic species used in Maximum Likelihood phylogenetic trees of the Supplementary Figure 1-4.

Genera	Specie	Access Number
Clostridium	botulinum	X73844.1
Clostridium	botulinum	ON567373.1
Clostridium	botulinum	MF062277.1
Clostridium	botulinum	JN617091.2
Clostridium	butyricum	AB687551.1
Clostridium	butyricum	OP984128.1
Clostridium	butyricum	MK156151.1
Clostridium	butyricum	KX148520.1
Clostridium	cadaveris	MK968718.1
Clostridium	cadaveris	MN537449.1
Clostridium	cadaveris	NR104695.1
Clostridium	cadaveris	MF145697.1
Clostridium	chauvoei	NR026013.1
Clostridium	chauvoei	EU081297.1
Clostridium	chauvoei	EU106373.1
Clostridium	chauvoei	EU106368.1
Clostridium	difficile	AB638443.1
Clostridium	difficile	JX073558.1
Clostridium	difficile	JX073560.1
Clostridium	difficile	JF693896.1
Clostridium	paraputrificum	AB032556.1
Clostridium	paraputrificum	OR398739.1
Clostridium	paraputrificum	OP727735.1
Clostridium	paraputrificum	MT903094.1
Clostridium	tertium	NR114728.1
Clostridium	tertium	OR398745.1
Clostridium	tertium	OP727740.1
Clostridium	tertium	ON142170.1
Pseudomonas	aeruginosa	LN874213.1
Pseudomonas	aeruginosa	AM419153.2
Pseudomonas	aeruginosa	KJ995745.1
Pseudomonas	aeruginosa	EU344794.1
Pseudomonas	oryzihabitans	OR648088.1
Pseudomonas	oryzihabitans	OQ381240.1
Pseudomonas	oryzihabitans	MW187500.1
Pseudomonas	oryzihabitans	MN911431.1
Pseudomonas	plecoglossicida	GQ398824.1
Pseudomonas	plecoglossicida	NR024662.1
Pseudomonas	plecoglossicida	MF574326.1
Pseudomonas	plecoglossicida	JQ034609.1
Pseudomonas	syringae	MK072691.1
Pseudomonas	syringae	NR074597.1
Pseudomonas	syringae	KY460679.1


Pseudomonas	syringae	MF074184.1
Pseudomonas	tolaasii	OQ790026.1
Pseudomonas	tolaasii	MW326080.1
Pseudomonas	tolaasii	MK355664.1
Pseudomonas	tolaasii	HQ591342.1
Staphylococcus	aureus	ON564600.1
Staphylococcus	aureus	ON222796.1
Staphylococcus	aureus	MT860724.1
Staphylococcus	epidermidis	KJ623584.1
Staphylococcus	epidermidis	OR449114.1
Staphylococcus	epidermidis	OR098502.1
Staphylococcus	saprophyticus	MH371235.1
Staphylococcus	saprophyticus	OQ062584.1
Staphylococcus	saprophyticus	OQ719862.1
Staphylococcus	haemolyticus	MF578766.1
Staphylococcus	haemolyticus	KJ623587.1
Staphylococcus	haemolyticus	OQ787674.1
Staphylococcus	carnosus	OP263296.1
Staphylococcus	carnosus	MF678927.1
Staphylococcus	carnosus	NR116434.1
Staphylococcus	equorum	MK214690.1
Staphylococcus	equorum	MZ603727.1
Staphylococcus	equorum	KF439745.1
Staphylococcus	xylosus	MW585598.1
Staphylococcus	xylosus	MN294563.1
Staphylococcus	xylosus	KY672997.1
Streptococcus	pyogenes	OQ913600
Streptococcus	pyogenes	OR397759
Streptococcus	pyogenes	MT378375
Streptococcus	mutans	LC720109
Streptococcus	mutans	LC720109
Streptococcus	mutans	OR563799
Streptococcus	pneumoniae	MF805759
Streptococcus	pneumoniae	NR 115239
Streptococcus	pneumoniae	MZ007495
Streptococcus	bovis	AF104109
Streptococcus	bovis	AF429766
Streptococcus	bovis	AY500370
Streptococcus	canis	NR 115729
Streptococcus	canis	AJ413203
Streptococcus	canis	JX876619
Streptococcus	agalactiae	OP290417
Streptococcus	agalactiae	OP290418

Tree scale: 0.1




Supplementary Figure 1. Phylogenetic diversity of *Pseudomonas* ASVs. Phylogenetic tree of *Pseudomonas* ASVs recovered from bat fecal samples based on 16S rRNA sequences. Several ASVs clustered near known pathogenic species, while others formed distinct lineages, indicating high genetic diversity and the presence of ancestral lineages related to clinically relevant taxa.


Tree scale: 0.1



Supplementary Figure 2. Phylogenetic relationships among *Staphylococcus* ASVs. Phylogenetic analysis of *Staphylococcus* ASVs identified in bat fecal samples. Although numerous ASVs were detected, most did not cluster directly with established pathogenic species, suggesting predominance of non-pathogenic lineages with phylogenetic proximity to clinically relevant taxa.



Supplementary Figure 3. Phylogenetic structure of *Clostridium* ASVs. Phylogenetic tree based on *Clostridium* ASVs detected in fecal samples. A small cluster of ASVs formed potential ancestral lineages related to pathogenic species, while the majority grouped into a well-defined clade distant from known pathogens, indicating evolutionary relatedness without direct association with pathogenic strains.



Supplementary Figure 4. Maximum Likelihood phylogenetic tree based on *Streptococcus* ASV obtained from 16S rRNA sequencing data and selected pathogenic species (see Supplementary Table 1).